



US008758773B2

(12) **United States Patent**
Talaat(10) **Patent No.:** **US 8,758,773 B2**
(45) **Date of Patent:** **Jun. 24, 2014**(54) **VACCINE CANDIDATES AGAINST JOHNE'S DISEASE**(75) Inventor: **Adel Mohamed Talaat**, Madison, WI (US)(73) Assignee: **Wisconsin Alumni Research Foundation**, Madison, WI (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 307 days.

(21) Appl. No.: **11/636,025**(22) Filed: **Dec. 8, 2006**(65) **Prior Publication Data**

US 2007/0134274 A1 Jun. 14, 2007

Related U.S. Application Data

(60) Provisional application No. 60/749,128, filed on Dec. 9, 2005.

(51) **Int. Cl.**
A61K 39/04 (2006.01)
C07H 21/02 (2006.01)
C07H 21/04 (2006.01)(52) **U.S. Cl.**
USPC **424/248.1**; 424/9.1; 424/9.2; 424/184.1;
424/234.1; 424/282.1; 435/253.1; 536/23.1;
536/23.7(58) **Field of Classification Search**
USPC 424/9.2, 184.1, 248.1, 278.1, 282.1,
424/9.1, 234.1; 435/243, 253.1, 69.1
See application file for complete search history.(56) **References Cited****U.S. PATENT DOCUMENTS**4,816,567 A 3/1989 Cabilly et al.
5,830,475 A * 11/1998 Aldovini et al. 424/200.1
6,503,514 B1 1/2003 Flesselles et al.
2003/0175725 A1 9/2003 Kapur et al.
2003/0204070 A1 10/2003 Chen et al.
2004/0213817 A1 10/2004 Miller et al.**OTHER PUBLICATIONS**AE017242 (AE017242 Jan. 30, 2004 NCBI Website).
Fleishmann et al., "Whole-Genome Comparison of *Mycobacterium tuberculosis* Clinical and Laboratory Strains," J of Bacteriology, 2002, 184:5479-5490.
<http://www.tigr.org/tigr-scripts/CMR2>, last visited Jul. 2008.
<http://www.ncbi.nlm.nih.gov/COG/>, printed on Oct. 14, 2008.
<http://www.tigr.org>, printed on Oct. 14, 2008.
<http://www.pasteur.fr>, printed on Oct. 14, 2008.
Ausubel, *Current Protocols in Molecular Biology* vols. 1-3, John Wiley & Sons, Inc., New York, NY (1987-2004).
Altschul et al., "Gapped BLAST and PSI-BLAST: a new generation of protein database search programs," *Nucl. Acids Res.*, 25(17):3389-3402 (1997).
Bardarov et al., "Conditionally replicating mycobacteriophages: A system for transposon delivery to *Mycobacterium tuberculosis*," *Proc. Natl. Acad. Sci. USA*, 94:10961-10966 (1997).Cavaignac et al., "Construction and screening of *Mycobacterium paratuberculosis* insertional mutant libraries," *Archives of Microbiology*, 173:229-231 (2000).
Davis, "Plasmid DNA expression systems for the purpose of immunization," *Current Opinion Biotech.*, 8:635-640 (1997).
Dieffenbach et al., *PCR Primer: A Laboratory Manual*, Cold Spring Harbor Laboratory Press (1995).
Furth et al., "Gene Transfer into Somatic Tissues by Jet Injection," *Analytical Biochem.*, 205:355-368 (1992).
GenBank access No. AE016958, 2008.
Glasner, et al., "ASAP, a systematic annotation package for community analysis of genomes," *Nucleic Acids Res.*, 31(1):147-151 (2003).
Harris et al., Development of a transposon mutagenesis system for *Mycobacterium avium* subsp. *paratuberculosis*, *FEMS Microbiology Letters*, 175:21-26 (1999).
Karin and Altschul, Methods for assessing the statistical significance of molecular sequence features by using general scoring schemes, *Proc. Natl. Acad. Sci. USA*, 87:2264-2268 (1990).
Kriegler, *Gene Transfer and Expression: A Laboratory Manual*, Stockton Press, New York (1990).
Li et al., The complete genome sequence of *Mycobacterium avium* subspecies *paratuberculosis*, *Proc. Natl. Acad. Sci. USA*, 102:12344-12349 (2005).
McAdam et al., "Characterization of a *Mycobacterium tuberculosis* H37Rv transposon library reveals insertions in 351 ORFs and mutants with altered virulence," *Microbiology*, 148:2975-2986 (2002).
Pearson and Lipman, "Improved tool for biological sequence comparison," *Proc. Natl. Acad. Sci. USA*, 85:2444-2448 (1988).
Riechmann et al., "Reshaping human antibodies for therapy," *Nature*, 332:323-327 (1988).
Sambrook et al., *Molecular Cloning: A laboratory Manual*, third edition, Cold Spring Harbor Laboratory Press (2000).
Smith and Waterman, Comparison of Biosequences, *Adv. Appl. Math.*, 2:482-489 (1981).
Talaat et al., "A combination vaccine confers full protection against co-infections with influenza, herpes simplex and respiratory syncytial viruses," *Vaccine*, 20:538-544 (2002).
Tang et al., Genetic immunization is a simple method for eliciting an immune response, *Nature*, 356:152-154 (1992).
Valentin-Weigand and Goethe, "Pathogenesis of *Mycobacterium avium* subspecies *paratuberculosis* infections in ruminants: still more questions than answers," *Microbes & Infection*, 1:1121-1127 (1999).
Verhoeven et al., "Reshaping Antibodies: Grafting an Antilysozyme Activity", *Science*, 239:1534-1536 (1988).
Wolf et al., Direct Gene Transfer into Mouse Muscle in Vivo, *Science* 247:1465-1468 (1990).
International Search Report for related PCT Application No. PCT/US06/47089.

* cited by examiner

Primary Examiner — Rod P Swartz

(74) Attorney, Agent, or Firm — Quarles & Brady LLP

(57) **ABSTRACT**A composition and method for immunizing a mammal infected with *Mycobacterium* are disclosed. The genes gcpE, pstA, kdpC, papA2, impA, umaA1, fabG2_2, aceAB, mbtH2, lpqP, map0834c, cspB, lipN, and map1634 of *M. paratuberculosis* and the products that they encode are vaccine targets for Johne's and Crohn's disease. Eighteen *M. paratuberculosis*-specific genomic islands (MAPs) were identified. Three inverted large genomic fragments in *M. paratuberculosis* (INV) were also identified. These genomic identifiers represent novel virulence determinants that can be used as targets for vaccines and for developments of drugs against Johne's disease. The methods can be used to deliver an immunizing compound to a mammal, to provide an immune response against Johne's or Crohn's disease in the mammal.**9 Claims, 7 Drawing Sheets**

FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5





FIGURE 6



M. avium 104 vs.
M. paratuberculosis k10
(inner circle)

1

VACCINE CANDIDATES AGAINST JOHNE'S DISEASE

CROSS-REFERENCE TO RELATED APPLICATIONS

This invention claims priority to U.S. Provisional Patent Application Ser. No. 60/749,128 filed Dec. 9, 2005.

GOVERNMENT INTERESTS

This invention was made with United States government support awarded by the following agency: USDA/CSREES grants 2004-35204-14209, 2004-35605-14243, 04-CRHF-0-6055. The United States may have certain rights in this invention.

FIELD OF THE INVENTION

This invention relates to nucleic acid sequences from *Mycobacterium avium* subspecies paratuberculosis (hereinafter referred to as *Mycobacterium paratuberculosis* or *M. paratuberculosis*), the products encoded by those sequences, compositions containing those sequences and products, and compositions and methods for prevention and treatment of *M. paratuberculosis* infection.

BACKGROUND OF THE INVENTION

Mycobacterium paratuberculosis causes Johne's disease (paratuberculosis) in dairy cattle. The disease is characterized by chronic diarrhea, weight loss, and malnutrition, resulting in estimated losses of \$220 million per year in the USA alone. World-wide, the prevalence of the disease can range from as low as 3-4% of the examined herds in regions with low incidence (such as England), to high levels of 50% of the herds in some areas within the USA (Wisconsin and Alabama). Cows infected with Johne's disease are known to secrete *Mycobacterium paratuberculosis* in their milk. In humans, *M. paratuberculosis* bacilli have been found in tissues examined from Crohn's disease patients indicating possible zoonotic transmission from infected dairy products to humans.

Unfortunately, the virulence mechanisms controlling *M. paratuberculosis* persistence inside the host are poorly understood, and the key steps for establishing the presence of paratuberculosis are elusive. Mechanisms responsible for invasion and persistence of *M. paratuberculosis* inside the intestine remain undefined on a molecular level (Valentin-Weigand and Goethe, 1999, *Microbes & Infection* 1: 1121-1127). Both live and dead bacilli are observed in sub-epithelial macrophages after uptake. Once inside the macrophages, *M. paratuberculosis* survive and proliferate inside the phagosomes using unknown mechanisms.

M. paratuberculosis is closely related to *Mycobacterium avium* subspecies *avium* (hereinafter referred to as *Mycobacterium avium* or *M. avium*), which is a persistent health problem for immunocompromised humans, particularly HIV-positive individuals. Limited tools are available to researchers to definitively identify *M. paratuberculosis* and to distinguish it from *M. avium*. Existing methods are subject to high cross-reactivity, poor sensitivity, specificity, and predictive value. This dearth of knowledge translates into a lack of suitable vaccines for prevention and treatment of Johne's disease in animals, and of Crohn's disease in humans.

The current challenge in screening *M. paratuberculosis* is to identify those targets that are essential for survival of the

2

bacilli during infection. Recently, random transposon mutagenesis-based protocols were employed for functional analysis of a large number of genes in *M. paratuberculosis* (Harris et al., 1999, *FEMS Microbiology Letters* 175: 21-26; Cavaignac et al., 2000, *Archives of Microbiology* 173: 229-231). When *M. paratuberculosis* was used as a target for mutagenesis, the libraries were screened to identify auxotrophs or genes responsible for survival under in vitro conditions. In these reports, six auxotrophs and two genes responsible for cell wall biosynthesis were identified (Harris et al., 1999; Cavaignac et al., 2000). So far, none of these libraries have been screened for virulence determinants.

Many clinical methods for detecting and identifying *Mycobacterium* species in samples require analysis of the bacterium's physical characteristics (e.g., acid-fast staining and microscopic detection of bacilli), physiological characteristics (e.g., growth on defined media) or biochemical characteristics (e.g., membrane lipid composition). These methods require relatively high concentrations of bacteria in the sample to be detected, may be subjective depending on the clinical technician's experience and expertise, and are time-consuming. Because *Mycobacterium* species are often difficult to grow in vitro and may take weeks to reach a useful density in culture, these methods can also result in delayed patient treatment and costs associated with isolating an infected individual until the diagnosis is completed.

More recently, assays that detect the presence of nucleic acid derived from bacteria in the sample have been preferred because of the sensitivity and relative speed of the assays. In particular, assays that use in vitro nucleic acid amplification of nucleic acids present in a clinical sample can provide increased sensitivity and specificity of detection. Such assays, however, can be limited to detecting one or a few *Mycobacterium* species depending on the sequences amplified and/or detected.

The genome sequences of both *M. avium* (Institute for Genomic Research) and of *M. paratuberculosis* (Li et al., 2005, *Proc. Natl. Acad. Sci. USA* 102: 12344-12349; GenBank accession No. AE016958) are currently available. It would be useful to analyze these genomes to provide a higher resolution analysis of *M. avium* subspecies genomes. A better understanding of the virulence mechanisms and pathogenesis of *M. paratuberculosis* is required to develop more effective vaccine and chemotherapies directed against *M. paratuberculosis*.

In view of the problems with bacterial specificity, the present inventors have focused their attention on identification of putative virulence factors that may contribute to the pathogenicity of *M. paratuberculosis*. This information could be used to design vaccines against pathogenic subspecies of *M. avium*. Such vaccines can be used for prevention and treatment of Johne's disease in animals or Crohn's disease in humans.

SUMMARY OF THE INVENTION

This invention relates to immunogenic compositions and methods for prevention and treatment of Johne's disease in animals or Crohn's disease in humans.

This invention provides a vaccine composition that includes an antigen selected from *Mycobacterium* strain-specific polynucleotide sequences and their products. In one embodiment, the antigen includes at least one of the gcpE (SEQ ID NO:7), pstA (SEQ ID NO:8), kdpC (SEQ ID NO:9), papA2 (SEQ ID NO:10), impA (SEQ ID NO:11), umaA1 (SEQ ID NO:12), fabG2_2 (SEQ ID NO:13), aceAB (SEQ ID NO:14), mbtH2 (SEQ ID NO:15), lpqP (SEQ ID NO:16),

3

map0834c (SEQ ID NO:17), cspB (SEQ ID NO:18), lipN (SEQ ID NO:19), or map1634 (SEQ ID NO:20) genes of *M. paratuberculosis*, or homologs of these genes. In another aspect, the invention is directed to an antigen that includes at least one of the genomic islands MAP-1 (SEQ ID NO:21), MAP-2 (SEQ ID NO:22), MAP-3 (SEQ ID NO:23), MAP-4 (SEQ ID NO:24), MAP-5 (SEQ ID NO:25), MAP-6 (SEQ ID NO:26), MAP-7 (SEQ ID NO:27), MAP-8 (SEQ ID NO:28), MAP-9 (SEQ ID NO:29), MAP-10 (SEQ ID NO:30), MAP-11 (SEQ ID NO:31), MAP-12 (SEQ ID NO:32), MAP-13 (SEQ ID NO:33), MAP-14 (SEQ ID NO:34), MAP-15 (SEQ ID NO:35), MAP-16 (SEQ ID NO:36), MAP-17 (SEQ ID NO:37), or MAP-18 (SEQ ID NO:38) of *M. paratuberculosis*, or homologs of these genomic islands. In addition to the antigens, the vaccine composition includes a pharmaceutically acceptable carrier. The vaccine composition may optionally include an adjuvant.

This invention provides an immunological composition that includes a eukaryotic expression vector that encodes an antigen. In one aspect, the eukaryotic expression vector includes at least one of the gcpE, pstA, kdpC, papA2, impA, umaA1, fabG2_2, aceAB, mbtH2, lpqP, map0834c, cspB, lipN, or map1634 genes of *M. paratuberculosis* or their homologs. In another aspect, the invention is directed to a eukaryotic expression vector that includes at least one of the genomic islands MAP-1, MAP-2, MAP-3, MAP-4, MAP-5, MAP-6, MAP-7, MAP-8, MAP-9, MAP-10, MAP-11, MAP-12, MAP-13, MAP-14, MAP-15, MAP-16, MAP-17, or MAP-18 of *M. paratuberculosis*, or their homologs. In addition to the eukaryotic expression vector, the immunological composition includes a pharmaceutically acceptable carrier. The immunological composition may optionally include an adjuvant.

This invention provides a method of treating Johne's disease in mammals. The method includes administering to a mammal a vaccine composition against *M. paratuberculosis*. The vaccine composition includes an antigen selected from the group of gcpE, pstA, kdpC, papA2, impA, umaA1, fabG2_2, aceAB, mbtH2, lpqP, map0834c, cspB, lipN, or map 1634 genes of *M. paratuberculosis*, or their homologs, or at least one of the genomic islands MAP-1, MAP-2, MAP-3, MAP-4, MAP-5, MAP-6, MAP-7, MAP-8, MAP-9, MAP-10, MAP-11, MAP-12, MAP-13, MAP-14, MAP-15, MAP-16, MAP-17, or MAP-18 of *M. paratuberculosis*, or their homologs. In addition to the antigen, the vaccine composition includes a pharmaceutically acceptable carrier. The vaccine composition may optionally include an adjuvant.

In another aspect, this invention provides a method of treating Johne's disease, which includes administering to a mammal an immunological composition comprising a vector expressing a nucleotide sequence that includes at least one of the gcpE, pstA, kdpC, papA2, impA, umaA1, fabG2_2, aceAB, mbtH2, lpqP, map0834c, cspB, lipN, or map 1634 genes of *M. paratuberculosis*, or their homologs, or at least one of the genomic islands MAP-1, MAP-2, MAP-3, MAP-4, MAP-5, MAP-6, MAP-7, MAP-8, MAP-9, MAP-10, MAP-11, MAP-12, MAP-13, MAP-14, MAP-15, MAP-16, MAP-17, or MAP-18 of *M. paratuberculosis*, or their homologs. In addition to the expression vector, the immunological composition includes a pharmaceutically acceptable carrier. The immunological composition may optionally include an adjuvant.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a schematic representation of the transposon Tn5367 from strain ATCC19698 used for insertion mutagenesis of *M. paratuberculosis*.

4

FIG. 2 depicts a genomic map showing the distribution of 1,128 transposon-insertion sites on the chromosome of *M. paratuberculosis*.

FIG. 3 depicts charts showing colonization levels of variable *M. paratuberculosis* strains to different mice organs.

FIG. 4 depicts charts showing liver and intestinal colonization levels of variable *M. paratuberculosis* strains to different mice organs.

FIG. 5 depicts a chart showing the histopathology of mice infected with *M. paratuberculosis* strains.

FIG. 6 is a genomic map showing the identification of genomic islands in the *M. avium* genome (A), and a map showing the strategy used for design of PCR primers to confirm the genomic island deletions (B).

FIG. 7 is a genomic map showing the synteny of *M. avium* and *M. paratuberculosis* genomes.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides genomic identifiers for mycobacterial species. These genomic identifiers can be used as targets for developments of vaccines and drugs against Johne's disease.

1. General Overview

The practice of the present invention employs, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry, immunology, protein kinetics, and mass spectroscopy, which are within the skill of art. Such techniques are explained fully in the literature, such as Sambrook et al., 2000, *Molecular Cloning: A Laboratory Manual*, third edition, Cold Spring Harbor Laboratory Press; *Current Protocols in Molecular Biology* Volumes 1-3, John Wiley & Sons, Inc.; Kriegler, 1990, *Gene Transfer and Expression: A Laboratory Manual*, Stockton Press, New York; Dieffenbach et al., 1995, *PCR Primer: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, each of which is incorporated herein by reference in its entirety. Procedures employing commercially available assay kits and reagents typically are used according to manufacturer-defined protocols unless otherwise noted.

Generally, the nomenclature and the laboratory procedures in recombinant DNA technology described below are those well known and commonly employed in the art. Standard techniques are used for cloning, DNA and RNA isolation, amplification and purification. Generally enzymatic reactions involving DNA ligase, DNA polymerase, restriction endonucleases and the like are performed according to the manufacturer's specifications.

2. Definitions

The phrase "nucleic acid" or "polynucleotide sequence" refers to a single or double-stranded polymer of deoxyribonucleotide or ribonucleotide bases read from the 5' to the 3' end. Nucleic acids may also include modified nucleotides that permit correct read-through by a polymerase and do not alter expression of a polypeptide encoded by that nucleic acid.

The phrase "nucleic acid sequence encoding" refers to a nucleic acid which directs the expression of a specific protein or peptide. The nucleic acid sequences include both the DNA strand sequence that is transcribed into RNA and the RNA sequence that is translated into protein. The nucleic acid sequences include both the full length nucleic acid sequences as well as non-full length sequences derived from the full length sequences. It should be further understood that the sequence includes the degenerate codons of the native sequence or sequences which may be introduced to provide codon preference in a specific host cell.

5

A “coding sequence” or “coding region” refers to a nucleic acid molecule having sequence information necessary to produce a gene product, when the sequence is expressed.

“Homology” refers to the resemblance or similarity between two nucleotide or amino acid sequences. As applied to a gene, “homolog” may refer to a gene similar in structure and/or evolutionary origin to a gene in another organism or another species. As applied to nucleic acid molecules, the term “homolog” means that two nucleic acid sequences, when optimally aligned (see below), share at least 80 percent sequence homology, preferably at least 90 percent sequence homology, more preferably at least 95, 96, 97, 98 or 99 percent sequence homology. “Percentage nucleotide (or nucleic acid) homology” or “percentage nucleotide (or nucleic acid) sequence homology” refers to a comparison of the nucleotides of two nucleic acid molecules which, when optimally aligned, have approximately the designated percentage of the same nucleotides or nucleotides that are not identical but differ by redundant nucleotide substitutions (the nucleotide substitution does not change the amino acid encoded by the particular codon). For example, “95% nucleotide homology” refers to a comparison of the nucleotides of two nucleic acid molecules which, when optimally aligned, have 95% nucleotide homology.

A “genomic sequence” or “genome” refers to the complete DNA sequence of an organism. The genomic sequences of both *M. avium* and of *M. paratuberculosis* are known and are currently available. The genomic sequence of *M. avium* can be obtained from the Institute for Genomic Research. The genomic sequence of *M. paratuberculosis* can be obtained from the GenBank, under accession number AE016958.

A “genomic island” (GI) refers to a nucleic acid region (and its homologs), that includes three or more consecutive open reading frames (ORFs), regardless of the size. A “MAP” genomic island means any genomic island (and its homologs) that is present in the *M. paratuberculosis* genome, but is not present in the *M. avium* genome. A “MAV” genomic island means any genomic island (and its homologs) that is present in the *M. avium*-genome, but is not present in the *M. paratuberculosis* genome.

A “junction” between two nucleic acid regions refers to a point that joins two nucleic acid regions. A “junction sequence” refers to a nucleic acid sequence that can be used for identification of the junction point. For example, a “junction sequence”, or a “junction region” of an inverted region (INV) and a corresponding flanking sequence refers to a nucleic acid segment that crosses the point that joins the inverted region with the flanking sequence. Such a nucleic acid segment is specific to the corresponding junction region (junction sequence), and can be used as its identifier.

The term “nucleic acid construct” or “DNA construct” is sometimes used to refer to a coding sequence or sequences operably linked to appropriate regulatory sequences so as to enable expression of the coding sequence, and inserted into an expression cassette for transforming a cell. This term may be used interchangeably with the term “transforming DNA” or “transgene”. Such a nucleic acid construct may contain a coding sequence for a gene product of interest, along with a selectable marker gene and/or a reporter gene.

A “label” is a composition detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. Useful labels include ³²P, fluorescent dyes, electron-dense reagents, enzymes (e.g., as commonly used in an ELISA), biotin, digoxigenin, or proteins for which antisera or monoclonal antibodies are available. For example, labels are preferably covalently bound to a genomic island, directly or through the use of a linker.

6

A “nucleic acid probe sequence” or “probe” is defined as a nucleic acid capable of binding to a target nucleic acid of complementary sequence through one or more types of chemical bonds, usually through complementary base pairing, usually through hydrogen bond formation. A probe may include natural (i.e., A, G, C, or T) or modified bases (7-deazaguanosine, inosine, etc.). In addition, the bases in a probe may be joined by a linkage other than a phosphodiester bond, so long as it does not interfere with hybridization. Thus, for example, probes may be peptide nucleic acids in which the constituent bases are joined by peptide bonds rather than phosphodiester linkages. Probes may bind target sequences lacking complete complementarity with the probe sequence depending upon the stringency of the hybridization conditions. The probes are preferably directly labeled as with isotopes, chromophores, lumiphores, chromogens, or indirectly labeled such as with biotin to which a streptavidin complex may later bind. By assaying for the presence or absence of the probe, one can detect the presence or absence of the select sequence or subsequence.

The term “recombinant” when used with reference, e.g., to a cell, or nucleic acid, protein, expression cassette, or vector, indicates that the cell, nucleic acid, protein, expression cassette, or vector, has been modified by the introduction of a heterologous nucleic acid or protein or the alteration of a native nucleic acid or protein, or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, underexpressed, or not expressed at all.

“Antibodies” refers to polyclonal and monoclonal antibodies, chimeric, and single chain antibodies, as well as Fab fragments, including the products of a Fab or other immunoglobulin expression library. With respect to antibodies, the term, “immunologically specific” refers to antibodies that bind to one or more epitopes of a protein of interest, but which do not substantially recognize and bind other molecules in a sample containing a mixed population of antigenic biological molecules. The present invention provides antibodies immunologically specific for part or all of the polypeptides of the present invention, e.g., those polypeptides encoded by the genes *gcpE*, *pstA*, *kdpC*, *papA2*, *impA*, *umaA1*, *fabG2_2*, *aceAB*, *mbtH2*, *lpqP*, *map0834c*, *cspB*, *lipN*, and *map1634* of *Mycobacterium paratuberculosis*.

An “expression cassette” refers to a nucleic acid construct, which when introduced into a host cell, results in transcription and/or translation of a RNA or polypeptide, respectively. Expression cassettes can be derived from a variety of sources depending on the host cell to be used for expression. An expression cassette can contain components derived from a viral, bacterial, insect, plant, or mammalian source. In the case of both expression of transgenes and inhibition of endogenous genes (e.g., by antisense, or sense suppression) the inserted polynucleotide sequence need not be identical and can be “substantially identical” to a sequence of the gene from which it was derived.

The term “vector” refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. One type of vector is a “plasmid”, which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors

can be integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as “recombinant expression vectors” (or simply, “expression vectors”). In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, “plasmid” and “vector” may be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

The terms “isolated,” “purified,” or “biologically pure” refer to material that is substantially or essentially free from components that normally accompany it as found in its native state. Purity and homogeneity are typically determined using analytical chemistry techniques such as polyacrylamide gel electrophoresis or high performance liquid chromatography. A protein that is the predominant species present in a preparation is substantially purified. In particular, an isolated nucleic acid of the present invention is separated from open reading frames that flank the desired gene and encode proteins other than the desired protein. The term “purified” denotes that a nucleic acid or protein gives rise to essentially one band in an electrophoretic gel. Particularly, it means that the nucleic acid or protein is at least 85% pure, more preferably at least 95% pure, and most preferably at least 99% pure.

In the case where the inserted polynucleotide sequence is transcribed and translated to produce a functional polypeptide, because of codon degeneracy a number of polynucleotide sequences will encode the same polypeptide. These variants are specifically covered by the term “polynucleotide sequence from” a particular gene. In addition, the term specifically includes sequences (e.g., full length sequences) substantially identical (determined as described below) with a gene sequence encoding a protein of the present invention and that encode proteins or functional fragments that retain the function of a protein of the present invention, e.g., a disease causing agent of *M. paratuberculosis*.

The term “immunization” is the process by which an individual is exposed to a material that is designed to stimulate his or her immune system against that material. The material is known as an “immunizing agent” or “immunogen”. When the immunizing agent is administered to a subject, the subject develops an immune response, which can be used for prevention and treatment against Johne’s disease or Crohn’s disease.

The term “vaccine” refers to an antigenic preparation used to produce immunity to a disease, in order to prevent or ameliorate the effects of infection. Vaccines are typically prepared using a combination of an immunologically effective amount of an immunogen together with an adjuvant effective for enhancing the immune response of the vaccinated subject against the immunogen. The process of distributing and administering vaccines is referred to as “vaccination”.

Vaccine formulations will contain a “therapeutically effective amount” of the active ingredient, that is, an amount capable of eliciting an immune response in a subject to which the composition is administered. In the treatment and prevention of Johne’s disease, for example, a “therapeutically effective amount” would preferably be an amount that enhances resistance of the vaccinated subject to new infection and/or reduces the clinical severity of the disease. Such protection will be demonstrated by either a reduction or lack of symptoms normally displayed by a subject infected with Johne’s

disease, a quicker recovery time and/or a lowered count of *M. paratuberculosis* bacteria. Vaccines can be administered prior to infection, as a preventative measure against Johne’s or Crohn’s disease. Alternatively, vaccines can be administered after the subject already has contracted a disease. Vaccines given after exposure to mycobacteria may be able to attenuate the disease, triggering a superior immune response than the natural infection itself.

A “pharmaceutically acceptable carrier” means any conventional pharmaceutically acceptable carrier, vehicle, or excipient that is used in the art for production and administration of vaccines. Pharmaceutically acceptable carriers are typically non-toxic, inert, solid or liquid carriers.

The term “adjuvant” refers to a compound that enhances the effectiveness of the vaccine, and may be added to the formulation that includes the immunizing agent. Adjuvants provide enhanced immune response even after administration of only a single dose of the vaccine. Adjuvants may include, for example, muramyl dipeptides, avridine, aluminum hydroxide, dimethyldioctadecyl ammonium bromide (DDA), oils, oil-in-water emulsions, saponins, cytokines, and other substances known in the art. Examples of suitable adjuvants are described in U.S. Patent Application Publication No. US2004/0213817 A1.

In the case of polynucleotides used to immunize a subject, the introduced sequence need not be perfectly identical to a sequence of the target endogenous gene. The introduced polynucleotide sequence is typically at least substantially identical (as determined below) to the target endogenous sequence.

Two nucleic acid sequences or polypeptides are said to be “identical” if the sequence of nucleotides or amino acid residues, respectively, in the two sequences is the same when aligned for maximum correspondence as described below.

The term “complementary to” is used herein to mean that the sequence is complementary to all or a portion of a reference polynucleotide sequence.

The term “biologically active fragment” is intended to mean a part of the complete molecule which retains all or some of the catalytic or biological activity possessed by the complete molecule, especially activity that allows specific binding of the antibody to an antigenic determinant.

“Functional equivalents” of an antibody include any molecule capable of specifically binding to the same antigenic determinant as the antibody, thereby neutralizing the molecule, e.g., antibody-like molecules, such as single chain antigen binding molecules.

Optimal alignment of sequences for comparison may be conducted by methods commonly known in the art, e.g., the local homology algorithm (Smith and Waterman, 1981, *Adv. Appl. Math.* 2: 482-489), by the search for similarity method (Pearson and Lipman 1988, *Proc. Natl. Acad. Sci. USA* 85: 2444-2448), by computerized implementations of these algorithms (GAP, BESTFIT, BLAST, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group (GCG), Madison, Wis.), or by inspection.

Protein and nucleic acid sequence identities are evaluated using the Basic Local Alignment Search Tool (“BLAST”) which is well known in the art (Karlin and Altschul, 1990, *Proc. Natl. Acad. Sci. USA* 87: 2267-2268; Altschul et al., 1997, *Nucl. Acids Res.* 25: 3389-3402). The BLAST programs identify homologous sequences by identifying similar segments, which are referred to herein as “high-scoring segment pairs,” between a query amino or nucleic acid sequence and a test sequence which is preferably obtained from a protein or nucleic acid sequence database. Preferably, the statistical significance of a high-scoring segment pair is

evaluated using the statistical significance formula (Karlin and Altschul, 1990), the disclosure of which is incorporated by reference in its entirety. The BLAST programs can be used with the default parameters or with modified parameters provided by the user.

"Percentage of sequence identity" is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

The term "substantial identity" of polynucleotide sequences means that a polynucleotide comprises a sequence that has at least 25% sequence identity. Alternatively, percent identity can be any integer from 25% to 100%. More preferred embodiments include at least: 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% compared to a reference sequence using the programs described herein; preferably BLAST using standard parameters, as described. These values can be appropriately adjusted to determine corresponding identity of proteins encoded by two nucleotide sequences by taking into account codon degeneracy, amino acid similarity, reading frame positioning and the like.

"Substantial identity" of amino acid sequences for purposes of this invention normally means polypeptide sequence identity of at least 40%. Preferred percent identity of polypeptides can be any integer from 40% to 100%. More preferred embodiments include at least 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 98.7%, or 99%.

Polypeptides that are "substantially similar" share sequences as noted above except that residue positions which are not identical may differ by conservative amino acid changes. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulfur-containing side chains is cysteine and methionine. Preferred conservative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, aspartic acid-glutamic acid, and asparagine-glutamine.

3. Identification of Vaccine Targets of the Present Invention

The invention described here utilizes large-scale identification of disrupted genes and the use of bioinformatics to select mutants that could be characterized in animals. Employing such an approach, novel virulence determinants were identified, based on mutants that were investigated in mice. These virulence determinants can be used for designing vaccines. Compared to similar protocols established for identifying virulence genes such as signature-tagged mutagenesis

(Ghadiali et al., 2003, *Nucleic Acids Res.* 31: 147-151), the approach employed here is simpler and uses a smaller number of animals.

The present invention also provides immunogenic preparations and vaccines containing at least one plasmid encoding and expressing at least one immunogen against *M. paratuberculosis* compositions formulated with an adjuvant.

The target nucleic acid sequences of the present invention include the gcpE, pstA, kdpC, papA2, impA, umaA1, fabG2_2, aceAB, mbtH2, lpqP, map0834c, cspB, lipN, and map1634 genes of *M. paratuberculosis*, their homologs, and the corresponding gene products. Presence of these genes, their homologs, and/or their products in a sample is indicative of a *M. paratuberculosis* infection.

The start and end coordinates of the *M. paratuberculosis* polynucleotides of this invention (e.g., genes, genomic islands, inverted regions, junction sequences) are based on the genomic sequence of *M. paratuberculosis* strain K10 (Li et al., 2005, Proc. Natl. Acad. Sci. USA 102: 12344-12349; GenBank No. AE016958). The start and end coordinates of the *M. avium* polynucleotides of this invention (e.g., genes, genomic islands, inverted regions, junction sequences) are based on the genomic sequence of *M. avium* strain 104, as obtained from The Institute for Genomic Research.

The size of gcpE is 1167 base pairs (bp), and it is located at positions 3272755 through 3273921 of the *M. paratuberculosis* genomic sequence.

The size of pstA is 12084 base pairs (bp), and it is located at positions 1309241 through 1321324 of the *M. paratuberculosis* genomic sequence.

The size of kdpC is 876 base pairs (bp), and it is located at positions 1038471 through 1039346 of the *M. paratuberculosis* genomic sequence.

The size of papA2 is 1518 base pairs (bp), and it is located at positions 1854059 through 1855576 of the *M. paratuberculosis* genomic sequence.

The size of impA is 801 base pairs (bp), and it is located at positions 1386766 through 1387566 of the *M. paratuberculosis* genomic sequence.

The size of umaA1 is 861 base pairs (bp), and it is located at positions 4423752 through 4424612 of the *M. paratuberculosis* genomic sequence.

The size of fabG2_2 is 750 base pairs (bp), and it is located at positions 2704522 through 2705271 of the *M. paratuberculosis* genomic sequence.

The size of aceAB is 2288 base pairs (bp), and it is located at positions 1795784 through 1798072 of the *M. paratuberculosis* genomic sequence.

The size of mbtH2 is 233 base pairs (bp), and it is located at positions 2063983 through 2064216 of the *M. paratuberculosis* genomic sequence.

The size of lpqP is 971 base pairs (bp), and it is located at positions 4755529 through 4756500 of the *M. paratuberculosis* genomic sequence.

The size of map0834c is 701 base pairs (bp), and it is located at positions 851908 through 852609 of the *M. paratuberculosis* genomic sequence.

The size of map1634 is 917 base pairs (bp), and it is located at positions 1789023 through 1789940 of the *M. paratuberculosis* genomic sequence.

In another aspect, the virulence determinants of the present invention include genomic islands (GIs). These GIs are strain-specific. The inventors have identified 18 *M. paratuberculosis*-specific genomic islands (MAPs), that are absent from the *M. avium* genome (Table 8).

11

The size of MAP-1 is 19,343 base pairs (bp). MAP-1 includes 17 ORFs. MAP-1 is located at positions 99,947 through 119,289 of the *M. paratuberculosis* genomic sequence.

The size of MAP-2 is 3,858 base pairs (bp). MAP-2 includes 3 ORFs. MAP-2 is located at positions 299,412 through 303,269 of the *M. paratuberculosis* genomic sequence.

The size of MAP-3 is 2,915 base pairs (bp). MAP-3 includes 3 ORFs. MAP-3 is located at positions 410,091 through 413,005 of the *M. paratuberculosis* genomic sequence.

The size of MAP-4 is 16,681 base pairs (bp). MAP-4 includes 17 ORFs. MAP-4 is located at positions 872,772 through 889,452 of the *M. paratuberculosis* genomic sequence.

The size of MAP-5 is 14,191 base pairs (bp). MAP-5 includes 17 ORFs. MAP-5 is located at positions 989,744 through 1,003,934 of the *M. paratuberculosis* genomic sequence.

The size of MAP-6 is 8,971 base pairs (bp). MAP-6 includes 6 ORFs. MAP-6 is located at positions 1,291,689 through 1,300,659 of the *M. paratuberculosis* genomic sequence.

The size of MAP-7 is 6,914 base pairs (bp). MAP-7 includes 6 ORFs. MAP-7 is located at positions 1,441,777 through 1,448,690 of the *M. paratuberculosis* genomic sequence.

The size of MAP-8 is 7,915 base pairs (bp). MAP-8 includes 8 ORFs. MAP-8 is located at positions 1,785,511 through 1,793,425 of the *M. paratuberculosis* genomic sequence.

The size of MAP-9 is 11,202 base pairs (bp). MAP-9 includes 10 ORFs. MAP-9 is located at positions 1,877,255 through 1,888,456 of the *M. paratuberculosis* genomic sequence.

The size of MAP-10 is 2,993 base pairs (bp). MAP-10 includes 3 ORFs. MAP-10 is located at positions 1,891,000 through 1,893,992 of the *M. paratuberculosis* genomic sequence.

The size of MAP-11 is 2,989 base pairs (bp). MAP-11 includes 4 ORFs. MAP-11 is located at positions 2,233,123 through 2,236,111 of the *M. paratuberculosis* genomic sequence.

The size of MAP-12 is 11,977 base pairs (bp). MAP-12 includes 11 ORFs. MAP-12 is located at positions 2,378,957 through 2,390,933 of the *M. paratuberculosis* genomic sequence.

The size of MAP-13 is 19,977 base pairs (bp). MAP-13 includes 19 ORFs. MAP-13 is located at positions 2,421,552 through 2,441,528 of the *M. paratuberculosis* genomic sequence.

The size of MAP-14 is 19,315 base pairs (bp). MAP-14 includes 19 ORFs. MAP-14 is located at positions 3,081,906 through 3,101,220 of the *M. paratuberculosis* genomic sequence.

The size of MAP-15 is 4,143 base pairs (bp). MAP-15 includes 3 ORFs. MAP-15 is located at positions 3,297,661 through 3,301,803 of the *M. paratuberculosis* genomic sequence.

The size of MAP-16 is 79,790 base pairs (bp). MAP-16 includes 56 ORFs. MAP-16 is located at positions 4,140,311 through 4,220,100 of the *M. paratuberculosis* genomic sequence.

12

The size of MAP-17 is 3,655 base pairs (bp). MAP-17 includes 5 ORFs. MAP-17 is located at positions 4,735,049 through 4,738,703 of the *M. paratuberculosis* genomic sequence.

The size of MAP-18 is 3,512 base pairs (bp). MAP-18 includes 3 ORFs. MAP-18 is located at positions 4,800,932 through 4,804,443 of the *M. paratuberculosis* genomic sequence.

The inventors have also identified 24 *M. avium*-specific genomic islands (MAVs), that are absent from the *M. paratuberculosis* genome (Table 9).

The GIs of the present invention (both MAPs and MAVs) can be used as target nucleic acid sequences for design of vaccines and drugs that are strain-specific. Thus, the targets enable one skilled in the art to distinguish between the presence of *M. paratuberculosis* or *M. avium* in a sample. Should both *Mycobacterium* strains be present in a sample, one should be able to identify the presence of both classes of target polynucleotides in the sample.

It is possible to diagnose the presence of *M. paratuberculosis* or *M. avium* in a sample due to the inversion of three large genomic fragments in *M. paratuberculosis* in comparison to *M. avium*. It was unexpectedly discovered that, when the GIs associated with both genomes were aligned, three large genomic fragments in *M. paratuberculosis* were identified as inverted relative to the corresponding genomic fragments in *M. avium*. These inverted nucleic acid regions (INV) had the sizes of approximately 54.9 Kb, 863.8 Kb and 1,969.4 Kb (FIG. 7).

The target polynucleotide may be DNA. In some variations, the target polynucleotide may be obtained from total cellular DNA, or in vitro amplified DNA.

The specificity of single stranded DNA to hybridize complementary fragments is determined by the "stringency" of the reaction conditions. Hybridization stringency increases as the propensity to form DNA duplexes decreases. In nucleic acid hybridization reactions, the stringency can be chosen to either favor specific hybridizations (high stringency), which can be used to identify, for example, full-length clones from a library. Less-specific hybridizations (low stringency) can be used to identify related, but not exact, DNA molecules (homologous, but not identical) or segments.

Identification of target sequences of the present invention may be accomplished by a number of techniques. For instance, oligonucleotide probes based on the sequences disclosed here can be used to identify the desired gene in a cDNA or genomic DNA library from a desired bacterial strain. To construct genomic libraries, large segments of genomic DNA are generated by random fragmentation, e.g. using restriction endonucleases, and are ligated with vector DNA to form concatemers that can be packaged into the appropriate vector. The cDNA or genomic library can then be screened using a probe based upon the sequence of a cloned gene such as the polynucleotides disclosed here. Probes may be used to hybridize with genomic DNA or cDNA sequences to identify homologous genes in the same or different bacterial strains.

Alternatively, the nucleic acids of interest can be amplified from nucleic acid samples using amplification techniques. For instance, polymerase chain reaction (PCR) technology can be used to amplify the sequences of the genes directly from mRNA, from cDNA, from genomic libraries or cDNA libraries. PCR and other in vitro amplification methods may also be useful, for example, to clone nucleic acid sequences that code for proteins to be expressed, to make nucleic acids to use as probes for detecting the presence of the desired mRNA in samples, for nucleic acid sequencing, or for other purposes.

Appropriate primers and probes for identifying the target sequences of the present invention from a sample are generated from comparisons of the sequences provided herein, according to standard PCR guides. For examples of primers used see the Examples section below.

Polynucleotides may also be synthesized by well-known techniques described in the technical literature. Double-stranded DNA fragments may then be obtained either by synthesizing the complementary strand and annealing the strands together under appropriate conditions, or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Once a nucleic acid is isolated using the method described above, standard methods can be used to determine if the nucleic acid is a preferred nucleic acid of the present invention, e.g., by using structural and functional assays known in the art. For example, using standard methods, the skilled practitioner can compare the sequence of a putative nucleic acid sequence thought to encode a preferred protein of the present invention to a nucleic acid sequence encoding a preferred protein of the present invention to determine if the putative nucleic acid is a preferred polynucleotide of the present invention.

Gene amplification and/or expression can be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA, dot blotting (DNA analysis), DNA microarrays, or in situ hybridization, using an appropriately labeled probe, based on the sequences provided herein. Various labels can be employed, most commonly fluorochromes and radioisotopes, particularly ^{32}P . However, other techniques can also be employed, such as using biotin-modified nucleotides for introduction into a polynucleotide. The biotin then serves as the site for binding to avidin or antibodies, which can be labeled with a variety of labels, such as radionuclides, fluoroscopes, enzymes, or the like. Alternatively, antibodies can be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn can be labeled and the assay can be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression can also be measured by immunological methods, such as immunohistochemical staining. With immunohistochemical staining techniques, a sample is prepared, typically by dehydration and fixation, followed by reaction with labeled antibodies specific for the gene product coupled, where the labels are usually visually detectable, such as enzymatic labels, fluorescent labels, luminescent labels, and the like. Gene expression can also be measured using PCR techniques, or using DNA microarrays, commonly known as gene chips.

4. DNA Vaccines

The use of deoxyribonucleic acid (DNA) molecules for vaccination is also known (Wolf et al., 1990, *Science* 247: 1465-1468). This vaccination technique induces cellular and humoral immunity (stimulation of the production of antibodies specifically directed against the immunogen) after in vivo transfection of cells of the subject to be vaccinated with nucleic acids encoding immunologically active proteins.

A "DNA vaccine" or "immunogenic" or "immunological composition" is composed of at least one vector (e.g., plasmid) which may be expressed by the cellular machinery of the subject to be vaccinated or inoculated and of a pharmaceutically acceptable carrier, vehicle, or excipient. The nucleotide sequence of this vector encodes one or more immunogens,

such as proteins or glycoproteins capable of inducing, in the subject to be vaccinated or inoculated, a cellular immune response (mobilization of the T lymphocytes) and a humoral immune response (Davis, 1997, *Current Opinion Biotech.* 8: 635-640).

The present invention provides DNA vaccines or immunogenic or immunological compositions for mammals. These DNA vaccines can be generated using the information on target polynucleotides that constitute virulence determinants of Johne's disease or Crohn's disease. In one aspect, the immunized mammals develop an immune response, which can be used for prevention and treatment against Johne's disease or Crohn's disease.

Various routes of administration of the DNA vaccine have been proposed (intraperitoneal, intravenous, intramuscular, subcutaneous, intradermal, mucosal, and the like), and they are useful for the practice of this invention. Various means of administration have also been proposed. Some means include the use of gold particles coated with DNA and projected so as to penetrate into the cells of the skin of the subject to be vaccinated (Tang et al., 1992, *Nature* 356: 152-154). Other means include the use of liquid jet injectors which make it possible to transfect both skin cells and cells of the underlying tissues (Furth et al., 1992, *Analytical Bioch.* 205: 365-368).

The invention also relates to small nucleic acids that selectively hybridize to the exemplified target polynucleotide sequences, including hybridizing to the exact complements of these sequences. Such small nucleic acids include oligonucleotides or small interfering ribonucleic acid (siRNA) molecules.

The invention further provides small interfering ribonucleic acid (siRNA) molecules for prevention and treatment of Johne's or Crohn's diseases. RNA interference (RNAi) using siRNA has been shown to be an effective means of silencing gene expression in cells. For example, retroviral vectors that express small RNAs as hairpin loops can be used for therapeutic purposes.

The oligonucleotide or siRNA may be partially complementary to the target nucleic acid sequence. Alternatively, the oligonucleotide may be exactly complementary to the target nucleic acid sequence. The oligonucleotide or siRNA molecule may be greater than about 4 nucleic acid bases in length and/or less than about 48 nucleic acid bases in length. In a further variation, the oligonucleotide or the siRNA molecule may be about 20 nucleic acid bases in length.

This invention provides a method for delivering an isolated polynucleotide to the interior of a cell in a mammal, comprising the interstitial introduction of an isolated polynucleotide into a tissue of the mammal where the polynucleotide is taken up by the cells of the tissue and exerts a therapeutic effect on the mammal. The method can be used to deliver a therapeutic polypeptide to the cells of the mammal, to provide an immune response upon in vivo transcription and/or translation of the polynucleotide, or to deliver antisense polynucleotides.

It is possible to coadminister DNA vaccines encoding antigen with siRNA targeting the target nucleic acid sequences of this invention, to enhance the antigen-specific cell responses, and elicit potent antibacterial effects in vaccinated subjects. Similarly, a skilled artisan should know to use combined/composite vaccines (see e.g., Talaat et al., 2002, *Vaccine* 20: 538-544, incorporated herein in entirety by reference), to increase the efficacy while reducing the number of vaccinations. For example, two or more antigens of this invention may be combined in a composite vaccine directed against Johne's disease or Crohn's disease.

The vaccines may include other components to serve certain functions, for example, directing the nucleic acid to a

certain location in the cell or directing transcription of the antigen. Compositions for transport to the nucleus may be included, particularly members of the high mobility group (HMG), more particularly HMG-1, which is a non-histone DNA-binding protein. In combination with antisense molecules, RNAses such as RNaseH, may be used. Other proteins that will aid or enhance the function of the antigen may be included, such as peptide sequences that direct antigen processing, particularly HLA presentation, or movement in the cytoplasm.

In one embodiment, the immunized mammals are farm animals, in particular cattle. The immunized cattle develop an immune response, which can be used for prevention and treatment against Johne's disease.

5. Antibodies

The present invention further provides for antibodies immunologically specific for all or part, e.g., an amino-terminal portion, of a polypeptide at least 70% identical to a mycobacterial sequence that is a virulence determinant. Exemplary anti-Johne's and anti-Crohn's antibodies include polyclonal, monoclonal, humanized, bispecific, and hetero-conjugate antibodies. The invention also provides functional equivalents of anti-Johne's and anti-Crohn's antibodies, e.g., antibody-like molecules, such as single chain antigen binding molecules.

The antibodies of this invention may be polyclonal antibodies. Methods of preparing polyclonal antibodies are known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include any of the antigens of this invention, its homolog, or a fusion protein thereof. Examples of adjuvants which may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

The antibodies of this invention may alternatively be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods. In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized in vitro.

The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Pat. No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences (U.S. Pat. No. 4,816,567) or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin

polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

The antibodies of the invention may further comprise humanized antibodies or human antibodies. Humanized antibodies might be preferably used for prevention and treatment of Crohn's disease. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin.

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the methods described in Riechmann et al., 1988, *Nature*, 332: 323-327; and in Verhoeyen et al., 1988, *Science* 239: 1534-1536, by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Pat. No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Administration of Vaccines

In one aspect, a method of treating Johne's disease or Crohn's disease is disclosed. In one embodiment, the method includes production of antibodies directed to *M. paratuberculosis* virulence proteins. This invention discloses a variety of proteins that are virulence determinants, and are thus indicative of *M. paratuberculosis* infection. Methods known in the art can be used to immunize subjects (animals and humans) for purposes of prevention and treatment against Johne's disease or Crohn's disease. Pharmaceutically acceptable carriers are typically used for administration of vaccine compositions. For example, the use of solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like media and agents for pharmaceutical active carriers is well known in the art.

In one embodiment, administration of an immunizing agent includes administering in vivo into a tissue of a mammal a construct comprising a nucleotide sequence encoding

an antigen, in an amount sufficient that uptake of the construct into cells of the mammal occurs, and sufficient expression results, to generate a detectable antibody response. In a preferred embodiment, the nucleotide sequence encodes an antigen that includes at least one of gcpE, pstA, kdpC, papA2, impA, umaA1, fabG2₂, aceAB, mbtH2, lpqP, map0834c, cspB, lipN, or map1634 genes of *M. paratuberculosis*. In another preferred embodiment, the nucleotide sequence encodes an antigen that includes at least one of the MAP-1, MAP-2, MAP-3, MAP-4, MAP-5, MAP-6, MAP-7, MAP-8, MAP-9, MAP-10, MAP-11, MAP-12, MAP-13, MAP-14, MAP-15, MAP-16, MAP-17, or MAP-18 genomic islands of *M. paratuberculosis*.

The vaccine provided by this invention may be administered subcutaneously, intramuscularly, intradermally, or into an organ. Intramuscular injection has been shown in the past to be an important delivery route for induction of immunity. Skeletal muscle has properties such as high vascularization and multi-nucleation. In addition, it is nonreplicating and capable of expressing recombinant proteins. These properties are advantageous for gene therapy using DNA vaccines. One theory of the mechanism of how muscle presents the protein and induces immune response is that recombinant protein is produced and released into the vascular network of the muscle and eventually presented by professional antigen-presenting cells such as dendritic cells, myoblasts, or macrophages infiltrating the muscle. Another suggestion is that at the injection site muscle injury induces myoblast proliferation and activation of infiltrating macrophages or dendritic-like cells, and they then present antigens through MHC class II antigen. Thus, other tissues which have similar qualities also would be good delivery sites for the vaccine.

The chosen route of administration will depend on the vaccine composition and the disease status of subjects. Relevant considerations include the types of immune cells to be activated, the time which the antigen is exposed to the immune system and the immunization schedule. Although many vaccines are administered consecutively within a short period, spreading the immunizations over a longer time may maintain effective clinical and immunological responses.

To immunize a subject, the vaccine is preferably administered parenterally, usually by intramuscular injection. Other modes of administration, however, such as sub-cutaneous, intraperitoneal and intravenous injection, are also acceptable. The quantity to be administered depends on the subject to be treated, the capacity of the subject's immune system to synthesize antibodies, and the degree of protection desired. Effective dosages can be readily established by one of ordinary skill in the art through routine trials establishing dose response curves. The subject is immunized by administration of the vaccine in at least one dose, and preferably two to four doses. Moreover, the subject may be administered as many doses as is required to maintain a state of immunity to infection.

Additional vaccine formulations that are suitable for other modes of administration include suppositories and, in some cases, aerosol, intranasal, oral formulations, and sustained release formulations. For suppositories, the vehicle composition will include traditional binders and carriers, such as, polyalkaline glycols, or triglycerides. Oral vehicles include such normally employed excipients as, for example, pharmaceutical grades of mannitol, lactose, starch, magnesium stearate, sodium saccharin cellulose, magnesium carbonate, and the like. The oral vaccine compositions may be taken in the form of solutions, suspensions, tablets, pills, capsules, sustained release formulations, or powders.

Another aspect of the invention provides a pharmaceutical product for use in immunizing a mammal, comprising a pharmaceutically effective amount of a polynucleotide encoding an immunogenic polypeptide, a sealed container enclosing the polynucleotide in a sterile fashion, and means associated with the container for permitting transfer of the polynucleotide from the container to the interstitial space of a tissue, whereby cells of the tissue can take up and express the polynucleotide.

One skilled in the art will know that it is possible to enhance the immune response of an animal to a target immunogen by using a variety of adjuvants. Suitable adjuvants are, for example, described in U.S. Patent Application Pub. No. US 2004/0213817 A1, incorporated herein in entirety by reference.

The invention is also directed to a kit for vaccination against Johnes's or Crohn's disease. The kit may include one or more of a sample that includes a target polynucleotide, and one or more nucleic acid probe sequences at least partially complementary to a target nucleic acid sequence. The kit may include instructions for using the kit.

It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications, patents, and patent applications cited herein are hereby incorporated by reference in their entirety for all purposes.

EXAMPLES

It is to be understood that this invention is not limited to the particular methodology, protocols, patients, or reagents described, and as such may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention, which is limited only by the claims. The following examples are offered to illustrate, but not to limit the claimed invention.

Example 1

Animals

Groups of BALB/c mice (N=10-20) at 3 to 4 weeks of age were infected with *M. paratuberculosis* strains using intraperitoneal (IP) injection. Infected mice were sacrificed at 3, 6 and 12 weeks post-infection and their livers, spleens and intestines collected for both histological and bacteriological examinations. Tissue sections collected for histopathology were preserved in 10% neutralized buffer formalin (NBF) before embedding in paraffin, cut into 4-5 μ m sections, stained with hematoxylin and eosin (HE) or acid fast staining (AFS). Tissue sections from infected animals were examined by two independent pathologists at 3, 6 and 12 weeks post infection. The severity of inflammatory responses was ranked using a score of 0 to 5 based on lesion size and number per field. Tissues with more than 3 fields containing multiple, large-sized lesions were given a score of 5 using the developed scale.

Bacterial Strains, Cultures and Vectors

Mycobacterium avium subsp. *paratuberculosis* strain ATCC 19698 (*M. paratuberculosis*) was used for constructing the mutant library. This strain was grown at 37° C. in Middlebrook 7H9 broth enriched with 10% albumin dextrose

complex (ADC), 0.5% glycerol, 0.05% Tween 80 and 2 mg/ml of mycobactin J (Allied Monitor, IN).

The temperature-sensitive, conditionally replicating phasmid (phAE94) used to deliver the transposon Tn5367 was obtained from Bill Jacobs laboratory (Albert Einstein College of Medicine) and propagated in *Mycobacterium smegmatis* mc² 155 at 30° C. as described previously (Bardarov et al., 1997, *Proc. Natl. Acad. Sci. USA* 94: 10961-10966). The Tn5367 is an IS1096-derived insertion element containing a kanamycin resistance gene as a selectable marker.

After phage transduction, mutants were selected on Middlebrook 7H10 medium plates supplemented with 30 µg/ml of kanamycin. *Escherichia coli* DH5α cells used for cloning purposes were grown on Luria-Bertani (LB) agar or broth supplemented with 100 µg/ml ampicillin. The plasmid vector pGEM T-easy (Promega, Madison, Wis.) was used for TA cloning the PCR products before sequencing.

Construction of a Transposon Mutants Library

The phasmid phAE94 was used to deliver the Tn5367 to mycobacterial cells using a protocol established earlier for *M. tuberculosis*. For each transduction, 10 ml of *M. paratuberculosis* culture was grown to 2×10⁸ CFU/ml (OD₆₀₀ 0.6-0.8), centrifuged and resuspended in 2.5 ml of MP buffer (50 mM Tris-HCl [pH 7.6], 150 mM NaCl, 2 mM CaCl₂) and incubated with 10¹⁰ PFU of phAE94 at the non-permissive temperature (37° C.) for 2 h in a shaking incubator to inhibit a possible lytic or lysogenic cycle of the phage.

Adsorption stop buffer (20 mM sodium citrate and 0.2% Tween 80) was added to prevent further phage infections and this mixture was plated immediately on 7H10 agar supplemented with 30 µg/ml of kanamycin and incubated at 37° C. for 6 weeks. Kanamycin-resistant colonies (5,060) were inoculated into 2 ml of 7H9 broth supplemented with kanamycin in 96-well microtitre plates for additional analysis.

Construction of lipN mutant. The lipN gene was deleted from *M. paratuberculosis* K10 strain using a homologous recombination protocol based on phage transduction. The whole gene was deleted from *M. paratuberculosis* K10 and was tested in mice. This gene was selected for deletion because of its up-regulation when DNA microarrays were used to analyze in vivo (fecal samples) collected from infected cows with high levels of mycobacterial shedding.

Southern Blot Analysis

To examine the randomness of Tn5367 insertions in the *M. paratuberculosis* genome, 10 randomly selected mutants were analyzed by Southern blot using a standard protocol. Kanamycin-resistant *M. paratuberculosis* single colonies were grown separately in 10 ml of 7H9 broth for 10 days at 37° C. before genomic DNA extraction and digestion (2-3 µg) with BamH1 restriction enzyme. Digested DNA fragments from both mutant and wild-type strains were electrophoresed on a 1% agarose gel and transferred to a nylon membrane (Perkin Elmer, Calif.), using an alkaline transfer protocol as recommended by the manufacturer.

A 1.3-kb DNA fragment from the kanamycin resistance gene was radiolabeled with [α-³²P]-dCTP using a Random Prime Labeling Kit (Promega) in accordance with the manufacturer's direction. The radio-labeled probe was hybridized to the nylon membrane at 65° C. overnight in a shaking water bath before washing, exposure to X-ray film, and development to visualize hybridization signals.

Sequencing of the Transposon Insertion Site

FIG. 1 shows a schematic representation of the transposon Tn5367 from strain ATCC19698 used for insertion mutagenesis of *M. paratuberculosis*. To determine the exact transposon insertion site within the *M. paratuberculosis* genome, a protocol for sequencing randomly primed PCR products was

adopted from previous work on *M. tuberculosis* with slight modifications. For PCR amplification, the genomic DNA of each mutant was extracted from individual cultures by boiling for 10 min, centrifuged at 10,000×g for 1 min, and 10 µl of the supernatants were used in a standard PCR reaction. For the first round of PCR, a transposon-specific primer (AMT31: 5'-TGCAGCAACGCCAGGTCCACACT-3') (SEQ ID NO:1) and the degenerate primer (AMT38: 5'-GTMTAC-GACTCACTATAGGGCNNNNNCATG-3') (SEQ ID NO:2) were used to amplify the chromosomal sequence flanking the transposon-insertion site.

PCR was carried out in a total volume of 25 µl in 10 mM Tris/HCl (pH 8.3), 50 mM KCl, 2.0 mM MgCl₂, 0.01% (w/v) BSA, 0.2 mM dNTPs, 0.1 µM of primer AMT31, 1.0 µM of primer AMT38 and 0.75 U Taq polymerase (Promega). First-round amplification was performed with an initial denaturing step at 94° C. for 5 min, followed by 40 cycles of denaturing at 94° C. for 1 min, annealing at 50° C. for 30 s and extension at 72° C. for 90 s, with a final extension step at 72° C. for 7 min. Only 1 µl of the first round amplification was then used as a template for the second round PCR (nested PCR) using a nested primer (AMT32: 5'-CTCTTGCTCTTCCGCTTCT-TCTCC-3') (SEQ ID NO:3) derived from the Tn5367 and T7 primer (AMT 39: 5'-TAATACGACTCACTATAGGG-3') (SEQ ID NO:4) present within the degenerative primer sequence. Reactions were carried out in a total volume of 50 µl in 10 mM Tris/HCl (pH 8.3), 50 mM KCl, 1.5 mM MgCl₂, 0.2 mM dNTPs, 0.5 µM primers, 5% (v/v) DMSO and 0.75 U Taq polymerase.

A final round of amplification was performed with a denaturing step at 95° C. for 5 min followed by 35 thermocycles (94° C. for 30 s, 57° C. for 30 s and 72° C. for 1 min) with a final extension step at 72° C. for 10 min. For almost 2/3 of the sequenced mutants, no cloning was attempted and AMT152 primer (5'-TTGCTCTTCCGCTTCTTCT-3') (SEQ ID NO:5) present in Tn5367 was used to directly sequence gel-purified amplicons. The product of the second amplification was gel-purified (Wizard Gel-extraction kit, Promega, Madison, Wis.) and cloned into PGEM T-easy vector for plasmid mini-preparation followed by automatic sequencing. Inserts in pGEM T-easy vector was confirmed by EcoRI restriction digestion and the sequencing was carried out using SP-6 primer (5'-TATTTAGGTGACACTATAG-3') (SEQ ID NO:6).

To identify the precise transposon-insertion site in the *M. paratuberculosis* genome, the transposon sequence was trimmed from the cloning vector sequences and a BLASTN search was used against the *M. paratuberculosis* K-10 complete genome sequence (GenBank accession no. AE016958). Sequences with at least 100 bp of alignment matching to the *M. paratuberculosis* genome were further analyzed while others without any transposon sequence were not analyzed to avoid using amplicons generated by non-specific primer binding and amplification.

Statistical Analysis

All bacterial counts from mouse organs were statistically analyzed using the Excel program (Microsoft, Seattle, Wash.). All counts are expressed as the mean±standard deviation (S.D.). Differences in counts between groups were analyzed with a Student's t-test for paired samples. Differences were considered to be significant if a probability value of p<0.05 was obtained when the CFU count of mutant strains were compared to that of the wild-type strain.

Generation of *M. paratuberculosis* Mutant Library

A genome-wide random-insertion mutant library was generated for the *M. paratuberculosis* ATCC 19698 using the temperature-sensitive mycobacteriophage phAE94 developed earlier for *M. tuberculosis*. A library consisting of 5,060

21

kanamycin-resistance colonies was obtained by the insertion of transposon Tn5367 in the bacterial genome (FIG. 1). One transduction reaction of 10^9 mycobacterial cells with phAE94 yielded all of the kanamycin resistant colonies used throughout this study. None of the retrieved colonies displayed a variant colony morphology from that usually observed in members of the *M. avium* complex. A large-scale sequencing strategy was employed to identify disrupted genes.

Identification of the Transposition Sites in *M. paratuberculosis* Mutants

Among the library of 5,060 mutants, 1,150 were analyzed using a high-throughput sequencing analysis employing a randomly primed PCR protocol that was successful in characterizing an *M. tuberculosis*-transposon library. These sequences were used to search *M. paratuberculosis* K-10 complete genome using BLASTN algorithm to identify the insertion site in 20% of the library. Generally, unique insertion sites (N=970) were identified, and almost 2/3 of the insertions occurring in predicted open reading frames (ORFs) while the rest of the insertions occurred in the intergenic regions (N=330) (Table 1).

TABLE 1

Percentage and number of unique insertions in a library of 5,060 mutants analyzed			
Insertion Sites	Number	No. of unique Insertions	% Unique*
ORF	714	640	89.6
Intergenic region	436	330	75.7
Total	1150	970	84.3

*indicates the percentage of insertions in unique sites within ORF or intergenic regions.

Among the 970 unique insertions within ORFs, only 288 of the predicted mycobacterial ORFs were disrupted at least once by the transposition of Tn5367 indicating that more than an insertion occurred multiple times in some genes. In fact, only 10.4% of disrupted ORFs showed more than one insertion per ORF indicating the presence of "hot spots" for transposition with Tn5367. Compared to insertions in ORFs, a higher rate by at least two times was observed when intergenic regions (24.3%) were examined (Table 2). Overall, the structure of the *M. paratuberculosis* mutant library was similar to that constructed in *M. tuberculosis*.

More scrutiny of the DNA sequences in both coding and intergenic regions revealed that regions most susceptible to transposon insertions are those with G+C content ranging from 50.5% to 60.5%, which is considerably lower than the average G+C content of the whole *M. paratuberculosis* (69.2%) (Table 2). Analysis of the flanking regions of Tn5367 site of insertion in genes with high frequency of transposition (N>4) identified areas of AT or TA repeats (e.g. TTT(T/A), AA(A/T) or TAA) as the most predominant sequences.

To illustrate the randomness of the Tn5367 transposition in *M. paratuberculosis* genome, the gene positions of all sequenced mutants were mapped to the genome sequence of *M. paratuberculosis* K10 (GenBank No. AE016958). Additionally, several mutants showed insertion into ORFs that have multiple copies in the genome (e.g. gene families or paralogous genes). These were excluded from further analysis.

As shown in FIG. 2, the transposition insertions were distributed in all parts of the genome without any apparent bias to a particular area. Overall, 1,128 mutants underwent the second level of bioinformatic analysis. FIG. 2 shows the

22

distribution of 1,128 transposon-insertion sites on the chromosome of *M. paratuberculosis* K-10 indicated by long bars on the outer-most circle. The inner two circles of short bars show predicted genes transcribed in sense or antisense directions.

TABLE 2

Characterization of <i>M. paratuberculosis</i> mutants with high insertion frequency (>10 insertions)					
	Genome Coordinates	Gene ID	No. of insertions	G + C %	Gene products*
Coding regions	1297579-1298913	MAP1235	43	55.88	Hypothetical protein
	1719957-1721030	MAP 1566	42	58.19	Hypothetical protein
	878808-880535	MAP 0856C	25	57.98	Hypothetical protein
	877826-878770	MAP 0855	25	59.57	Hypothetical protein
	4266449-4267747	MAP 3818	15	59.66	Hypothetical protein
	1295719-1296441	MAP 1233	13	50.48	Hypothetical protein
	1296587-1297387	MAP 1234	13	57.42	Hypothetical protein
	4803081-4803626	MAP 4327C	12	60.25	Hypothetical protein
	299412-300203	MAP 0282C	11	60.47	Hypothetical protein
	2380554-2381286	MAP2149c-MAP2150	97	54.3	Hypothetical proteins
Intergenic regions	1276333-1276722	MAP1216c-MAP1217c	44	52.9	LpqQ & hypothetical protein
	1997030-1997898	MAP1820-MAP1821c	26	54.01	Hypothetical proteins
	4455022-4458337	MAP3997c-MAP3998c	21	53.9	SerB and hypothetical protein
	1409338-1410190	MAP1318c-MAP1319	20	57.4	Adenylate cyclase
	2383052-2384295	MAP2151-MAP2152c	20	54.1	Hypothetical proteins
	300204-301106	MAP0282c-MAP0283c	17	58.2	Hypothetical proteins
	31518-32640	MAP0027-MAP0028c	13	57.8	Hypothetical proteins
	4263656-4264948	MAP3815-MAP3816	13	60.4	Hypothetical proteins
	4810959-4811624	MAP4333-MAP4334	11	56.7	Hypothetical proteins

*Gene products were described based on cluster of proteins analysis with at least 50% identity to other mycobacterial spp. For intergenic regions, the products of both flanking genes were listed.

To further analyze the expected phenotypes of the disrupted genes, the flanking sequences of each disrupted gene were examined, to determine their participation in transcriptional units such as operons. This analysis could reveal potential polar effect that could be observed in some mutants. Using the operon prediction algorithm (OPERON), approximately 124 (43.0%) of disrupted ORFs were identified as members of 113 putative operons (Table 3), indicating possible phenotypes related to disruption of function encoded by the whole operon and not just the disrupted gene. A total of 52 of the disrupted genes were within the last gene of an operon and were unlikely to affect the expression of other genes.

A total of 23 of the Tn5367 insertions were counted in several genes of the same 12 operons suggesting preference of transpositions throughout these sequences. For example, in the kdp operon (encoding putative potassium translocating proteins), 4 genes were disrupted among the 5 genes consti-

tuting this operon. Overall, sequence analysis of transposon junction sites identified disruption of a unique set of genes scattered all over the genome.

TABLE 3

Operon analysis of 288 ORFs disrupted by transposons in this study		
	Operon (%)	Not in operon (%)
Number	124 (43.0)	164 (56.9)
First gene	40 (32.3)	N/A*
Middle gene	32 (35.8)	N/A
Last gene	52 (41.9)	N/A

*N/A: Not applicable

Sequence Analysis of Disrupted Genes

A total of 288 genes represented by 970 mutants were identified as disrupted from the initial screening of the transposon mutant library constructed in *M. paratuberculosis*. Examining the potential functional contribution of each disrupted gene among different functional classes encoded in the completely sequenced genome of *M. paratuberculosis* K10 strain will better characterize their roles in infection. With the help of the Cluster of Orthologous Group website, disrupted genes were sorted into functional categories (Table 4). Six genes did not have a match in the COG functional category of *M. paratuberculosis* and consequently were analyzed using *M. tuberculosis* functional category. These genes are involved in different cellular processes such as lipid metabolism (desA1), cell wall biosynthesis (mmpS4) and several possible lipoproteins (lppP, lppJ, lppN) including a member of the PE family (PE6).

TABLE 4

List of functional categories of 288 disrupted genes that were identified				
Functional Category	Coding Sequences		Mutants Number	
	Number in genome	% in genome	Number mutant	% in genome
Translation	154	3.5	6	3.9
RNA processing and modification	1	0.02	0	0.0
Transcription	262	6.0	8	3.1
Replication, recombination and repair	179	4.1	13	7.3
Chromatin structure and dynamics	1	0.02	0	0.0
Cell cycle control, mitosis and meiosis	34	0.8	3	8.8
Defense mechanisms	46	1.1	5	10.9
Signal transduction mechanisms	112	2.6	6	5.4
Cell wall/membrane biogenesis	132	3.0	12	9.1
Cell motility	10	0.2	0	0.0
Intracellular trafficking and secretion	20	0.5	0	0.0
Posttranslational modification, protein turnover, chaperones	102	2.3	5	4.9
Energy production and conversion	277	6.4	10	3.6
Carbohydrate transport and metabolism	187	4.3	18	9.6
Amino acid transport and metabolism	246	5.7	16	6.5
Nucleotide transport and metabolism	67	1.5	2	3.0

TABLE 4-continued

List of functional categories of 288 disrupted genes that were identified				
Functional Category	Coding Sequences		Mutants Number	
	Number in genome	% in genome	Number mutant	% in genome
Coenzyme transport and metabolism	126	2.9	3	2.4
Lipid transport and metabolism	326	7.5	20	6.1
Inorganic ion transport and metabolism	174	4.0	9	5.2
Secondary metabolites biosynthesis, transport and catabolism	357	8.2	26	7.3
General function, prediction only	375	8.6	30	8.0
Unknown function	248	5.7	16	6.5
Unknown	914	21.0	80	8.8

Interestingly, genes involved in cell motility, intracellular trafficking and secretions were not represented in the mutants that were analyzed so far despite their comprising a substantial number of genes (N=30) (Table 4). However, for most functional groups, the percentage of disrupted genes ranged between 3-11% of the genes encoded within the *M. paratuberculosis* genome.

In most of the functional classes, the percentage of disrupted genes among mutants agreed with the percentage of particular functional class to the rest of the genome. Only 2 gene groups (bacterial defense mechanisms and cell cycling) were over-represented in the mutant library indicating potential sequence divergence from the high G+C content of the rest of the genome, which favorably agreed with the Tn5367 insertional bias discussed before.

Colonization of Transposon Mutants to Mice Organs

To identify novel virulence determinants in *M. paratuberculosis*, the mouse model of paratuberculosis was employed to characterize selected transposon mutants generated in this study. Bioinformatic analysis was used to identify genes with potential contribution to virulence. Genes were selected if information on their functional role was available, especially genes involved in cellular process believed to be necessary for survival inside the host or genes similar to known virulence factors in other bacteria (Table 5).

The screen for virulence determinants was designed to encompass mutations in a broad range of metabolic pathways to determine whether any could play an essential role for *M. paratuberculosis* persistence during the infection. Genes involved in carbohydrate metabolism (e.g. gcpE, impA), ion transport and metabolism (e.g. kdpC, trpE2) and cell wall biogenesis (e.g. mmpL10, umaA1) were chosen for further investigating in the mouse model of paratuberculosis, and respective mutants were tested in vivo. Also chosen were: a probable isocitrate lyase (aceAB), a gene involved in mycobactin/exocholin synthesis (mbtH2), a possible conserved lipoprotein (lppP), as well as putative transcriptional regulators (map0834c and map1634).

TABLE 5

Characterization of transposon mutants tested in the mouse model of paratuberculosis		
Gene	Insertion %*	Known molecular function
mmpL10	18.6	Conserved transmembrane transport protein
fprA	56.5	Adrenodoxin-oxidoreductase
papA2	12.1	Conserved polyketide synthase associated protein
gcpE	56.8	Isoprenoid biosynthesis, 4-hydroxy-3-methylbut-2-en-1-yl diphosphate synthase
papA3_1	65.2	Probable conserved polyketide synthase associated protein
kdpC	45.1	Probable Potassium-transporting ATPase C chain
umaA1	63.5	Possible mycolic acid synthase
pstA	3.8	Non-ribosomal binding peptide synthetase
fabG2_2	70.1	Putative oxidoreductase activity
trpE2	81.2	Probable anthranilate synthase component I
impA	52.0	Probable inositol-monophosphatase
cspB	63.8	Small cold shock protein
aceAB	95.5	Probable isocitrate lyase
mbtH2	64.6	mbtH_2 protein family, mycobactin/exocholins synthesis
lpqP	1.6	Possible conserved lipoprotein
ptrA	83.6	Transcriptional regulatory, putative two-component system regulator
map1634	88.8	Transcription factor activity
lipN**	deletion	Lipase, esterase protein

*insertion % indicates the percentage from start codon of gene.

**lipN mutant was generated by homologous recombination.

Before animal infection, the growth curve of all mutants in Middlebrook 7H9 broth supplemented with kanamycin was shown to be similar to that of the parent strain. However, most mutants reached an $OD_{600}=1.0$ at 35 days compared to 25 days for the ATCC19698, parent strain, which could be attributed to the presence of kanamycin in the growth media. Once mycobacterial strains reached $OD_{600}=1.0$, they were appropriately diluted and prepared for intraperitoneal (IP) inoculation of 10^7 - 10^8 CFU/mouse. In each case, the bacterial colonization and the nature of histopathology induced post-challenge were compared to the parent strain of *M. paratuberculosis* inoculated at similar infectious dose.

FIG. 3 shows colonization levels of variable *M. paratuberculosis* strains to mice organs. Groups of mice were infected via intraperitoneal injection (10^7 - 10^8 CFU/mouse) with the wild-type strain (ATCC19698) or one of 11 mutants. Colonization by only 8 mutants is shown in liver (A), spleen (B) and intestine (C) after 3, 6 and 12 weeks post infection. Bars represent the standard errors calculated from the mean of colony counts estimated from organs at different times post infection.

All challenged mice were monitored for 12 weeks post infection with tissue sampling at 3, 6 and 12 weeks post infection. For samples collected at 3 weeks post-infection, only the strains with a disruption in gcpE or kdpC genes displayed significantly ($p<0.05$) lower colonization levels compared to the parent strain (FIG. 3), especially in the primary target of *M. paratuberculosis*, the intestine. Some of the mutants (gcpE and kdpC) displayed a significant reduction in the intestinal colony counts starting from 3 weeks post infection and throughout the experiment. At 6 weeks post infection, both papA2 and pstA mutants showed significant colony reduction in the intestine that was maintained in the later time point. At 12 weeks post infection, umaA1, fabG2_2, and impA genes displayed significantly decreased colonization in the intestine ($p<0.05$) with a reduction of at least 2 logs (FIG. 3C). Colonization levels of the spleen did not show a significant change while levels in the liver and intestine were vari-

able between mutants and wild-type and therefore, they were the most informative organs (FIG. 3).

The four mutants mmpL10, fprA, papA3_1, and trpE2 showed a 10-fold reduction in mycobacterial levels at least in one examined organ by 12 weeks post infection although, this reduction was not statistically significant ($p>0.05$).

Additional mutants with colonization levels significantly lower in both intestine and liver were identified. Shown in FIG. 4 are data obtained using attenuated mutants with disruption in one of aceAB, mbtH2, lpqP, map0834c, cspB, lipN, or map1634 genes. The graph in FIG. 4A depicts liver colonization of BALB/c mice following infection with 10^8 CFU/animal of *M. paratuberculosis* mutants compared to the wild type strain ATCC19698. IP injection was used as a method for infection. Colonization levels in the liver over 3, 6, and 12 weeks post infection were monitored and are shown in FIG. 4A. The graph in FIG. 4B depicts intestinal colonization of BALB/c mice following infection with 10^8 CFU/animal of *M. paratuberculosis* mutants compared to the wild type strain ATCC19698. IP injection was used as a method for infection. Colonization levels in the intestine over 3, 6, and 12 weeks post infection were monitored and are shown in FIG. 4B.

Histopathology of Mice Infected with Transposon Mutants
All animal groups infected with mutants or the parent strain displayed a granulomatous inflammatory reaction consistent with infection with *M. paratuberculosis* using the mouse model of paratuberculosis. Liver sections were the most reflective organ for paratuberculosis where a typical granulomatous response was found. It was exhibited as aggregation of lymphocytes surrounded with a thin layer of fibrous connective tissues.

FIG. 5 shows histopathological data from liver of mice infected with *M. paratuberculosis* strains as outlined in FIG. 3. At 3, 6 and 12 weeks post infection, mice were sacrificed and liver, spleen, and intestine were processed for histopathological examination. Liver sections stained with H&E with arrows indicating granulomatous inflammatory responses were shown in FIG. 4 of U.S. Provisional Patent Application Ser. No. 60/749,128, incorporated herein by reference. FIG. 5 is a chart showing the inflammatory scores of all mice groups.

Granuloma formation was apparent in animals infected with ATCC19698 strain and some mutants such as Δ mmpL10. Both the size and number of granulomas were increased over time indicating the progression of the disease. During early times of infection (3 and 6 weeks sampling), most mutants displayed only lymphocytic inflammatory responses while the formation of granulomas was observed only at the late time (12 weeks samples). Additionally, the severity of inflammation reached level 3 (out of 5) at 12 weeks post-infection for mice infected with ATCC19869 while in the group infected with mutants such as Δ gcpE and Δ kdpC, the granulomatous response was lower (ranged between levels 1 and 2).

When mice infected with Δ mmpL10 were examined, the lymphocyte aggregates were larger in size and were well-separated by fibrous tissues compared to the granuloma formed in mice infected with the ATCC19698. On the other hand, some mutants (e.g. Δ gcpE, Δ impA) began with relatively minor lesions and remained at this level as time progressed while others (Δ pap3_1, fabG2_2) started with mild lesions and progressively increased in severity over time.

A third group of mutants (Δ fprA, Δ kdpC) began with a similar level of response to that of the parent strain and continue to be severely affected until the end of the sampling time.

Generally, by combining the histopathology and colonization data it was possible to assess the overall virulence of the examined mutants and classify disrupted genes into 3 classes.

In Class I (early growth mutants), the disruption of genes (e.g. *gcpE*, *KdpC*) generated mutants that are not able to multiply efficiently in mice tissues and therefore, a modest level of lesions was generated and their colonization levels were significantly lower than that of wild-type. In Class II (tissue specific mutants), levels of bacterial colonization were significantly reduced in only specific tissues such as *umaA1* for liver and *papA2* in the intestine at 6 weeks samples. No characteristic pathology of this group could be delineated since only liver sections were reflective of the paratuberculosis using the mouse model employed in this study. In Class III (persistence mutants), levels of colonization were maintained unchanged in the first 6 weeks and then reduced significantly at later times (e.g. *fabG2_2* and *impA*). The lesions formed in animals infected with Class III mutants showed a similar pattern of lesion progression to those of animals infected with the parent strain.

Generally, there was an inverse relationship between granuloma formation scores and mycobacterial colonization levels of mutants for samples collected at 12 weeks post infection. The decline of *M. paratuberculosis* levels could be attributed to the initiation of a strong immune response represented by an increase of granuloma formation. However, in the case of animals infected with Δ *pstA* and Δ *impA*, the decline of colonization level was consistent with the reduction in granuloma scores.

Overall, large scale characterization of mutant libraries for virulence determinants is shown to be possible, especially when the genome sequence of a given genome is known. The employed approach can be applied in other bacterial systems where there is little information available on pathogen virulence determinants.

Histopathological analyses of mice infected with the attenuated *M. paratuberculosis* mutants *aceAB*, *mbtH2*, *lpqP*, *map0834c*, *cspB*, *lipN*, or *map1634* showed a decrease in granuloma formation in the liver, compared to the mice infected with the wild type *M. paratuberculosis* strain ATCC19698.

Characterization of Transposon Mutants

The list of diagnostic targets, i.e., potential virulence determinants disclosed here includes the *gcpE* gene encoding a product that controls a terminal step of isoprenoid biosynthesis via the mevalonate independent 2-C-methyl-D-erythritol-4-phosphate (MEP) pathway. Because of its conserved nature and divergence from mammalian counterpart, *gcpE* and its products are considered a suitable target for drug development.

Another diagnostic target, i.e., potential virulence gene, is *pstA*, which encodes non-ribosomal peptide synthetase in *M. tuberculosis* with a role in glycopeptidolipids (GPLs) synthesis. The GPLs is a class of species-specific mycobacterial lipids and major constituents of the cell envelopes of many non-tuberculous mycobacteria as well, such as *M. smegmatis*.

Disruption of *umaA1* also resulted in lower colonization levels in all organs examined at 6 weeks post infection and forward.

Additional potential virulence determinants include *papA3_1* and *papA2*, genes that are members of the polyketide synthase associated proteins family of highly conserved genes. Members of the *pap* family encode virulence-enhancing lipids. Nonetheless, these two mutants displayed different attenuation phenotypes. The *papA2* mutant showed significantly lower CFU than the *papA3_1* mutant.

The *kdpC* gene encodes an inducible high affinity potassium uptake system. The *kdpC* mutant was significantly reduced mostly in the intestinal tissue at early and late stages of infection.

The *impA* mutant showed significantly reduced levels at late times of infection indicating that *impA* may possibly play a role in *M. paratuberculosis* entry into the persistence stage of the infection.

The *aceAB* mutant showed significantly reduced levels at late times of infection indicating that *aceAB* may possibly play a role in *M. paratuberculosis* entry into the persistence stage of the infection. Deletion of a homologue of the gene in *M. tuberculosis* rendered this mutant attenuated.

The *mbtH2* mutant showed significantly reduced levels at early times of infection indicating that *mbtH2* may possibly play a role in *M. paratuberculosis* entry into the intestinal cells or survival in macrophage during early infection. This gene was induced during animal infection using DNA microarrays conducted in the inventor's laboratory.

The *lpqP* mutant showed significantly reduced levels at late times of infection indicating that *lpqP* may possibly play a role in *M. paratuberculosis* entry into the persistence stage of the infection.

The *prpA* mutant showed significantly reduced levels at late times of infection indicating that *prpA* may possibly play a role in *M. paratuberculosis* entry into the persistence stage of the infection. The *prpA* homologue in *M. tuberculosis* is two-component transcriptional regulator. This gene was induced at low pH using DNA microarrays conducted in the inventor's laboratory.

The *map1634* mutant showed significantly reduced levels at late times of infection indicating that *map1634* may possibly play a role in *M. paratuberculosis* entry into the persistence stage of the infection.

The *lipN* mutant showed significantly reduced levels at mid and late times of infection indicating that *lipN* may play an important role in *M. paratuberculosis* during early and persistent stages of the infection. *LipN* encodes a lipase which could be important degrading fatty acids. This gene was induced in cow samples using DNA microarrays conducted in the inventor's laboratory.

Example 2

Bacterial Strains

Mycobacterial isolates (N=34) were collected from different human and domesticated or wildlife animal specimens representing different geographical regions within the USA (Table 6). *Mycobacterium avium* subsp. *paratuberculosis* K10 strain, *M. avium* subsp. *avium* strain 104 (*M. avium* 104) and *M. intracellulare* were obtained from Raul Barletta (University of Nebraska). *M. paratuberculosis* ATCC19698 and other animal isolates were obtained from the John's Testing Center, University of Wisconsin-Madison, while the *M. paratuberculosis* human isolates were obtained from Saleh Naser (University of Central Florida). All strains were grown in Middlebrook 7H9 broth supplemented with 0.5% glycerol, 0.05% Tween 80 and 10% ADC (2% glucose, 5% BSA fraction V, and 0.85% NaCl) at 37° C. For *M. paratuberculosis* strains, 2 µg/ml of mycobactin-J (Allied Monitor, Fayette, Mo.) also was added for optimal growth.

TABLE 6

<i>Mycobacterium</i> strains tested in Example 2 of the present invention				
Species	Strain	Host	Sample origin	Location
<i>M. avium</i> subsp. <i>paratuberculosis</i>	K10	Cow	Feces	Wisconsin
	ATCC 19698	Cow	Feces	Unknown
	JTC33666	Turkomen markhor (Goat)	Feces	California
	JTC33770	Cow	Feces	Wisconsin
	CW303	Cow	Feces	Wisconsin
	1B	Human	Ileum	Florida
	3B	Human	Ileum	Florida
	4B	Human	Ileum	Florida
	5B	Human	Ileum	Florida
	DT3	British red deer	Feces	Unknown
	DT9	African Eland	Feces	Unknown
	DT12	Chinese Reeve's muntjac (Deer)	Ileum	Unknown
	DT19	White rhino	Feces	Unknown
	JTC1281	Oryx	Lymph Node	Florida
	JTC1282	Cow	Lymph Node	Wisconsin
	JTC1283	Cow	Feces	Georgia
	JTC1285	Goat	Feces	Virginia
	JTC1286	Cow	Ileum	Wisconsin
	104	Human	Blood	Unknown
	T93	Cow	Feces	Texas
	T99	Cow	Feces	Texas
	T100	Cow	Feces	Texas
	DT30	Angolan springbok	Feces	Unknown
	DT44	Fornosan Reeve's muntjac (Deer)	Lymph Node	Unknown
	DT78	Water buffalo	Ileum	Unknown
	DT84	Lowland wisent	Lymph Node	Unknown
<i>M. avium</i> subsp. <i>avium</i>	DT247	Cuvier's gazelle	Lymph Node	Unknown
	JTC956	Ankole	Feces	Florida
	JTC981	Bongo	Feces	Florida
	JTC982	Nyala	Feces	Florida
	JTC1161	Cow	Feces	Florida
	JTC1262	Bison	Lymph Node	Montana
	JTC33793	Dama gazelle	Feces	Indiana
	mc ² 76	Human	Sputum	Unknown
<i>M. intracellulare</i>				

Microarray Design

Oligonucleotide microarrays were synthesized in situ on glass slides using a maskless array synthesizer. Probe sequences were chosen from the complete the genome sequence of *M. avium* 104. Sequence data of *M. avium* 104 strain was obtained from The Institute for Genomic Research. Open reading frames (ORFs) were predicted using GeneMark software. For every ORF, 18 pairs of 24-mer sequences were selected as probes. Each pair of probes consists of a perfect match (PM) probe, along with a mismatch (MM) probe with mutations at the 6th and 12th positions of the corresponding PM probes. A total of ~185,000 unique probe sequences were synthesized on derivatized glass slides by NimbleGen Systems (Madison, Wis.).

Genomic DNA Extraction and Labeling

Genomic DNA was extracted using a modified CTAB-based protocol followed by two rounds of ethanol precipitation. For each hybridization, 10 µg of genomic DNA was digested with 0.5 U of RQ1 DNase (Promega, Madison, Wis.) until the fragmented DNA was in the range of 50-200 bp (examined on a 2% agarose gel). The reaction was stopped by adding 5 µl of DNase stop solution and incubating at 90° C. for 5 minutes. Digested DNA was purified using YM-10 microfilters (Millipore, Billerica, Mass.).

Genomic DNA hybridizations were prepared by an end-labeling reaction. Biotin was added to purified mycobacterial DNA fragments (10 µg) using terminal deoxynucleotide transferase in the presence of 1 µM of biotin-N6-ddATP at 37° C. for 1 hr. Before hybridization, biotin-labeled gDNA was heated to 95° C. for 5 minutes, followed by 45° C. for 5 minutes, and centrifuged at 14,000 rpm for 10 minutes before adding to the microarray slide.

After microarray hybridization for 12-16 hrs, slides were washed in non-stringent (6×SSPE and 0.01% Tween-20) and stringent (100 mM MES, 0.1 M NaCl, and 0.01% Tween 20) buffers for 5 min each, followed by fluorescent detection by adding Cy3 streptavidin (Amersham Biosciences Corp., Piscataway, N.J.). Washed microarray slides were dried by argon gas and scanned with an Axon GenPix 4000B (Axon Instrument, Union City, Calif.) laser scanner at 5 µm resolution. Replicate microarrays were hybridized for every genome tested. Two hybridizations of the same genomic DNA with high reproducibility (correlation coefficient>0.9) were allowed for downstream analysis.

Data Analysis and Prediction of Genomic Deletions

The images of scanned microarray slides were analyzed using specialized software (NimbleScan) developed by NimbleGen Systems. The average signal intensity of a MM probe was subtracted from that of the corresponding PM probe. The median value of all PM-MM intensities for an ORF was used to represent the signal intensity for the ORF. The median intensities value for each slide was normalized by multiplying each signal by a scaling factor that was 1000 divided by the average of all median intensities for that array.

To compare hybridization signals generated from each of the genomes to that of *M. avium* 104, the normalized data from replicate hybridizations were exported to R language program with the EBarrays package version 1.1, which employs a Bayesian statistical model for pair-wise genomic comparisons using a log-normal-normal model. Genes with the probability of differential expression larger than 0.5 were considered significantly different between the genomes of *M. avium* and *M. paratuberculosis*.

The hybridization signals corresponding to each gene of all investigated genomes were plotted according to genomic location of *M. avium* 104 strain using the GenVision software (DNASTar Inc., Madison, Wis.). The same data set was also analyzed by MultiExperiment Viewer 3.0 to identify common cluster patterns among mycobacterial isolates.

Microarray Analysis of *M. avium* and *M. paratuberculosis* Genomes

Genomic rearrangements among *M. avium* and *M. paratuberculosis* isolated from variable hosts were investigated, to identify diagnostic targets for microbial infection. The analysis began using 5 mycobacterial isolates employing DNA microarrays and was expanded to include an additional 29 isolates employing a more affordable technology of PCR followed by direct sequencing. All of the isolates were collected from human and domesticated or wildlife animal sources and had been previously identified at the time of isolation using standard culturing techniques for *M. avium*.

and *M. paratuberculosis*. The identity of each isolate was confirmed further by acid-fast staining and positive PCR amplification of IS900 sequences from all *M. paratuberculosis*. Additionally, the growth of all *M. paratuberculosis* isolates were mycobactin-J dependent while all *M. avium* isolates were not.

Before starting the microarray analysis, an hsp65 PCR typing protocol was performed to ensure the identity of each isolate. The PCR typing protocol agreed with earlier characterization of all mycobacterial isolates used throughout this study. FIG. 5A of U.S. Provisional Patent Application Ser. No. 60/749,128, incorporated by reference, depicts the PCR confirmation of the identity of the examined genomes.

To investigate the extent of variation among *M. avium* and *M. paratuberculosis* on a genome-wide scale, oligonucleotide microarrays were designed from the *M. avium* 104 strain genome sequence. The GeneMark algorithm was used to predict potential ORFs in the raw sequence of *M. avium* genome obtained from TIGR. A total of 4987 ORFs were predicted for *M. avium* compared to 4350 ORFs predicted in *M. paratuberculosis*. Relaxed criteria for assigning ORFs were chosen (at least 100 bp in length with a maximal permitted overlap of 30 bases between ORFs) to use a comprehensive representation of the genome to construct DNA microarrays.

Similar to other bacterial genomes, the average ORF length was ~1 Kb. Using the ASAP comparative genomic software suite, the ORFs shared by *M. paratuberculosis* and *M. avium* had an average percent identity of 98%, a result corroborated by others. BLAST analysis of the ORFs from both genomes show that about 65% (N=2557) of the genes have a significant match (E<10⁻¹⁰) in the other genome.

To test the reliability of genomic DNA extraction protocols and microarray hybridizations, the signal intensities of replicate hybridizations of the same mycobacterial genomic DNA were compared using scatter plots. ORFs with positive hybridization signals in at least 10 probe pairs were normalized and used for downstream analysis to ensure the inclusion of only ORFs with reliable signals. In all replicates, independently isolated hybridized samples of gDNA had high correlation coefficients (r>0.9).

To investigate the genomic relatedness among isolates compared to the *M. avium* 104 strain, a hierarchical cluster analysis was used to assess the similarity of the hybridization signals among isolates on a genome-wide level. FIG. 5C of U.S. Provisional Patent Application Ser. No. 60/749,128, incorporated by reference, shows a dendrogram displaying the overall genomic hybridization signals generated from biological replicates of different mycobacterial isolates from animal or human (HU) sources.

Within the *M. paratuberculosis* cluster, the human and the clinical animal isolates were highly similar to each other than to the ATCC19698 reference strain, implying a closer relatedness between human and clinical isolate of *M. paratuberculosis*. Interestingly, despite the high degree of similarity between genes shared among isolates, hundreds of genes appeared to be missing from different genomes relative to *M. avium* genome. Most of the genes were found in clusters in the *M. avium* 104 genome, the reference strain used for designing the microarray chip. Consequently, regions absent in *M. avium* 104 but present in other genomes could not be identified in this analysis.

PCR Verification and Sequence Analysis

To confirm the results predicted by microarray hybridizations, a 3-primer PCR protocol was used to amplify the regions flanking predicted genomic islands. For every island, one pair of primers (F—forward and R1—reverse 1) was

designed upstream of the target region and a third primer (R2—reverse 2) was designed downstream of the same region. The primers were designed so that expected lengths of the products were less than 1.5 Kb between F and R1 and less than 3 Kb between F and R2 when amplified from the genomes with the deleted island. Each PCR contained 1 M betaine, 50 mM potassium glutamate, 10 mM Tris-HCl pH 8.8, 0.1% of Triton X-100, 2 mM of magnesium chloride, 0.2 mM dNTPs, 0.5 μ M of each primer, 1 U Taq DNA polymerase and 15 ng genomic DNA. The PCR cycling condition was 94° C. for 5 minutes, followed by 30 cycles of 94° C. for 1 minute, 59° C. for 1 minute and 72° C. for 3 minutes.

All PCR products were examined using 1.5% agarose gels and stained with ethidium bromide. To further confirm sequence deletions, amplicons flanking deleted regions were sequenced using standard BigDye® Terminator v3.1 (Applied Biosystems, Foster City, Calif.) and compared to the genome sequence of *M. paratuberculosis* or *M. avium* using BLAST alignments.

Large Genomic Deletions among *M. avium* and *M. paratuberculosis* Isolates

To better analyze the hybridization signals generated from examined genomes, a Bayesian statistical principle (EBarrays package) was used to compare the hybridization signals generated from different isolates relative to the signals generated from *M. avium* 104 genome. The Bayesian analysis estimates the likelihood of observed differences in ORF signals for each gene between each isolate and the *M. avium* 104 reference strain.

FIG. 6A depicts a genome map based on *M. avium* sequence displaying GIs deleted in the examined strains as predicted by DNA microarrays. Inner circles denote the microarray hybridization signals for each examined genome (see legend in center). The outermost dark boxes denote the location of all GIs associated with *M. avium*. A large number of differences were seen among isolates, including many ORFs scattered throughout the genome.

PCR and sequencing were used to confirm deletions identified by microarrays. FIG. 6B depicts a diagram illustrating the PCR and sequence-based strategy implemented to verify the genomic deletions. Three primers for each island were designed including a forward (F) and 2 reverse primers. When regions included 3 or more consecutive ORFs, they were defined as a genomic island (GI) regardless of the size. Applying such criterion for genomic islands (GIs), 24 islands were present in *M. avium* 104 but absent from all *M. paratuberculosis* isolates, regardless of the source of the *M. paratuberculosis* isolates (animal or human). The GIs ranged in size from 3 to 196 Kb (Table 7) with a total of 846 Kb encoding 759 ORFs. Interestingly, a clinical strain of *M. avium* (JTC981) was also missing 7 GIs (nearly 518 Kb) in common with all *M. paratuberculosis* isolates, in addition to the partial absence of 5 other GIs. This variability indicated a wide-spectrum of genomic diversity among *M. avium* strains that was not evident among *M. paratuberculosis* isolates.

To confirm the absence of GI regions from isolates, a strategy based on PCR amplification of the flanking regions of each GI was used, followed by sequence analysis to confirm the missing elements. Because the size of most of the genomic island regions exceeds the length of the amplification capability of a typical PCR reaction, 3 primers for each island were designed, including one forward and 2 reverse primers (FIG. 6B). This strategy was successfully applied on 21 genomic islands, while amplification from the rest of the islands (N=3) was not possible due to extensive genomic rearrangements.

FIG. 7 depicts the synteny of *M. avium* and *M. paratuberculosis* genomes.

PCR confirmation of genomic deletions was performed. For example, amplicons from *M. avium*-specific islands #5, 8, 11, 18 and 20 were obtained using DNA templates from 5 different isolates of *M. avium*. Additionally, PCR analysis of the distribution of *M. paratuberculosis*-specific island #1 was performed within 21 clinical isolates of *M. avium* and *M. paratuberculosis*. Electrophoresed DNA samples showed PCR confirmations of the genomic deletions.

Overall, the PCR and sequencing verified the GI content as predicted by comparative genomic hybridizations (Table 7). The success of this strategy in identifying island deletions provided a protocol to examine several clinical isolates that could not be otherwise analyzed by costly DNA microarrays.

fairly recent ancestor. Looking for consecutive ORFs from *M. paratuberculosis* that do not have a BLAST match in *M. avium* identified sets of ORFs representing 18 GIs comprising 240 Kb that are present only in *M. paratuberculosis* genome (Table 8).

Genes encoded within *M. avium* and *M. paratuberculosis* specific islands were analyzed by BLASTP algorithm against the GenPept database (Oct. 19, 2004 release) to identify their potential functions. The BLAST results allowed the assignment of signature features to each island. As detailed in Tables 8 and 9, with the presence of a large number of ORFs encoding mobile genetic elements (e.g. insertion sequences and prophages), several ORFs encode transcriptional regulatory elements, especially from TetR-family of regulators. The polymorphism in TetR regulators could be attributed to their sequences allowing them to be amenable for rearrangements.

TABLE 7

List of genomic regions that displayed different hybridization signals using DNA microarrays designed from the genome of <i>M. avium</i> 104 strain							
Island Number	Start (bp) ^a	End (bp) ^a	<i>M. parat.</i> K10 ^b	<i>M. parat.</i> 19698	<i>M. parat.</i> human	<i>M. avium</i> JTC981	PCR and sequence confirmation ^c
1	254,394	294,226	—	—	—	—	Yes
2	461,414	492,800	—	—	—	—	Yes
3	666,033	675,725	—	—	—	—	Yes
4	747,095	794,450	—	—	—	—	Yes
5	1,421,722	1,439,626	—	—	—	+	Yes
6	1,444,205	1,463,365	—	—	—	+	Yes
7	1,795,281	1,991,691	—	—	—	+/-	Yes
8	2,097,907	2,100,883	—	—	—	—	Yes
9	2,220,320	2,241,163	—	—	—	+/-	Yes
10	2,259,120	2,271,610	—	—	—	—	Yes
11	2,462,693	2,466,285	—	—	—	+	Yes
12	2,549,555	2,730,999	—	—	—	—	ND
13	2,815,625	2,821,149	—	—	—	+	Yes
14	3,008,716	3,036,980	—	—	—	+	Yes
15	3,214,820	3,219,550	—	—	—	+	ND
16	3,340,393	3,384,549	—	—	—	+	Yes
17	3,392,586	3,413,804	—	—	—	+	ND
18	3,523,417	3,527,334	—	—	—	+/-	Yes
19	3,670,518	3,675,686	—	—	—	+	Yes
20	3,917,752	3,939,034	—	—	—	+/-	Yes
21	4,254,594	4,261,488	—	—	—	+/-	Yes
22	5,122,371	5,132,301	—	—	—	+	Yes
23	5,174,641	5,270,187	—	—	—	+	Yes
24	5,378,903	5,395,102	—	—	—	+	Yes

^aCoordinates of start and end of island based on the genome sequence of *M. avium* strain 104.

^b+ or — denotes presence or absence of genomic regions in examined genomes while +/- denotes incomplete deletion.

^cND—not done.

Bioinformatic Analysis of Genomic Islands

Pair-wise BLAST analysis of the genome sequences of *M. avium* 104 and *M. paratuberculosis* K10 was used to further refine the ability to detect genomic rearrangements, especially for regions present in *M. paratuberculosis* K10 genome but deleted from *M. avium* 104 genome. The pair-wise comparison allowed to better analyze the flanking sequences for each GI and to characterize the mechanism of genomic rearrangements among examined strains.

BLAST analysis (E scores >0.001 and <25% sequence alignment between ORFs) correctly identified the deleted GIs where ORFs of *M. avium* were missing in *M. paratuberculosis* detected by using the comparative genomic hybridization protocol. A large proportion of ORFs in each genome (>75%) are likely orthologous (>25% sequence alignment of the ORF length and >90% sequence identity at nucleotide level). This high degree of similarity between orthologues indicates a

Alternatively, it is possible that the bacteria are able to differentially acquire specific groups of genes suitable for a particular microenvironment.

Further analysis of the GIs identified islands in both *M. avium* and *M. paratuberculosis* (such as MAV-7, MAV-12 and MAP-13) encoding different operons of the mce (mammalian cell entry) sequences that were shown to participate in the pathogenesis of *M. tuberculosis*. Another island (MAV-17) encodes the drrAB operon for antibiotic resistance, which is a well-documented problem for treating *M. avium* infection in HIV patients. The GC % of the majority of *M. paratuberculosis* specific islands (11/18) was at least 5% less than the average GC % of the *M. paratuberculosis* genome (69%) compared to only 3 GIs (out of 24) specific for *M. avium* genome (Table 9) with lower than average GC %.

TABLE 8

<i>M. paratuberculosis</i> -specific (MAP) genomic islands deleted in <i>M. avium</i> genome					
Island Number	No. of ORFs	GC %	Island Type	Size (bp)	Signature Features
MAP-1	17	63.90	I	19,343	Transposition and TetR-family transcriptional regulator genes
MAP-2	3	60.43	I	3,858	Conserved hypothetical proteins
MAP-3	3	66.16	I	2,915	Formate dehydrogenase alpha subunit
MAP-4	17	60.66	I	16,681	Transposition, unknown genes and a possible prophage
MAP-5	12	69.56	I	14,191	Transposition and oxidoreductase genes, PPE family domain protein
MAP-6	6	57.73	II	8,971	Variable genes such as <i>drnC</i>
MAP-7	6	67.26	II	6,914	Transcriptional regulator <i>psrA</i> and biosynthesis genes
MAP-8	8	61.59	II	7,915	TetR-family transcriptional regulator and unknown genes
MAP-9	10	65.49	II	11,202	Transposition, metabolic and TetR-family transcriptional regulator genes
MAP-10	3	66.68	II	2,993	Biosynthesis of cofactors, prosthetic groups, and carriers transcriptional regulator, TetR family domain protein
MAP-11	4	62.89	I	2,989	Serine/threonine protein kinase and glyoxalase genes
MAP-12	11	61.08	I	11,977	Transposition, iron metabolism genes and a prophage
MAP-13	19	66.01	II	19,977	TetR-family transcript, regulator and <i>mce</i> family proteins
MAP-14	19	65.76	II	19,315	Possible prophage and unknown proteins
MAP-15	3	62.93	I	4,143	Unknown proteins and a prophage function genes
MAP-16	56	64.32	I	79,790	Transposition and iron regulatory genes
MAP-17	5	61.60	I	3,655	Unknown proteins and a multi-copy phage resistance gene
MAP-18	3	60.36	I	3,512	Hypothetical proteins
Total	204			239,969	

TABLE 9

Characteristics of <i>M. avium</i> -specific (MAV) genomic islands					
Island Number	No. of ORFs	GC %	Island Type	Size (bp)	Signature Features
MAV-1	38	68.93	I	39,833	Eukaryotic genes with an integrase gene
MAV-2	32	65.87	I	31,387	Transposition and <i>M. tuberculosis</i> genes
MAV-3	10	63.34	I	9,693	Insertion sequence and <i>M. tuberculosis</i> or <i>M. avium</i> genes
MAV-4	53	66.83	I	47,356	PPE family and eukaryotic genes
MAV-5	16	64.10	I	17,905	Transposition and insertion sequences genes
MAV-6	23	68.80	I	19,161	Transposition, transcript, regulator and heavy metal resistance genes
MAV-7	187	65.50	II	196,411	Transposition, transcript, regulators, cell entry, iron regulation genes
MAV-8	3	65.18	I	2,977	Transposition and transcriptional regulator genes
MAV-9	15	62.43	I	20,844	Transposition and type III restriction system endonuclease genes
MAV-10	12	63.87	I	12,491	Transposition genes
MAV-11	5	65.45	I	3,593	Reductases and hypothetical proteins
MAV-12	168	65.05	II	181,445	Transposition, transcriptional regulators and cell entry genes
MAV-13	7	67.78	II	5,525	Transcriptional regulator
MAV-14	26	67.32	I	28,265	Transposition and <i>M. tuberculosis</i> genes

TABLE 9-continued

Characteristics of <i>M. avium</i> -specific (MAV) genomic islands					
Island Number	No. of ORFs	GC %	Island Type	Size (bp)	Signature Features
MAV-15	3	64.12	II	4,731	<i>Streptomyces</i> and <i>M. leprae</i> genes
MAV-16	6	69.64	I	44,157	Transposition and Pst genes
MAV-17	20	65.23	II	21,219	Transposition and drrAB genes (antibiotic resistance)
MAV-18	4	68.13	I	3,918	Transcriptional regulator and <i>Streptomyces</i> genes
MAV-19	4	65.30	I	5,169	Transposition genes
MAV-20	15	63.93	I	21,283	Transposition, transcriptional regulator and membrane-protein genes of <i>M. tuberculosis</i>
MAV-21	8	65.93	I	6,895	Transposition and antigen genes
MAV-22	9	67.71	I	9,931	Transcriptional regulator and metalloprotease genes
MAV-23	77	64.08	I	95,547	Transposition, transcript, regulators, secreted proteins, cell entry genes
MAV-24	18	70.25	I	16,200	Hypothetical and unknown proteins from <i>M. tuberculosis</i> and <i>Streptomyces</i>
Total	759			845,936	

Genomic Deletions Among Field Isolates of *M. avium*

Microarrays and PCR analysis of 5 mycobacterial isolates identified the presence of variable GIs between *M. avium* and *M. paratuberculosis* genomes. To analyze the extent of such variations among clinical isolates circulating in both human and animal populations, PCR and a sequencing-based strategy were used to examine 28 additional *M. avium* and *M. paratuberculosis* isolates collected from different geographical locations within the USA (Table 6). An additional isolate of *M. intracellulare* was included as a representative strain that belongs to the MAC group but not a subspecies of *M. avium*.

For PCR amplification, GIs spatially scattered throughout the *M. avium* and *M. paratuberculosis* genomes were examined (Tables 10, 11) to identify any potential rearrangements in all quarters of the genome. Because of the wide-spectrum diversity observed among *M. avium* genomes, 4 GIs (MAV-3, 11, 21 and 23) were chosen to assess genomic rearrangements in clinical isolates. Because of the limited diversity observed among *M. paratuberculosis* genomes, a total of 6 *M. paratuberculosis*-specific GIs (MAP-1, 3, 5, 12, 16 and 17) were chosen for testing genomic rearrangements. As suggested from the initial comparative genomic hybridizations, clinical isolates of *M. paratuberculosis* showed a limited diversity in the existence of *M. avium*-specific islands (DT9 clinical isolate from a red deer) indicating the clonal nature of this organism (Table 10).

To the contrary, *M. avium* isolates showed a different profile from both *M. avium* 104 and *M. avium* JTC981 indicating extensive variability within *M. avium* isolates. A similar pattern of genomic rearrangements was observed when *M. paratuberculosis*-specific GIs were analyzed using *M. avium* and *M. paratuberculosis* isolates (Table 11). Most of the *M. paratuberculosis* clinical isolates with deleted GIs were from wildlife animals suggesting that strains circulating in wildlife animals could provide a potential source for genomic rearrangements in *M. paratuberculosis*.

TABLE 10

PCR identification of selected MAV-island regions from 29 clinical isolates of *M. paratuberculosis* and *M. avium* collected from different states

Clinical Isolate	Subspecies	Genomic island			
		MAV-3	MAV-11	MAV-21	MAV-23
JTC33666	<i>M. paratuberculosis</i>	-	-	-	-
JTC33770	<i>M. paratuberculosis</i>	-	-	-	-
CW303	<i>M. paratuberculosis</i>	-	-	-	-
1B	<i>M. paratuberculosis</i>	-	-	-	-
3B	<i>M. paratuberculosis</i>	-	-	-	-
4B	<i>M. paratuberculosis</i>	-	-	-	-
5B	<i>M. paratuberculosis</i>	-	-	-	-
DT3	<i>M. paratuberculosis</i>	-	-	-	-
DT9	<i>M. paratuberculosis</i>	+	N/A	-	-
DT12	<i>M. paratuberculosis</i>	-	-	-	-
DT19	<i>M. paratuberculosis</i>	-	-	-	-
JTC1281	<i>M. paratuberculosis</i>	-	-	-	-
JTC1282	<i>M. paratuberculosis</i>	-	-	-	-
JTC1283	<i>M. paratuberculosis</i>	-	-	-	-
JTC1285	<i>M. paratuberculosis</i>	-	-	-	-
JTC1286	<i>M. paratuberculosis</i>	-	-	-	-
T93	<i>M. avium</i>	+	-	-	-
T99	<i>M. avium</i>	+	-	-	-
T100	<i>M. avium</i>	+	+	-	-
DT30	<i>M. avium</i>	-	+	+	+
DT44	<i>M. avium</i>	-	+	+	+
DT78	<i>M. avium</i>	-	+	+	+
DT84	<i>M. avium</i>	-	+	-	+
DT247	<i>M. avium</i>	-	+	+	+
JTC956	<i>M. avium</i>	N/A	N/A	N/A	-
JTC982	<i>M. avium</i>	N/A	+	N/A	+
JTC1161	<i>M. avium</i>	+	+	-	-
JTC1262	<i>M. avium</i>	+	-	-	-
JTC33793	<i>M. avium</i>	+	+	+	+

Symbols (+ or -) denote presence or absence of genomic regions; N/A denotes no amplification of DNA fragments.

Combined with the hierarchical cluster analysis employed on the whole genome hybridizations, PCR and sequence analyses provided more evidence that genomic diversity is quite extensive among *M. avium* strains but much less limited in *M. paratuberculosis*.

Large DNA Fragment Inversions Within the Genomes of *M. avium* Subspecies.

Because of the high similarity among the genomes of *M. paratuberculosis* and *M. avium* reported earlier, considerable conservation in the synteny between genomes (gene order) within *M. avium* strains was expected. The order of GIs was used as markers for testing the conserved gene order and the overall genome structure between *M. paratuberculosis* and *M. avium* genomes.

It was unexpectedly discovered that, when the GIs associated with both genomes were aligned, three large genomic fragments in *M. paratuberculosis* were identified as inverted relative to the corresponding genomic fragments in *M. avium*. These fragments had the sizes of approximately 1969.4 Kb, 863.8 Kb, and 54.9 Kb (FIG. 7). The largest inverted region (INV-1) of approximately 1969.4 Kb is flanked by MAV-4 and MAV-19. INV-1 encompasses bases 1075033 through 3044433 of the *M. paratuberculosis* genomic sequence. The second inverted region (INV-2) of approximately 863.8 Kb is flanked by MAV-21 and MAV-24. Located near the origin of replication, INV-2 encompasses bases 3885218 through 4748979 of the *M. paratuberculosis* genomic sequence. The smallest inverted region (INV-3) of approximately 54.9 Kb is flanked by MAV-1 and MAV-2. INV-3 encompasses bases 320484 through 377132 of the *M. paratuberculosis* genomic sequence.

Because the sequences of the inverted regions and of the flanking MAVs are known, it is possible to use the junction

regions (sequences) to identify the presence of either *M. paratuberculosis* or *M. avium* in a sample. For example, using the right sets of primers, one skilled in the art would know to detect sequences that are specific to the junction regions that are characteristic for either *M. avium* or *M. paratuberculosis*.

Referring to FIG. 7, the location of genomic islands present in *M. avium* (dark grey boxes numbered 1-24, outer circle) or in *M. paratuberculosis* (light grey boxes numbered 1-18, inner circle) genomes are drawn to scale on the circular map of *M. avium* (outer circle) as well as the map of *M. paratuberculosis* (inner circle). The sequences of *M. paratuberculosis* K10 (query sequence) compared with the whole genome sequence *M. avium* 104 ORFs (target sequence) using BLAST algorithm with cut off values of $E > 0.001$ and alignment percentage $< 25\%$ of the whole gene were accepted as indications for gene deletion. The numerous short bars represent predicted ORFs in forward (outermost) or reverse (innermost) orientations. Large arrows indicate sites of genomic inversions.

Because the bioinformatics analysis used raw genome sequences, PCR and sequencing approach were used to substantiate the genomic inversions in 7 mycobacterial isolates (3 isolates of *M. avium* and 4 isolates of *M. paratuberculosis*). As predicted from the initial sequence analysis, primers flanking the junction sites of the inverted regions gave the correct DNA fragment sizes and orientations consistent with the sequence of *M. avium* and *M. paratuberculosis* genomes.

TABLE 11

Clinical		Genomic island					
Isolate	Subspecies	MAP-1	MAP-3	MAP-5	MAP-12	MAP-16	MAP-17
JTC33666	<i>M. paratub.</i>	+	+	+	+	+	+
JTC33770	<i>M. paratub.</i>	+	+	+	+	+	+
CW303	<i>M. paratub.</i>	+	+	+	+	+	+
1B	<i>M. paratub.</i>	+	+	+	+	+	+
3B	<i>M. paratub.</i>	+	+	+	+	+	+
4B	<i>M. paratub.</i>	+	+	+	+	+	+
5B	<i>M. paratub.</i>	+	+	+	+	+	+
DT3	<i>M. paratub.</i>	-	+	+	+	+	+
DT9	<i>M. paratub.</i>	-	+	+	+	+	+
DT12	<i>M. paratub.</i>	+	+	+	+	+	+
DT19	<i>M. paratub.</i>	+	+	+	+	+	+
JTC1281	<i>M. paratub.</i>	-	+	+	+	+	+
JTC1282	<i>M. paratub.</i>	-	+	+	+	+	+
JTC1283	<i>M. paratub.</i>	-	+	+	+	+	+
JTC1285	<i>M. paratub.</i>	-	-	+	+	+	-
JTC1286	<i>M. paratub.</i>	+	+	+	+	+	+
T93	<i>M. avium</i>	-	-	-	-	-	-
T99	<i>M. avium</i>	-	N/A	+	-	+	+
T100	<i>M. avium</i>	+	N/A	+	+	-	+
DT30	<i>M. avium</i>	-	-	-	-	-	-
DT44	<i>M. avium</i>	-	-	-	-	-	-
DT78	<i>M. avium</i>	-	-	+	-	-	+
DT84	<i>M. avium</i>	-	-	-	-	-	-
DT247	<i>M. avium</i>	-	-	+	-	-	-
JTC956	<i>M. avium</i>	N/A	-	N/A	-	+	+
JTC982	<i>M. avium</i>	-	-	+	-	-	-
JTC1161	<i>M. avium</i>	-	-	+	N/A	+	+
JTC1262	<i>M. avium</i>	-	-	-	-	-	-
JTC33793	<i>M. avium</i>	-	-	-	-	-	-

Symbols (+ or -) denote presence or absence of genomic regions; N/A denotes no amplification of DNA fragments.

41

It is to be understood that this invention is not limited to the particular devices, methodology, protocols, subjects, or reagents described, and as such may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention, which is limited only by the claims. Other suitable modifications and

42

adaptations of a variety of conditions and parameters normally encountered in clinical prevention and therapy, obvious to those skilled in the art, are within the scope of this invention. All publications, patents, and patent applications cited herein are incorporated by reference in their entirety for all purposes.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 38

<210> SEQ ID NO 1
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: AMT31 transposon-specific primer

<400> SEQUENCE: 1

tgcagcaacg ccaggtccac act 23

<210> SEQ ID NO 2
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: AMT38 degenerate primer
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (23)..(26)
 <223> OTHER INFORMATION: n is a, c, g, or t

<400> SEQUENCE: 2

gtaatacgac tcactatagg gcnnnncatg 30

<210> SEQ ID NO 3
 <211> LENGTH: 24
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: AMT32 nested primer

<400> SEQUENCE: 3

ctcttgcctc tccgcttctc ctcc 24

<210> SEQ ID NO 4
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: AMT39 T7 primer

<400> SEQUENCE: 4

taatacgact cactataggg 20

<210> SEQ ID NO 5
 <211> LENGTH: 19
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: AMT 152 primer

<400> SEQUENCE: 5

ttgctcttcc gcttcttctc 19

<210> SEQ ID NO 6
 <211> LENGTH: 19
 <212> TYPE: DNA

-continued

<213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: SP-6 primer

<400> SEQUENCE: 6

tatttaggtg acactatag 19

<210> SEQ ID NO 7
 <211> LENGTH: 1167
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 7

tcagcttacg gtgacaaccg gtgaaccgct ggatgttgcc tccggcccg cctcaactgtt 60
 catctgcgcg gcgatgcgca tcgctcttc gatcagcgtc tccacgatct gggattcggg 120
 caccgtcttg atcacctcgc cgcgaacgaa gatctgcccc ttgccatttc ccgacgcgac 180
 gcccaggctg gcctcgcggg cctcccccg ccggttgacc acgcagccca tcaccgcgac 240
 ccgcagcggc acgtccaggc cgtccaggcc ggcgctgacc tcgttgccca gctgtagac 300
 gtgcacctgc gcgcgccccg acgacgggca ggacacgata tcgagcgaa ggcgcgcag 360
 gttcagcgac tccaggatct ggatgcgcac cttgacctcc tcgaccggcg gcgcgcgacg 420
 cgacaccccg atggtgtcgc cgataccgcg cgacagcagc gcaccgaacg cgaccgcgga 480
 cttgatggcg ccctgaaacg cgggccccgc ctcggtgacg ccaggtgca gcgggtagtc 540
 gcaactgctg gcgagttgct cgtaggcggc gaccatcacc acggggctgt tgtgcttgac 600
 gctgatcttg atgtcgaaa agccgtgctc ctgaaacagc gaggcctccc acagcgcgga 660
 ctgcaccagc gcctcggggg tggccttgcc gtacttgcc atgaaccgct tgtccagcga 720
 gcccgcgttg acgcgatgc ggatcggaat gcccgctgcc gcagcggtt tggcgacctc 780
 gcccaccccg ccgtcgaatt ccttgatgtt gccgggggtc acccgaccg cggcgcgacc 840
 ggctgcgatg gccgcgaaga tgtacttggt ctggaagtgg atgtcggcga tcaccgggat 900
 ctggctgtgc cggcgatct cggccagcgc gtggcgctcc tccctggcgc ggcaggccac 960
 ccggacgatg tcgcagcccg ccgcggtcag ctggcgatc tgctgcagcg tcgagttcac 1020
 gtctggggtc ttggtgtgac acatcgactg caccgagatc ggatagtcgc tgccgacccc 1080
 gacgtcgcgc accatcagtt ggcgcgtgcg gcgcgggggc gcaagcgtgg gcgcgggggt 1140
 ctgcggcatg cccaggcctg tcgtcac 1167

<210> SEQ ID NO 8
 <211> LENGTH: 12084
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 8

gtgggtggag acttggggga gtcgatggcg agggatgata gggcgtttc gctgacgcga 60
 gggcagcttg atatctggtt atcccaggaa gccggtttcg ccggcaccga gtggcagctc 120
 ggtctgctgg tcaagatcga cggcaaggtc catcgcgacg cgctcgagca ggccatcacc 180
 caggccgtcg ccgaagccga acccggccgg gtctcgttct tcgagctcga cggccagggtg 240
 gtgcagaagc cgatcgacta ccgcacgctc gagctggctt ttcacgacct gaccgaccac 300
 gccgaccccg tggccgaggc gcgggagatg tcttcggcca ttcagcgac gccgatgccg 360
 ttgaacggcc aaatgttcaa attcgtgctt ttccaaacag ggcacgacga attctatttg 420
 tttggtgtct gccaccacat agctatcgac ggtctgggca tggctcttgt ttgccgacgg 480

-continued

gtggccacca	tttattcggc	aatggtggcc	ggtaagccga	ttccggacgc	ttacttcggc	540
acggtgcagg	acctgatcga	cctggagtcg	ggctacgagg	cctccccgga	ctatgccgag	600
gacaaggcgt	actggagcga	gcacctcccg	ccggagagcg	ggccggtcga	ccggtgccc	660
gacgccgaag	gggagcgcga	ccactactcg	ccgtccgcgt	cggtgcagct	ggaccgcgcc	720
gtcgccaacc	ggatcaagga	gctgtccaaa	aagcttgcca	tccgccgctt	ttcggtcacc	780
accgccgcgt	gcgcgctggt	ggtcgcgggc	tggtcgggta	gcggatcggg	ggcggcgctg	840
gacttccccg	tcagccgacg	ggcgcgtccg	gagtcacaaa	cgctgcccg	gatgctggcc	900
ggcgtggtgc	cgctggtgct	cagcaccgcg	cccagatcga	cggtggccga	cttctgcaag	960
cacgtcgaca	agcgcacccg	cgagctgctg	gcgcaccagc	gcttcccggc	gcacaccctc	1020
gaaggcgacg	ggttgcggca	ggcgcccaac	cggttcggga	tcaacttcac	cccgtcccgg	1080
ctgacgctgg	acctggccgg	ttccccggcg	acggcgtcgt	acaccaacca	cgccccggcg	1140
gggcacttcg	ggctgttctt	cctgggcggc	ggcgaccagc	tgttctctag	caccgcgggc	1200
ccgggcccaac	cgcttcgcag	cttcggcgtc	gcggacctgg	ccggtcggct	gcagcagatc	1260
ctggccgcga	tgaccgagga	cccgaccgcg	ccgctgtcct	cgatcgaact	gctgaccggc	1320
gacgagcccc	cgctgatcga	ccggtggagc	aaccgtcccg	cgctgaccga	gcccgcaccc	1380
gccccgggtg	cgatccccca	ggccttcggc	gaacacgtgc	agcgaccccc	cgacgcgggtg	1440
gcggtgacgt	tcggggcgac	ctcgtgacc	tacgcccagc	tcgacgaggc	gtccaaccgg	1500
ctggggccatc	tgtcgcgcga	ccacggcggtg	ggccccggcg	actgcgtggc	ggtgatgttc	1560
ccccgctgcg	ccgacgccat	cgtctcgatg	ctggcgggtg	tcaagaccgg	ggcggcctac	1620
gtgccgatcg	acccggcgca	cgcgtcgtcg	cggatggact	tcgtgctcgc	cgacgcgcgc	1680
cccagcgcgg	tgatcaccac	ctccgacctg	cgtcgcgggc	tggacgatca	cgacctcctc	1740
gtcgtcgacg	tgacgacccc	ggcgttcgaa	gcccagcccc	gcaccgcgct	gccgtggccg	1800
gcgccggagg	acaccgccta	catcatctac	acctcgggaa	ccaccgggac	ccccaaaggt	1860
gttgccattc	ctcatctcaa	cgtcacctgg	ctgatcgagt	cgctggacgc	cggcctgccg	1920
cccgaaaacg	tgtggagcga	gtgccactcg	tcggcgcttcg	acttctcggg	gtgggagatc	1980
ttcggcgcgc	tgtcgcgcgg	ccggcgactg	ctggtggtgc	ccgagtcggg	ggcgtcgtcg	2040
ccggaggact	tccacgcctc	gctggtcgcg	gagcaggtca	gcgtgctcac	ccagacgcgc	2100
tcggcgggtg	cgatgctctc	acccgagggc	ctggagtcca	ccgcgctagt	ggtggccggc	2160
gaggcctgcc	cgaccgacgt	ggtcgaccgg	tgggcggcgc	ccggtcgggt	gatgctggac	2220
gcctacggcc	cgaccgagac	cacggtgtgc	gcgtccatca	gcacaccgct	gacggccggc	2280
gacccgggtg	tgccgatcgg	ctcgcgcgac	gcccggggcg	cgatgttcgt	gctcgacaag	2340
tggctgcagc	cgggtccccc	cggcggtggtg	ggcgagctgt	acctggccgg	ccgcggcggtg	2400
gggcacggct	acgtgcgcgc	gcccggcctg	accgcctcgc	ggttcgtgcc	caaccggttc	2460
ggcgcccccg	gctcgcggat	gtaccgcacc	ggcgacctgg	tgtgctgggg	ccccgacggg	2520
cagctgcagt	acctggggcg	cgcgcgagag	caggccaaga	tccgcggctt	ccgcacgcag	2580
ctcggcgaaa	cccagtcggg	gctggccggg	ttggacgggg	tggagcaggc	ggcgggtggc	2640
gcccgcgagg	accggccccg	cgacaagcgc	ctggtcgggt	acatcaccgg	caccgcgcac	2700
ccggccgagc	tgcgcgcgca	gctggccgac	cggctgcgcg	cctacatggt	cccgaaccgc	2760
gtgatggtgc	tggacgcgct	gccgctgacc	ggcaacggca	agctggacaa	gcgcgcgctg	2820

-continued

ccctcgccgg	aatacgccgc	cggcgaatac	cgggcgcccc	gcgacgcgat	cgaggagatc	2880
ctggccgaca	tctacgcccc	ggtgctgggc	gtggagcggg	tcggggtgga	cgactcgttc	2940
ttcgacctgg	gcggcgacag	catcctgtcc	atgcaggtgg	tggcccgccg	ccgcgcggcg	3000
ggggtgatct	gtcggcccg	cgacgtgttc	gtcgagcaga	cggtagcccg	gctggccgg	3060
gtgtcccaag	tggctgtcga	cggcgagctg	ggcgccggcg	acgaggggat	cgggccgggtg	3120
cagccccccc	cgatcatcgg	ctggctgcag	gacatcgacg	gcccgatcga	cgagttcaac	3180
cagaccatgg	tgtcggccgc	gcccgcgggg	gtcgggtgtc	acgacgtcgc	ggtggtgctg	3240
caggcactgc	tggaccggca	cgcgatgctg	cggctgtgcc	tcgacgacga	cggcgcgggc	3300
ggctgggacc	tgacgtgcc	gcccccggt	tcggttgacg	ccgcgcgcct	cctgcgcacg	3360
gtcgacgtgc	tctccgaggc	cgcgctggcg	cgggcgcggg	cccggctgaa	ccccggcgcc	3420
ggcctgatgc	tgtcccggtt	atgggcaagc	gccaccaacg	aattggccct	ggtcgttcac	3480
cacctggcgg	ttgacggggg	gtcgtggcgg	acgttgatcg	aggacatcaa	catcgccctg	3540
gcgcagcatc	acagcgggtc	ggagatcgcg	ttgccggtgc	cgggcacgtc	gtttgcgcgg	3600
tggctcgtcga	ttctggccga	gtacgccaa	agcccgccag	tggtagctgc	ggcggcgccg	3660
tggcagcagg	tggtagccac	gcccggcggt	ctgccggcgg	tggggcccca	tgacacctat	3720
gcctcggagg	ggcagttgtc	ggcgtcgtc	gatgtgcaga	ccacccggtt	gttgttgggg	3780
gaggtgcccg	cggcgtttca	cgtcgggggt	caagacattt	tgctgattgc	gttcgggttg	3840
gcctgcacgg	agtctcgtgg	tggtagccgc	ccgatcggtc	tcgacgtgga	gggtcacggg	3900
cggcacgagg	agatcgccct	gggggtggat	ctgtctcgca	cggtaggctg	gttcaccacg	3960
aaatacctg	tggcactgac	gatcagtcag	cgtctggatt	gggcgcgggt	ggtggcgggg	4020
gaggccgcgc	tgggcgcggg	gatcaaggat	gccaaaggag	agttgcgggc	gctgcccga	4080
ggcctgagct	acgggttgct	gcgctacctg	aaccccgaga	tcgaggtgca	ggggccggat	4140
cgggtgatcg	gattcaacta	cctgggccc	ctcggcgccg	cggccgcga	cctgtccgac	4200
gagcactggc	gcctcagccc	cgacagtcgg	tcggtagcgg	ccgcggccgc	ggcgatccca	4260
ctgccgttgg	gacataccgt	cgaactcaac	gccggcacca	tggacaccga	cgcgggccc	4320
cagttgcacg	ccaactggac	ctgggcgcgc	tccgtgctca	ccgacgagca	gctaaaccgg	4380
ttgagccggg	tgtggttcga	ggcgtgacc	ggcatctcgg	cgcacgtgca	ggcggcgccg	4440
ggcgggctga	cgcggtccga	catcgccccc	accctcctcg	accaaggccg	gatcgagcag	4500
ctggaacggc	actacgacgt	cgcgacatc	ctgccgctga	ccccgctgca	gcaggggctg	4560
ctgttccacg	cgaccggaag	ccatgcgcag	ggcgacgtct	acgcggtgca	gctgagcgtc	4620
acgctgcgcg	gcgccctcga	cccgcaccgg	ctacaccgcg	ccctgcacac	cgtcgtcacc	4680
cgccacccga	acctggccgc	cgccttctgc	cccagctcgg	gcgagccggg	gcagatcatc	4740
ccggccgaac	ccgaaatggc	ttggcgctac	ctcgaactcg	acggcggcga	catcgacgaa	4800
cagctcgagc	agctgtccgc	ggacgaacgc	gccgcgggtc	gagagctcgg	cgaccgccc	4860
ccgtttggcg	cgcgctgat	cgcacccgcg	gacacggaac	accggttcgt	gctcaccgtc	4920
caccacctgg	tgatggacgg	ctggctcgtg	ccggtgctgc	tgcaggaaat	cttcgcctgc	4980
tactacggtg	cccggctgcc	ggcgccggcg	ccgtaccgcg	gcttcgtcac	ctggctggcg	5040
gcccgcgacg	tgccggccgc	cgcgcgcga	tggcgccggg	tgctcgacgg	tttcgacacc	5100
cccaccctgg	tggccccgcg	gggtgcccga	gcgccggggc	ggcgccgggt	cgcctcgttc	5160
cgggtggccg	ccgaaaccac	cagcgcggtc	agcgaactcg	cacgcgcgcg	ccgcaccacc	5220

-continued

gtcaacacccg	tgttcgaggc	cgcttgggcg	cagctgctga	tgatgctgac	cgcccgacac	5280
gacgtcgctg	tgggcaccgc	cgtctctggc	cgcccgcccg	agctgcccgg	cgccgagtcg	5340
atggtcgggc	tgttgatcaa	caccgtcccg	gtgcgcgccc	acgccaccgc	ggcgaccacc	5400
atcgcgagcc	tcgtagacca	gctgcaacgc	gccacaaacc	acaccgtgga	gcatcagcac	5460
ctggcgctca	acgaatcca	cgcctacc	ggacaggacc	aactcttcga	caccctgctg	5520
gtctacgaga	actatccgat	cgacaccgcc	gccctgtcgg	cgcccgacga	cctcaccgcc	5580
accgaattca	gctgccacga	ctacaaccac	taccgctgt	cgctgcaagt	ggcgcccgcc	5640
gacgaactgg	gccttcgcct	cgaattcgac	accgacgtgt	tcgacccggc	ggccatcgac	5700
accctggccg	accggttgcg	gaagctgctg	gccgccatgc	cgcccgaccc	ggaccgcccg	5760
ttgcgatcac	tggacctgct	cgacgccacc	gagcacaccc	ggctgcaacg	gtggggcaac	5820
cgcccgccgc	tgagccggcc	ggcaaccggg	ccgtcgctgc	cggagttgtt	cgccgcacag	5880
gtcgccaacg	ctccgcacgc	cgtcgcgctg	cgctacgccg	gccggtcgat	gacctaccgc	5940
gaactcgacg	aggcgctcgac	ccggctggcc	cacctgctgg	ccggccacgg	cgccaccccg	6000
ggttgctttg	tggcactgct	gttttcccg	tcggccgagg	cgatcgctgc	gatgctggcg	6060
gtgctgaaaa	cgccgcggcg	ctacctgccg	atcgaccccg	cgctgccggc	gacccgcatc	6120
gagttcatgc	tcggcgacgc	cgcacccgtc	gtcgcggtca	gcaccgccga	cctgcgcgcc	6180
cggtggaggg	ccttcggcct	gccggctcgt	gacgtcgccg	ccaccggcgc	ccagcccgcc	6240
ggcccggttc	cgccgcccgc	gcccgacaac	atcgccctatc	tgctctacac	gtccgggacc	6300
accggcgctc	ccaagggcgt	tgccgtcacc	caccgcaacg	tcgcccagct	gctcgagctc	6360
ctgcacgcgt	cgctgcccgg	caccgggggtg	tggtcgcagt	gccactccta	cggttcgac	6420
gtctcggtcc	aggagatctg	ggcgccctcg	gccggcgggc	gccggctggt	ggtggtgccc	6480
gagtcgggtg	ccagctcacc	cgacgagctg	cacgcgctgc	tgatcgccga	gaacgtcacc	6540
gtgctcagcc	agacaccgtc	ggcgctggcg	gcgctgtcac	cacgaaacct	gcacgcggcg	6600
ttggtgatcg	gcggcgagcc	ctgcccggcc	gcgctcgccg	accggtgggc	gcccgggccg	6660
gtgatgatca	acgcctacgg	ccccacggaa	accaccgtcg	acgcggtgct	cagcacaccg	6720
ctggcccgcc	gcgcgggagc	acccccactc	ggctcccccg	tagcgggtgc	gacgctgttc	6780
gtgctggagc	cgtggctcgc	gcagggtcct	gccggtgtga	ccggcgagct	ctatatcgcc	6840
ggcgccgggg	tggccgcggg	ctatctgggc	cgcccccgtc	tgacggcgcc	gcggttcgtg	6900
gcctgcccgt	tcggcgacgc	cgggtgcggg	atgtaccgca	ccggcgatct	ggtgcgctgg	6960
gacgcgagc	gccgactgca	ctacgtcgcc	cgggccgcatc	agcagggtcaa	gattcgcgcc	7020
caccgcatcg	agctggggcg	aatccattct	gcgctggccg	aattggacgg	cgtcggggaa	7080
gtagcggtga	tcgcccgcga	ggaccgcccc	ggcgagaaac	ggatcgctcg	ctacctcacc	7140
ggcaccgccg	acccggcgcc	gatccgcgcc	cggctggccg	agcggttgcc	ggcctacatg	7200
gttcccgcgc	cgggtgctgg	gatcgaggcg	ctgccgttga	ctcccaacgg	gaaactggac	7260
gcccggggccc	tgccggcgcc	ggaatacgcg	ggcggggcat	accggggccc	gtccactccc	7320
accgaggaga	tcacgcggcg	catctacacc	cagggtgctcg	gcctgcacag	ggttggtgtc	7380
gacgactcgt	tcttcgacct	ggcggggtgat	tcgctgtcgg	cgatgcgggt	gatcgccgcc	7440
gtcaacgccg	gcctcgacgc	ccggctgtcg	gtgcgagtg	tggtcgaaag	gccaccattt	7500
gcgcaactgg	cggcgcgccct	cggcgaaagg	gggcaccggg	tcgccgcggg	ggtggccgcc	7560

-continued

gagcggcccg	cggtggtgcc	gttgctgttc	gcgcagtcgc	ggctgtggtt	catcggccag	7620
ctgcacgggc	cgtccccggt	gtacaacatg	gtggccgcgc	tgcggctgca	cgggccgggtg	7680
gacatcggcg	cgctgggcgc	cgcaactgcat	gatgtcgtga	cccggcagca	gagcctgcgc	7740
acggtgttcg	ccgcgaccga	cgggacgccc	gcccagtggt	tgtgcccgc	cgaccgtgcc	7800
gacatcggct	ggcaggtcat	cgacgccagt	ggctggtcgc	cgggcccgagt	ggatgacgcc	7860
atccgcgaca	ccgcccggca	tacctttgac	ctggctgctg	aaattccgtt	gcgtgcagtg	7920
cttttgcggt	gtggcgcgga	ggagcatttg	ttggtggcgg	tggtgcatca	tattgccgcg	7980
gacgggtggt	cgttgacgcc	gttggtgcgt	gacttggcgc	gggcgtatgc	gagtcggtcg	8040
gcggggcggg	tcccggattg	gggtgccgtt	ccgggtcagt	atgtcgatta	cacgttgttg	8100
cagcgcgccc	agttcggtag	cctcgacgac	ccgcacagcc	tgatcgccgg	tcagctgcgc	8160
tactgggagc	acaccctggc	gggcatgccc	gagcggtttg	aattgcccac	cgatcggccg	8220
tatccggtgg	tggccgattt	ccgcggcgcc	agtgtcgcgg	tggagtggcc	ggcgcagttg	8280
cagcagcaaa	tatcgcggtt	ggcgcggggc	cataatgcca	ccagtttcat	ggtggtgcag	8340
gccgcgttgg	cggtgctgct	ggccaagggt	agcgcgagtt	cggtgtgggc	ggtgggcttt	8400
ccgatcgccg	ggcgccgcga	ccggcgctg	gatgacgtgg	tgggtttttt	cgtaacacc	8460
ctggtgctgc	gggtggacgt	ctccggtgat	cccacggtgg	gcgagctgct	ggcgcgggtg	8520
cggcaacgca	gcctggctgc	ctatgagcat	caggatgtgc	cgttcgaggt	gctggtggag	8580
cggctcaacc	cgccccgcag	tctggcgcac	catccgctgg	tgcaggtgat	gttggcctgg	8640
cagaacatcg	agcccaccga	gctgagcctg	gggcaggtgc	gggtgactcc	gctgccggtg	8700
gatacccgca	ccgcccggat	ggatctggct	tggtcgctgg	ccgaacgctg	ggcaccgat	8760
ggctcaccgg	ccggtatcgg	gggagcggtc	gaattccgca	ccgacgtggt	cgataccgcc	8820
acggtggagg	cgttgaccca	gcggttgccg	cggtgctgg	cggccatgac	cgccgacccc	8880
ggccgcgggt	tgtcctcgat	cgacctgctc	gaccccgacg	agcacgcccg	cctcgacgcc	8940
ctcggcaacc	gcgcagcact	gacccgacca	caaaacccgc	ccacctccat	cccgcgatg	9000
ttcgcgcgcc	agatggcgcg	caccccgcac	gccgtggcgc	tgaccgcca	cggtcgctcg	9060
gtcacctatc	gccggctcga	ggaacacgca	aaccaattag	cgcaccaact	tattcgttac	9120
ggcgcagggc	cgggcgattg	cgtggcgctg	ctgctggagc	gttcgcgcga	ggcgtcgcg	9180
gccatcctgg	gggtgctcaa	ggccggggcc	gcctacctgc	ccatcgaccc	cagcctgccc	9240
agtgcgcgga	tcgagttcat	gctcacggac	gccgcacccg	cggccgtgct	caccagcacc	9300
gaattccatt	gccgtctaca	ggattaccac	cagaccgtca	tcgacgtcga	cgacccgtcg	9360
atccgggaac	aaccgcgtac	cgcaccaccg	gcgcccgcgc	cggacaatat	cgctacctc	9420
atctacacct	cgggcaccac	cggcgtcccc	aaaggcgtcg	cggtcaccca	ccgcaacgcc	9480
acccagctgt	tcgcgtcgct	gggagccgcc	ggcctgcccc	ccgcacccgg	aaaggtgtgg	9540
ggccagtgcc	attcgctggc	cttcgacttc	tcagtgtggg	agatcttcgg	cgcgctcctg	9600
aacggcgggc	gcgtgctggt	ggtgcccgc	gacgtggtgc	gtccccgga	agacctgtgc	9660
gccttgctga	tcgaggaaacg	ggtcgacgtg	ctcagccaaa	cgccgtcggc	attcgatgcg	9720
ctgcagcgcg	ccgactccgc	ccggcggtc	aacccgcaga	cgggtgatctt	cgggggcgaa	9780
gcgctgatcc	cgcaccggct	ggcgggctgg	ctggaagggc	atcccgcaacg	cccgcggctg	9840
atcaacatgt	acggcatcac	cgagacgacg	gtgcacgcct	ccttcgggga	gatcgctgac	9900
ggcgacatcg	acggcaacgt	cagcccgatc	ggaatgccct	tggcgcaact	gggattcttc	9960

-continued

```

gtgctggatg gatggctgcg gcctgtgcct gccggtgtga ccggcgagct gtacatcgcc 10020
ggcgccgggg tggccgcccgt ctatctgggc cggcccggtc tgacggcgtc gcggttcgtg 10080
gcctgcccgt tcggcgggcg cggcgagcgg atgtaccgca ccggggacct ggcccgggtg 10140
ggcgccgacg ggcagctgca atacctgggc cggcgcgacg aacaggtaaa gattcgcggc 10200
taccgcatcg aactcggcga aatccagtcc gccctggcgg aattggacag cgtcgagcag 10260
gcggcggtga tcgcccgcga ggaccgtccc ggcgacgagc ggctggctgc atacgtcacc 10320
gggacggcgg acccggcgca gctccgcacc gcgctgaccg aacggctgcc cgcctacctg 10380
gtcccgcggc cgggtgctggt gctggacgcg ctgccgttga caccagcggg caaactcgac 10440
accggcgccc tgcccgcggc cgactaccag ggcccggagg actacctggc cccggccggc 10500
gcggtggagg agatcctggc ctggctctac gccacgggtg tggggctgcc gcggcggttc 10560
ggggtgcagg aatccttctt cgacctgggc ggcgactcgc tgtcgcccat gcggtggttc 10620
gcggccatct acaacgcgct ggacatccac ctgccgggtg gggccgtctt cgaggcgccc 10680
tcggtgcgca gcctgagcca gcggctgaac gccgatcccg ctgtggcgca aggccttcgg 10740
gccgacttcg catcggtgca cggccgcgac gccaccgagg tgtacgccag cgacctgacc 10800
ctggacaagt tcacgcagc cgcgacgctg tcgcgcgcac ccgcgctgcc cggccccggc 10860
gccgaggtgc gcaccgtgct gctgaccggc gccaccggct ttctggggcg ctacctggtg 10920
ctgcaatggc tggaacgcct ggaactggcc gacgggaaac tcactgtctt ggtgcggggc 10980
gcctcggacg acgacgcgcg gcgcgcctc gagcgcaact tcgacagcgg cgatcccggc 11040
ctgctcgggt acttccacga actggccgcc gaccacctcg aagtcacgc cggcgacaag 11100
ggccgcgcca acctcggcct ggacgatcgg acctggcagc ggctggccga caccgtcgac 11160
ctgatcgtcg acgcgggcgg cgtggtcaac ggctgctgct cctaccagga actgttcggc 11220
cccaacgtcg ccggcacgcg cgagctgacg ccgctggcgc tgtccaccag actcaagccg 11280
tacagctacg tgtcgaccgc caacgtcggc gaccagatcg agccgtcggc gttcaccgag 11340
gacgcccaca tcgggtgctc cgggcccgat cgcaccatcg acggcggtta cggcaacggc 11400
tacggcaaca gcaagtgggc cggcgagggt ttgctgcgcg aggcgcacga cctgtgcggg 11460
ctgccggtct cgggtgttcg ttgcgacatg atcctggccg acaccagcta cgcggggcag 11520
ctcaacctgt cggacatggt caccggctg ctgttcagcg tgggtggcag cggcgtggcg 11580
ccgcgctcgt tctaccggct cgacgcgcac ggcaaccggc agcgcgcgca cttcgacgcg 11640
ctgccggtcg agttcgtcgc cgaggcgatc gccacgctgg gcgcccaggg ggcccgggac 11700
gccggcatcg ggttcgcgac ctaccacgtg atgaaccgc acgacgacgg catcgggctc 11760
gacgagtacg tcgactggct gatcgaggcg ggctacctga tcgagcgggt cgacgaattc 11820
gaccagtggc tgcaccggat ggagaccgcg ctgcacgccc tgccggaacg gcagcgccac 11880
cagtcggtgc tgcagctgct ggcgttgctc aaagcccggc acgtgccgcc ggccgacccg 11940
gcccgcggct gcctggggcc caccgagcgg ttccgcgctg cgggtgcaaga agccaaaatc 12000
ggcgccgaca acgacatccc gcacatcacc gcccgggtca tcgtcaaata cgtgaccgac 12060
ctgcagttgc tgggcttgcct ctga 12084

```

<210> SEQ ID NO 9

<211> LENGTH: 876

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

-continued

<400> SEQUENCE: 9

ctagctggcg acccgatagc ggtggctcgag ttgcagattt agctgcagca cggtgacgca	60
cggttcacccg aaaaagccca gcgcgcgcgc gctgctgttc tgcgccacca gctcccggat	120
ctgctccggg cggacgtggc gcaccttggc caccgcgcgc acctggatgt cggcgtaggc	180
cggcgagatg ttcgggtcca ggcgctgcc gccggcggtg accgcgtcgg cgggcacggc	240
cggggcgcgc ggtgcgcgc gcgcgatggg cagcatctgg ccgagcgaat agtcctcccc	300
ggtcttggcg cattcgacgc gcacccctc gtagaggctc acgaacggtg tcggggtcga	360
ttcgcacggt tcggtgacgc tgaccaccgc ggcgggtgg gtgacgttgc cgcgctcgtc	420
gcgcggcccg atcacgcaca gcacgcgcgc cacacgcgc ccggtgcaga acggccgcga	480
cccgtcgacg cctccagtt tggcgacggc ggcgctgcgc gagcacaccg tggtcagcag	540
gcggggtttg ccgggtgcgt cgcagatgct ctccggcccc aggttgctgc cgcgctgga	600
ggtcgggtcg tagccggtgc cggccgcga gggcgactc tgaagtagt gcgcgacggg	660
gttgccgtcc ttgtcggtga acagctggcc gatcagtcgt ctgccaccgc gcttgccgtt	720
ggcggtcagg atcgaccctc cgcgctggc gcgtaatccg gggaactcgc ccaccaccca	780
gacgagcagc ggataggcca ggcgggtgat caggttcagc accagcagcg cccgcaacgc	840
cgccagtcgc aggcgaatga agttcgacag tgtcat	876

<210> SEQ ID NO 10

<211> LENGTH: 1518

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 10

gtggtcacat tcgggacggt gcataactgg gatccgggaa ccggttcggt catttcctgg	60
cacgcgactc cggcggcgcg ggaanaagcc cggcaggcgc cgattagcga cgtgccggcc	120
agttatcagc agttgcatca tcttcgcgcg ttcagcgaac acgcgcgcgc gggcctggac	180
atggcgcggc tgaacatcgg ggtctgggat atctccggcg tgtgcgacgt ggcggccatg	240
accgaagcga tcaatgccc tctgcgcgcg cagcacacct accacagttg gttcgagcat	300
cggaccgacg gccgtatcgt ccggcacacc ttgcgcgac ccgcgcacat cgagttcgtc	360
gcgctgcagc gcggcgagat gacgcgcacc gagctgcgcg cccacatcct ggccaccccg	420
aatccgctgc gctgggactg cttcaccttc ggtctcgtcc agcaccgcga tcaactcacc	480
ttctatatga gtgcggacca tctggtcac gacgggatgt cggtcggcgt gatcttcctc	540
gaaatccatc tgacgtatgc cgttttgggt agcggcgggc gcccgctgcc gctgcccgag	600
ccggccagct accacgacta ctgccgcgcg cagcaccagc acaccgaggc gctgaccctg	660
cagtcacccc aggtgcgcgc atggattcga ttgcgcgagg acaacggcgc aaccctcccg	720
agtttccccc tgccacttgg cgatccgtcg gtgcgctgcg gaagcggtgt ggtggtcgca	780
ccgctcatgg acgagagtca gaccgagcgc ttcgacgcca cctgcacgaa ggcgggggct	840
cgttcacgcg gtggcggtgt cgctgcgcgc gccttcgcgc agtacagct gaccggggcg	900
gagacgtatt gcgcgatcac gccgtatgac caccgcagca cgcgggcga attcgtgact	960
ccgggctggt tcgccagctt catcccggtc accgtcccg tcgcgggggc ctggttcggt	1020
gacgcggtga tcgcgcgca ggcgtccttt gattccgcta tcggtctggc cgacgtcccc	1080
ttcgaccgcg ttttgagct gtcgtcttcc ggcgggcgaa tcagcaaac gacgggcgac	1140
gtccacatgc tctcgttcgc cgcgcgcgcg gggatcccg tcagcgggca atgggacggc	1200

-continued

ctgaacgcgcg ggatttacgg cgaacggccg tcgtccgato aggtgcttat gtgggtcaac	1260
cgattcgaca ccgagaccac cttgacggtg gcgttccac agaateccgt cgcgcgcgat	1320
tcggtcgagc gctacatccg ggcggtcagg gcgatgtgtc tgcgggtggt ggagcacggc	1380
gcgcgcgcgcg tgccgaaccg ccggcgcgtc gttgcgcgcg tcaacgcgtc ggccgcgcga	1440
tcgaccgcca acgcgcgcga ccgcaccgac cgtcgccctgc aaggcgcggg cttcggcgc	1500
aaccgcggtcc ttcggtga	1518

<210> SEQ ID NO 11
 <211> LENGTH: 801
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 11

atggacctgg acgcgttggt ggcgcgggcg tcggcgatcc tcgacgacgc ctgaaagccc	60
ttcctcgccg gtcaccgcgc cgactccgcg gtccgcaaga agggcaacga ctttgccacg	120
gacgtcgacc tcgccatcga gcggcagggtg gtgcgcgcgc tggtgagggc caccgggatc	180
ggcgtgcacg gcgaggagtt cggcggttcg gccgtcgact cggaatgggt gtgggtgctc	240
gaccgcggtg acggcacggt caactacgcg gccgggtccc cgatggccgg gatcctgctg	300
gcccctgctgc accacggcga cccggtggcc gggctgacct ggttgccgtt cctcgaccag	360
cgctacaccg cggtgaccgg cggcccctg cgcaagaacg aaatcccgcg gccgccactg	420
acttccatcg acctggccga cgccttggtc ggggcgggat cgttcagcgc ggacgcccgc	480
ggccggttcc cggggcgcta ccggatggcg gtgctggaac acctcagccg ggtctcgtcc	540
cggttgcgca tgcacggctc caccggcctg gatctggcct atgtcgccga cgggatattg	600
ggcgcggcgcg tcagtttcgg cgggcacgtg tgggaccacg ccgcgggggt ggcgctggtg	660
cgggcgcgcg gcggcggtgt gaccgacctg gccgggcggc cgtggacccc ggcctcggtat	720
tcggcgctgg ccgccggacc cggcgcgcac gccgagatcc tggacatcct gcgcaatata	780
ggccgaccgg aggactactg a	801

<210> SEQ ID NO 12
 <211> LENGTH: 861
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 12

atgacgggct tggaaccgta ttacgaggaa tcgcagtcca tctacgacgt ctccgacgaa	60
ttctttgctg tatttctcga tccgacctg gcctacacgt gcgccttctt cgaacgcgac	120
gacatgacgc tcgaagaagc gcaaacgcgc aagttcgacc tggcgctggg caagctgaac	180
ctcgagcccc ggatgacgct gctggacatc gggtgccgct ggggcggcgc actgaagcgg	240
gccctggaga agtatgactt caacgtcatc ggaatcacgt tgagccgcaa ccaattcgaa	300
tacagcaaga agctgctggc cggatttcgc acggaccgca acatcgaggt gcgctgcag	360
ggctgggaag agttcgacga caaggctgac cggatcgta ccatcggcgc cttcgaggcg	420
ttcaaatagg agcgctattc cgcatttttc gaccgcgcct acgacatcct gcccgacgac	480
ggccggatgc ttttgacac catcttgacc tacaccaga agcagcagca cgagcgcggc	540
gtttcgatca cgatgagcga tttacggttc atgcggttca tcggccagga aattttcccg	600
ggcggccagt tgccggcgca ggaggacatt ttcaaatcgc gtgaggcggc gggtctttcg	660

-continued

gtggagcgagg tgcagctgct ggcgcgagcac tacgcgcgga ccctcaacat ctgggcggcg	720
aacctggagg ccaacaagga caaggccatc gccatccagt cgcaagcggt ctacgacgc	780
tacatgcact acctgacgg gtgcgagaac ttcttcgca agggcatcag caatgtggg	840
caattcacgc tggtaagta g	861

<210> SEQ ID NO 13
 <211> LENGTH: 750
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 13

tcacaggtgc cgccgcgcgc tgatctccat gacgggtgcg gtcagtacg acgacatgtc	60
ggaggccaaa aacagcgcca cgctggcgac ctgctgggc tcgccggccc ggcccatcga	120
cacctcgccc accttgaggt cccaaatgcg ttgcggcatg gcctctgtca tcgccgagcg	180
gatcaaacgc gggcgcatcg cgttcacccg cacacccagg taggccagt ccttggcggc	240
cgcttggtc atgcccacga tgcggccctt ggccgcgag tagttggtct ggccgacct	300
gcccagcttg cccgacacg acgacatgtt gatgatggcg ccgcgcttgt tttccgcat	360
gategcgccc gccaatcggt tgccgttcca ggtgcccttc aagtgcacg cgatgacctg	420
atcgaaatgc tctcggtca tcttgcgat ggtggcgctc cgggtgatcc cggcgttgtt	480
gacctgatg tccaggccgc cgaaccgctc gacggcggtc tggatcagcg tttcgacctc	540
ggacgacttg gtgacgtgc agcgacggc cagcgccacc tggtaacgc ccagctgttt	600
ggccgcgggt tgcgctgcct ccagattgac gtgcgcgagg acgacccgcg ccccttcggc	660
gacgaacctg tcggcgatcg cgaaccccaa accttgtgcg ccacctgtga tgaccgcggt	720
ctgacctg agcaaggaca cctgtaccac	750

<210> SEQ ID NO 14
 <211> LENGTH: 2289
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 14

atggcgatca tcgacaaaga gacgcaagtc cggccatcgt tcgacgacga ggtagccgc	60
acgcagcgct acttcgacga ccccgcttc gcccgcatca cgcgcctcta cacgctcgg	120
cagggtggcg agcagcgcg caccatccgc accgactaca ccgtggcacg cgacgcggcg	180
gcccgtttct acgaacggct ggcgcgagctg ttgcgccaga agaagagcat caccacctc	240
ggcccgtact cgcccgcca ggcggtcacc atgaagcggg tgggcatcga gggcatctac	300
ctcgcggggt gggccacctc ggccaagggc tccaccaccg aggacccggg gcccgacctc	360
gccagctacc cgctgagcca ggtgcccgac gacgcgcgg tggttggtgc cgcgctgctc	420
accgcccacc gcaaccagca gtacctcgcg ctgcagatga gcgaacgca gcgcgcggcc	480
accaaggagt acgactaccg gccgttcac atcgccgacg ccgacaccgg ccacggggga	540
gaccgcacg tgccgaacct gatccgcgt ttgctcgagg tcgggggtgcc cggctaccac	600
atcgaggacc agcggcccg caccaagaag tgtggccacc agggcgga ggtcttggtg	660
ccctccgacg agcagatcaa gcggctcaac gccgcgcgt tccagctcga catcatcgcg	720
gtgccgggga tcatcgctgc ccgcaccgac gccgaggccg ccaacctgct ggacagccgc	780
gccgacgagc gcgaccagcc gttcctgctg ggtgccacca acctcgacat cccgtcgta	840
aaggcgtgct tctggcgat ggtgcgcgc ttctacgagc tgggtgtcaa ggacctcaac	900

-continued

```

ggccacctgc tctacgcgct gccggaggcg gactacgccg aggccaccgc ctggctcgag    960
cgtcaggggca tccagggcgct catctccgac gccgtcaacg cctggcgcgga gaacggggcaa 1020
cagtcaccatcg acgacctgtt cgaccagggtg gactcgcggt tcgtggcggc ctgggaggagc 1080
gacgcggggc tgatgacctc cggcgaggcc gtgcgcgagg tgcctggagt gcgcgcgagc 1140
gagggcgagc cggctgacat gagcgccgac gactggcggg ccttcgcggc gcgcgcctcg 1200
ctctactccg ccaaggccaa ggcaaggag ctgggcttcg acccgggctg ggactgcgag 1260
ctggccaaga ccccgagggt ctactaccag atccggggcg gcaccccgta cgccatcgcc 1320
aagtcgctgg ccgcgcgcc gtgcgcgac atcctgtgga tggagaccaa gaccgcgcgac 1380
ctggccgacg ccaagcagtt cgccgacgct atccacgccg agttcccgca ccagatgctg 1440
gcctacaacc tgtgcgcgtc gttcaactgg gacaccaccg gcacgacgga cgagcagatg 1500
aagcagttcc ccgaggaact cggcaagatg ggcttcgtct tcaacttcat cacctacggc 1560
ggacaccaga tcgacggcgt gccgcgcgag gacttcgcca cctcgctgca acaggacggc 1620
atgctggcgc tggcccgctt gcagcgcaag atgcgtctgg tcgaatcccc ttaccgcaca 1680
ccgcaaaccg tcgtcggttg gccccgcagc gatgccgcac tggccgcctc gtcgggcgcg 1740
accgcgacca ccaaggcgat gccgcagggc tcgaccagc atcagcatct ggtgcagacc 1800
gaggtgcccc agaagctgct cgaggagtgg ctggcgatgt ggagcgagaa ctaccacctc 1860
ggcgagaagc tccgcgtgca gttgcggccc cgccggcggt gttcgacgt gctggaactc 1920
ggcatctacg cgcacggcga cgagcaactg gccaatgtcg tcgtcgacct gatcaaggac 1980
cggcacggcc gcagcatcct tcaggtgcgc gaccagaaca ccttcgcgga aaagctccgg 2040
cagaagcggc tgatgacgtt gatccacctg tggctggtgc accgcttcaa ggccgacgcg 2100
gtgatctacg tgacgccgac cgaggacaac ctgtaccaga cctcgaagat gaagtgcac 2160
ggcatcttca cgcaggtgta ccaggaggtc ggcgagatca tcgtcgcgga ggtgaaccgg 2220
ccccgcacgc ccgaactgct gcagcccgc cgggtggcgc tgcgcaagct gatcaccaaa 2280
gaggggttag                                     2289

```

```

<210> SEQ ID NO 15
<211> LENGTH: 234
<212> TYPE: DNA
<213> ORGANISM: Mycobacterium avium

```

```

<400> SEQUENCE: 15

```

```

ttacttgcca gcaccgcggc gctgcgcgc cagcctgtcc cgtaaagtct tcgggcggat    60
atcggggccag ttctgttcga tgtagtccag gcacgcagcg cgatccgctt cgccgtaaac 120
catctgccag ccggccggaa tgtcggcgaa cgccggccac aggtctgtgt gttcttcgtc 180
gttgaccagg acgaaaaagc tgccattgtc gtcgtcgaac ggattggtac tcac      234

```

```

<210> SEQ ID NO 16
<211> LENGTH: 972
<212> TYPE: DNA
<213> ORGANISM: Mycobacterium avium

```

```

<400> SEQUENCE: 16

```

```

atgcccgccg gctgccacca tggccctatg cgattcggat gtgcccgggc ggtcaacctc    60
ggcctgctgc cggttctggt cgtcgtgctc gccggtgcc tcgcggcgcg gcacgcgctg 120
ggaacgcctg actcccaatc gattccggtc gggccatcca cccacacct gcagtcgggc 180

```

-continued

ggcacgcccc gcagctatca cctgtaccgg ccgacggggc tgagcgaggc ggccccctg	240
gtggtgatgc tgcacggcgg ctttggcaac ggcgagcagg ccgagcgcg ctaaccactgg	300
gacgccgagg ccgacgcgg gcatttcctg gtgcgctacc ccgacggcct gggccgcgcc	360
tggaaacgcc gaacctgttg cggtgagccg gcacacgcc gcaccgacga cgtcggattc	420
gtcaacgcc tggtcggcgc catcgcgggc cagatcccg tggatcgtgc ccgcgtctac	480
gtcaccggca tgtcgaaegg cgccatgatg gcgctgcggc tgggctgcca gagcgacacc	540
ctgcgcgga tgcccccgt ggccggcacc ctgctcacc attgctccg ggcgcggccg	600
gcctcggtgc tgcagatcca cggcaccgcc gacgaccgc tgcctatgc gggcggaccc	660
ggaaaggcgt tcgcgctcaa cggctcccc cgggtcgacg ggcgctcgg ggaatccgtc	720
aacgccacct ggcgcgccat cgacgcctgc gggccgcca gctcgaccac cgccggcgat	780
gtcaccacc agaccgcgg ctgcgcggac ggcgcgacg tggagttag ctcggtggcc	840
gggtgcggcc accaatggc cggcggggcg ccagcccc tggcgaaaa ggtggccgga	900
attccggcgc cgtcgacggc gctggacgcc accgacaga tctggcaatt ctccgccgt	960
aatcacggtt ag	972

<210> SEQ ID NO 17

<211> LENGTH: 702

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 17

ttactgcatg cgcagcacga atcccactcc gcggacgggtg tgcagcagcc ggggcccgcc	60
gttggcctcg agcttgccgc gcaggatatc gatgaacacg tcgacgacgt tgggtgcggc	120
ggcgaagtca tagccccaca ccagctccag cagttgtgcg cgggacagca cggcggtctt	180
gtgctcggcc aggaccgcca gcaggtcgaa ctcccgttg gtcaaatcga cgtccacgcc	240
gttgaccccg gcgcggcggc cgggatgtc gacctccagc gggcccaccg tgatcgtttc	300
cgacgaggag gtggcggtgg cgccgcggcg gcgcagcagc gccttgacc gcgcgaccag	360
ctcgccagc acgaacggct tgaccaggta gtgctcggcc ccggcctcca ggcgcgccac	420
ccggtcgtcg accgagctgc gggccgacag cagcgacagc gggacgtcgt tgtccatggc	480
gcgcagcgcg gtgacgacg tgacgcgctc cagcaccggc atgttgatgt cgagcacgat	540
cgcgtccggg cgtgtctcgg tggcgtgcg cagcgctcgc gcgcgctga ccgcggtgga	600
cacctgaat ccggacagtc gcaggccacg ttccagcgac gcgagcacat cggagtcgtc	660
gtcgacgacc aagaccgcg gtgagctagc tgctgtgtcc at	702

<210> SEQ ID NO 18

<211> LENGTH: 417

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 18

gtgagaccag tgccgaccgg caaggttaag tggtagcagc ctgacaaggg cttcggcttc	60
ctgtcccagg aggacggcga agacgtgtac gtccgctcgt cggcgttgcc cgctggcgtc	120
gaggggctca aagcgggcca gcgcgtggaa ttccggcatc cctccggccg ccgcggaccg	180
caagcgttga gcctcaagct gattgagccg ccgcgagcc tgaccaaggc gcgcgcgag	240
gggcctgcc aacacaagca cagccccgac gagctgcacg gcatggtcga ggacatgac	300
acgctgtcgg aaagcaccgt ccaaccgag ctgcgcaagg gccgctaccc cgaccgcaag	360

-continued

acgcgcgcgc ggggtctcga ggtggtcagg gcgggtggccc gcgagctgga cgctga 417

<210> SEQ ID NO 19
 <211> LENGTH: 1122
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 19

tcagaccgcg ctcaggtgcg cgcgcagcgc cgagatcaac tcgctggtgg ccaccatgct	60
gtcgccgcgc agctggaaca ggttggcgaa gccgtgcgtc agcgaaccca ggtaccgcag	120
gtccaccggg gtgcggcgcg cgcgcagcgc ctgggcgtag ctctggccct cgtcgcgtag	180
cgggtcgaag ccggccaccg cgatcagcgc cggggccagg ccggccacgc attcggccag	240
cgcggcgcat acccgcggtt ccgtccggtc cagccgcgag ttccgcaggt actgcgattc	300
gaaccagtcg atgtcgcgct tggtcagcag gaagccgcgc gagaacaggc tcagcgaccg	360
ggtccgggcg gtgaagtcgg tgcgcgggta gatcagccac tgcagcacgc gtgccgggccc	420
gccccgcgtc cgggcccagct ggttgacgac ggccggccagg ttgcgcgcgc cgctgtcgcc	480
gccgaccgcc acccgcccgc ggatcgcgcc cagctcgccg gcattgctcat gcgccagggt	540
gaaggcgggc tacgcgtcgt cgatggcggc cggcgcgggg tgctcgggcg ccagccggta	600
gtcgatcgac agcacgtgga tgcggcggtc gccgcaggtc agccggcaca gcgagtcgtg	660
ggtgtccagg tcgccgatcg accagccgcc gccgtggtag aagaccagca gcggggcctg	720
ggtctcgccg ccggcgggcc gatagtccg cgcggggatg tcgccggcgc ggccgggcag	780
cgcagttcgc gcgacgtcga cgtggatctg cgggcggggg aacccccgcg tggactcgtg	840
catctgggcg cgggacagct cggggtcgtc gtcgaccacc aggcgcgtcg tgcgacggc	900
gcgcagaccg gacagcatca gctgcagggt cgggtcgagc gtgttgccgt cgatgatgac	960
cgagcgggcg cgcaccaggc cgcggcgcac ggccgtcggg atccacggga tgacctgac	1020
cccagcgtg gtgacggcgc cctggatcg gttggtcagc ggcatcgacc cgcgctgcgc	1080
gcgggggtcg acgggtgcgg tgcgggtcag cggttggtc at	1122

<210> SEQ ID NO 20
 <211> LENGTH: 918
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 20

atggatcttc atcagctgga gtgctttatc gcggtcgcgc aggaaggcac cttcaccgcc	60
gcggcacagc ggatacatct ggcgcaatct ggtgttagtg cacacattaa ggcgtcgaa	120
cgcgaaatcg ggcagcagct attcgaacgg ccggccacgc cggtagcggt tacgcgggt	180
ggaaacgcgc ttctgccata cgcgcgcgca gcaactcgatg cgctggccgc aggtcgcgcc	240
tcgatcgacg gactcaccgg tctactacat ggtcgccctg ccatcggcac catcacctcg	300
atctcaccgc gcagcatcga cctccccgag cttctagcag cgtttcacca tgagcaccg	360
ggggtagacc tgtcgctggt tgaggacact gcagcgatgc tcaatcgcca catatccaat	420
ggcgctttag acgtagcttt caccagcctg acggatgagg cagtagccgc cgtcgcgatg	480
cgcgagttgc atcgggagcc ggtgatcgcg atctttctac cgctccgatcc gttatctcct	540
tgccggaagc tcacattggc cgagctggcg gatagaccgc tcatcacgct cccggaggga	600
tcaggcttgc gctggcaact caaccgcgcg ctgcggcggg ccggcggtca agcccacatc	660

-continued

gccttcgaag cggcgatcc cgacgtactc gttgcgctcg ttgcgaaggg gctgggctg	720
ggtctcgttc ctcaatcagc cctcgcgcaa agcgatcagc taataggatt gccagtcagc	780
gatcatccgc cggggcgctt aggcattatc tggccagaag ggcaagccgc cagcccagcg	840
gctcgtgcct tcgttgaaca cgccaccacg gcgacaacta aacttcggcg accggccgag	900
ctacgccagg atcgctga	918

<210> SEQ ID NO 21
 <211> LENGTH: 1530
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 21

ttggccgatg accccgggtc aagcttcacc acggtgtgga atgcggtcgt ttcggagctc	60
aacggcgagc cgtcgccga cggcggagcc gccaacgca cgactctggt cactccctc	120
acccctcagc aaagagcgtg gctcaatctg gtccgcccgc tgaccatcgt cgaggggttt	180
gctctgctgt cgggtccgag cagtttcgtg cagaacgaga tcgaacggca cctgcgcgcc	240
ccgatcaccg acgcgctcag ccgcccctg ggtcagcaga tccagctggg agtccgcac	300
gtccccccac ccgacgagc cgaggacgcg cccatcccgc cggccgagcc gttccccgac	360
accgacgccg cctgttccgc cgacgacggc gccgacggcg agccggtgga gaacggggag	420
ccggtcaccg acaccagcc cggctggccc aactacttca ccgagcgcc gcacgccatc	480
gateccgcgc tcgcgcgcgc aacgagcctc aaccgcccgt acaccttcga cactgtctg	540
atcggggcgt ccaaccggtt cgcgcacgcc gccgccttgg ccacgcgca agcaccggcc	600
cgcgcctaca acccgttgtt catctggggc gagtccggtc tgggcaagac gcacctgctg	660
cacgcgcgcg gcaattacgc gcaacggctc tccccggca tgcgggtcaa gtacgtctc	720
accgaggaat tcacgaacga cttcatcaac tcgctgcgcg acgaccgcaa ggtcgccttc	780
aagcgcagct atcgcgagc cgacgtgctc ctggctcgatg acatccaatt catcgagggc	840
aaggaaggca tccaggagga gttcttccac accttcaaca cgctgcacaa cgccaacaag	900
cagatcgtca tctcctccga ccggccgccc aaacagctgg ccacctgga agaccgctg	960
cgaacccggt tcgagtgggg cctgatcacc gacgtgcagc cccccaact cgaacccgc	1020
atcgcgatcc tgcgaagaa ggcgcagatg gagcgccctg cggtgcccga cgacgtgctg	1080
gaactcatcg ccagcagcat cgagcgcaac atccgcgaac tcgagggcgc cctgatccgg	1140
gtcaccgcgt tcgcctcgtt gaacaagacc ccgatcgaca agtcgctggc cgagatcgtg	1200
ctgcgcgatt tgattgcga cgccagcacc atgcagatca gcgcggccac catcatggcc	1260
gccaccgccg aatacttcga caccaccgtc gaggaactgc gcgggccggg caagaccgg	1320
gcgctggccc agtcccgcga aatcgcgatg tacctgtgcc gcgagctcac ggatctgtcg	1380
ctgcccaaga tcgggcaggc cttcgccgcg gaccacacca cggatgatga cgcccagcg	1440
aagatcctgt ccgagatggc cgagcgacgc gaggtgttcg atcacgtcaa ggagctcacc	1500
actcgcatto gtcagcgtc caagcgtga	1530

<210> SEQ ID NO 22
 <211> LENGTH: 1200
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 22

atggacgcgc cgacgacaac ggctggcctc agcgacttga agtttcgttt agtcggggag	60
---	----

-continued

tctttcgccg acgcggtgtc gtgggtggcc aagagcttgc cgtcgcgacc cgcggtgccg	120
gtgctctccg ggggtctgct gtccggcacc gacgaggggc tcaccatttc cggattcgac	180
tacgaggttt cgcgcaagc acaggtggcg gccgaaatcg cgtctccggg aagtgttttg	240
gtatccgggc ggttgctgtc tgacatcggt cgggcgctgc ccaacaagcc gatcgacttc	300
tacgtcgacg gcaatcgggt ggcgttgaac tgcggaagcg cccggttctc gctgccgacg	360
atggccgctg aggattaccc gacgtgccc acgctgccc aggagaccgg gacgtgccc	420
gccgatctgt tcgccgaggc gatcgggcag gtccgcatcg cggccggccg cgacgacacc	480
ttgcccattg tgaccggaat ccgggtcgag atttcggggg acacggtggt tttggccgcc	540
accgaccggt tccggtggc ggttcgcgag ctgacctggt cggcgccctc ccccgacatc	600
gaagcggcgg tcctggtgcc ggccaagacg ctggccgagg cggcgccgac cggcatcgac	660
ggttccgacg tgcggctgtc gctgggggcg ggccgagggt tcggcaagga tggcctgctg	720
ggtatcagcg gcaacggcaa gcgcagcacc acccgcttgc tggacgcgga attcccgaag	780
ttccgccagc tgttgccggc tgagcacacg gcggtggcca ccatcaacgt cgcgagctg	840
accgaggcca tcaagctggt ggcgtggtg gccgaccggg gcgcgcaggt gcgcatggaa	900
ttcagcgagg ggtcgctcgc gctgtccgcc ggccgacgac atgtcggccg ggccgaggag	960
gatctggccg tggatttcgc cggcgaaccg ctgacgatcg cggtcaacct cactatctg	1020
accgacggac tgggatcggg gcgtccgaa cgggtgtcgt tcggcttcac caccgccggc	1080
aagcccgcac tgctgcgccc ggctccgac gacgacagcc cggcgagcgg cagcgggccg	1140
ttcagcgccg tgcccaccga ttacgtctac ctgctgatgc ccgtcgggtt gccaggttag	1200

<210> SEQ ID NO 23

<211> LENGTH: 1158

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 23

gtgtacgtcc ggcatttagg actgcgcgac ttccggtcct gggcacacgc cgacctcgaa	60
ctgcagccgg gtccgacggt cttcatcggg tccaacggct tcgggaagac gaattctgctt	120
gaggcgctgt ggtattcgag cagctgggg tcacaccggg tgggcacgga cgcgccgttg	180
atccgcgccc gcgcgacgg ggcggtggtg tcgaccatcg tggtaacga cggccgggaa	240
tgtgcggctg atctggagat cgcgcgccgc cgggcgaaca aggcgcgggt gaaccggtca	300
cccgtagcga gcaccgcga ggtgctcggc gtgctgcgcg cgggtgctgt cgcgccgag	360
gacctggccc tggtagcggg ggtacctcc gagcggcgcc gttacctga cgacctggcg	420
acgtcgggc gcccgcgat cgcgcgggtg cgcgcgact acgacaaggt gttgcggcag	480
cgcaccgctg tgcataaat gctgtccggt gcccgccacc gggcgacgg cggcgctctg	540
gacacctcg acgtgtggga cagccgctg gccgaatacg gggccaatt gatggctgcc	600
cgaatcgatt tggtaacca gttggcgccg gaggtggaga aggcctatca gctgctggcc	660
ccgggatcgc gggcgcgctc gatcggctac cgatccagcc tgggcgcggc ggcctcggcc	720
gaggtgaacg cggcgacgg cgactatctg gaggcgcgcg tgctggccgg gttggcgcc	780
caccgggacg ccgaactgga acggggcatg tgctggctg gcccgaccg cgacgacctg	840
gagctgtggc tcggtgagca ggtggcgaaa ggttttgcca gccatgggga atcgtggtcg	900
ctggcgctgt ccctcgggct ggccgccttc gagttgctgc gggccgacga aagcgatccg	960

-continued

gtgttgctgc tgcagcacgt gttcgccgag ctgcagcccg cccgccgccc ggcactggcc	1020
gcgggtggccc aatccgccga acaggtgttg gtcaccgccc cggtgctcga agacatcccc	1080
acgggctggc aggtctggcg gctcttcgtc gagttgcgcg acaccgacgc gggccgggta	1140
tcggagctgc gcccatga	1158

<210> SEQ ID NO 24

<211> LENGTH: 501

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 24

atggacctgg tgcggcgac cctcgaggag gcccgggccc cgccccgggc acagggaaaa	60
gacgccggcc gcggggcgcc cgcagcacc acgccgccc gggtggcggg tcagcggcgc	120
agctggtcgg gaccgggacc cgacgctcgc gaccgcacac cgctgggccc gctggcgcgg	180
gacctggcca ggaagcgggg atggtcggcc caggtcgcgg agggcacctg gttggggaac	240
tggacggcgg tggtcgggtc ccagatcgcc gaccacgcgg tccccaccgg tctgcgcgac	300
ggtgtgctga cgtgttcgc cgagtcgaca gcttgggcca ccagtttgcg gatgatgcag	360
gcgcaactgt tggccaagat cgcgcgcgcg gtcggcaacg gggtggtgac ctgcctgaag	420
atcacccggc cggccgccc gtcctggcgc aaaggcccc ggcacatcgc cgggcgcggg	480
ccgcgcgaca cctatgggta g	501

<210> SEQ ID NO 25

<211> LENGTH: 2034

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 25

gtggctgccc agaagaagaa ggcgcaagac gaatacgggtg cttcagcgat caccgtcctg	60
gaaggactgg aggcgggtccg caaacgcccc ggcattgtaca tcggctctac cggcgcgcga	120
ggtctgcacc acctcatctg ggaggtggtc gacaaactcg tcgacgaagc gatggccggc	180
taegccgacc gggctgcagc gcggatcctg gacgacggca gcgttgaggt cgcgcgacaac	240
ggccgcggca tccccgtcgc gatgcacgcg accggcgcgc ccaccgtcga cgtgggtgatg	300
acgcagctgc acgccggcgg caagttcggc ggcgaaaaca gcggtataaa cgtcagcggc	360
ggtctgcacg gcgtcggcgt ctcggtggtc aacgcactgt ccaccggtc cgaggtcaac	420
atcgcgccg acggctacga gtggtcgcag tactacgacc acgccgtgcc cggcacccctc	480
aaacagggcg agggccaccaa gcgcaccggc accaccatcc ggttctgggc cgaccccgac	540
atcttcgaga ccaccagta cgacttcgaa acgggtggccc gacggctgca ggaatggcg	600
ttcctcaaca agggcctgac catcaacctc accgacgagc gggtagacaa cgaagaggtc	660
gtcgacgagg tggctcagca caccgccgac gcacccaagt cggcgcagga gaaggcggcg	720
gaatcggctg cgcgcgataa ggtcaagcac cgcaccttc actacccgg cggcctggtc	780
gaattcgtca aacacatcaa tcgcacaaaa aaccccatcc accagagcat catcgatttc	840
ggtgggaagg gccccggcca cgaggtcgag atcgcgatgc agtggaacgg cggtatttc	900
gaatcggctg acacgttcgc caacaccatc aacacgcacg agggcgggac ccacgaggag	960
ggcttcgcga cgcggttgac ctccgtggtc aacaagtacg ccaaggacaa gaagctgctc	1020
aaggacaagg accccaacct caccggcgac gacatccgcg agggtttggc cgcggtgac	1080
tcggtcaagg tgagcgaacc gcagttcgag ggccagacca agaccaaact gggcaacacc	1140

-continued

gaggtgaagt cgttcgtgca gaaggtgtgc aacgaacagc tcacccactg gttcgaagcc	1200
aaccccgagc acgccaagt cattgtcaac aaggcggttt cgtcagcgca ggcgcgatt	1260
gccgcgcgca aggcgcgaga gttggtgcgc cgcaagagcg caaccgacct gggcgggctg	1320
cccggaagc tcgcgactg ccggtcgacc gatccgcgca agtcggaatt gtatgtggtc	1380
gagggtgact cggccggcgg ctcggcgaaa agcggccggg actcgatgtt ccaggccatc	1440
cttcgcgtgc gcggcaagat catcaacgtc gaaaaggccc gcatcgaccg ggttttgaag	1500
aacaccgaag tgcaggcgat catcacgcgc ctgggcaccg ggattcacga cgagttcgac	1560
atcaccaagc tgcgtacca caagatcgtg ttgatggcgc acgcgcgact ggacggccag	1620
cacatctcga cgctgttgtt gacgctgctg ttccggttca tgcggccgct gatcgaacac	1680
gggcacgtgt tcttgcccca gccaccgctg tacaagctga aatggcagcg cagcgatcca	1740
gagttcgctt actccgaccg cgagcgggac gggctgctcg aggcggcct gaaggccggc	1800
aagaagatca acaaggacga cggtatccag cgctacaagg gtctgggcga gatggacgcc	1860
aaggaattgt gggaaaccac aatggatccc accgtcgggg tgcctgcgca ggtcacgctg	1920
gacgacgccg cggccgcgca cgagctgttc tccatcctga tggcgagga cgtcgacgcg	1980
cgccgcagct tcacacccg caatgccaaa gacgttcgct tcctagacgt ttaa	2034

<210> SEQ ID NO 26

<211> LENGTH: 2520

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 26

atgactgaca ccacgtgcc acccgggcgt gacgcgcgcg accgcgtcga accggtcgac	60
atccagcagg agatgcagcg cagctacatc gattacgcga tgagcgtaat cgtcggccgc	120
gcgctgcccc aggtgcgcga cggcctcaag ccggtgcacc gccgggtact ttacgccatg	180
tacgactcgg gtttcgccc ggaccgcagc cagcccaagt cggcgcggtc ggtcgccgag	240
acgatgggca actaccaccc gcacggcgac gcctcgatct acgacaccct ggtcgggatg	300
gcccagccgt ggtcgctgcg ctaccgctt gtcgacgggc agggcaactt tggttcgccg	360
ggcaacgacc cgcgggccgc gatgcggtac accgagggcg ggctgacccc gctggccatg	420
gagatgctgc gcgaaatcga cgaggagaca gtcgatttca ttcccaacta cgacggccgg	480
gtgcaagagc cgacgggtgt gccacgccg ttcccaacc tgctggccaa cgggtcgggg	540
ggcatcgcgg tcggcatggc cacgaacatc ccgccgcaca acctcggcga gctcgccgag	600
gcggtgttct gggcgctgga caattacgag gccgacgaag agggcacccct ggcgcgcgtg	660
atggaacggg tgaaaggacc cgacttcccg acctcgggcc tgatcgtcgg cagcagggc	720
atcgccgacg cctacaagac cggccgcggt tccatccgga tgcgcggagt cgttgagggtg	780
gaagaggatt cgcgcggccg cactcgtctg gtcacaccc agttgccgta tcagggtcaac	840
cacgacaact tcacacctc gatcgccgag cagggtgcgc acggcaagct ggcggccatc	900
tccaatatcg aggaccaatc cagcgaccgg gtcgggctgc gcatcgatc cgagctcaag	960
cgcgacgcgc tcgccaaggt ggtgctgaac aacctctaca agcacacca gctcgagacc	1020
agcttcggcg ccaacatgct ggccatcgtc gacggggtgc cgcgcaccct gcggctcgac	1080
cagctgatcc gccactacgt cgaccaccaa ctcgacgtca tcgtcggcg caccacctac	1140
cggttgcgca aggccaacga gcggggccac atcctgcgcg gtctgggtcaa ggcgctcgat	1200

-continued

gcgctcgacg aggtcatcgc cctgatccgg gcgtcggaag ccgtcgacat cgcgcggcag	1260
ggcttgatcg agctgctega catcgacgag atccaggcgc aggcgatcct ggacatgcag	1320
ctgcgcgggc tggccgcgct ggagcggcag cgcacatcgc acgacctggc caagatcgag	1380
gccgagatcg ccgacctgga ggacatcctg gccaaagccgg aacggcagcg cggcatcgtg	1440
cgcgacgagc tcgccgagat cgtcgaaaag cacggcgacg cgcggcgcac ccgcatcgtg	1500
gccgcggcag cgcagcttag cgacgaggat ctgatcgctc gcgaggacgt cgtcgtcacc	1560
atcaccgaga ccggctacgc caagcgcacc aagaccgacc tgtaccgcag ccagaagcgc	1620
ggcggcaagg gcgtgcaggg cgcgggcctc aaccaggacg acatcgtgcg gcacttcttc	1680
gtgtgctega cgcacgactg gatcctgttc ttcaccaccc agggcggggt ctaccgcgc	1740
aaggcctacg aactgcccga ggctgccgc accgcccgcg gtcagcacgt ggccaacctg	1800
ctggcgcttc agcccagga gccgatcgct cagggtgatcc agatccggag ctatgaggac	1860
gtccctacc tgggtgctggc caccgcgaac ggctgggtga agaagaccaa gctgaccgac	1920
ttcgactcga accgctcggg cggcatcgtg gcgatcaacc tgcgcgacaa cgacgaactc	1980
gtgggcgcgg tggtgtgctc gcccgaggac gatctgctgc tgggtgctggc caacggccag	2040
tccatccggt tctcggcgac cgacgaggcg ctgcgccga tgggcggcg cactccggt	2100
gtgcagggca tgcgcttcaa cgcgcgacgac tacctgctgt cgctcaacgt ggtccgcgag	2160
ggcacctacc tgctgggtggc gacgtccggc gggtagccca agcgcaccgc gatcgaggag	2220
tatccggtgc agggcggcg cggcaagggc gtgctgaccg tgatgtatga ccgccggcgt	2280
ggcaggctgg tgggtgctgc gattgtggac gaggacagcg agctgtacgc gatcacctcc	2340
ggcggcggtg tcacccgcac cgcggcgggc cagggtccga aggcgggacg gcagaccaag	2400
ggcgctccgc tgatgaatct gggtaggggc gacacgctgc tggccatcgc tcgcaacgcc	2460
gaggaagccg cggacgaggc cgtcgacgag agcgacgggt ccgcggggtc ggacggctag	2520

<210> SEQ ID NO 27

<211> LENGTH: 852

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 27

gtgagttcac cgaacgagcc gggcgacccc aagaaggggg agacccccaa cggggacggg	60
tcggcagagc gggccggggg gcggcgggcg acaccgccgg cgcggggtgg ccgcgcccc	120
gaggccggcg acggaccgcc gtggcagcgc ggcagacccc gggcccagca gcccgcgcg	180
cgcagaaacg agccggccac cgagaagcgg gccgacggcg cgggggcccga ggcccggctg	240
aaaccgttca tctcgggcac cgcgtcgac gccaaaggag cggagcgcgg tgaggggccc	300
cccgcgagg cctacgccag cgaactgccc gacctgtcgg ggcgggtgcc gcgcggccca	360
caccgcaagc cggccgcccga gcgcgggtgcc gagacgaccg cggggcaagg cgggtggcgt	420
ccggtgagcg gcgagagcgg cgagggccgg gacaaccggg accgggtgca ggtgtcgcg	480
cgcaccccg cgcgggttcg cgcacgatg cagatccgcc ggatcgaccc gtggagcacc	540
ctgaagggtg cgctgctgct ctcggtggcg ctattcttcg tctggatgat cgcgctcgcg	600
ttcctctacc tgggtgctcg tggcatgggc gtgtggtcga agttgaacag caacgtcggg	660
gaactgctga acaacaccag cggcagcagc gggaattgg tgtccagcgg caccattttc	720
ggcggcgccg tgctgatcgg cctgggtcaac atcgtgctgt tgaccgcgat ggcgaccatc	780
gcggcggtta tctacaacct ggccaccgat ctgatcggcg gcacgagggt caccctggcc	840

-continued

gatcgcgact ag 852

<210> SEQ ID NO 28
 <211> LENGTH: 579
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 28

ctaggctcca caaatcgaat ccctctcccc gcccgggcgg tacgacgcgc cgaggaagcg	60
gtgcaccagg gcgcgctcgg cggccggggtc cttgagcggc cagccccata acgccaggaa	120
gacgcggatc agccactgcg ccgccagcgg gtctgtcgtgg ccgggcccga gcatctcggc	180
ggccaggggc gtcaccgtgg gcgagttgat caccagtcg ctgaccgggg ccgcgtggat	240
cgaccgcacg agctgagcca gcggatcgga gcgcaaccgt tccagcgcca tgatcgtggc	300
ggtcaccacc cgctcggggc cgcgcaggtc tttgatggcc tcccgcgtgg tgctgacgat	360
gcgggcgggc tggataccga cgacggcatc gcggatcgtc gccttgccgc cggcgtgccg	420
gtagatggtc gccggtgaac agtgaccccg ggccgccagc gcctcgatcg tgaaccgcgc	480
gtagccgcgg cgccgatga ggtcggcggc cgcggcgtag atccgctcgg cgcccgcgct	540
gcgccgatcc cggcccacca accagtcgtc ccgcgccat	579

<210> SEQ ID NO 29
 <211> LENGTH: 1326
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 29

ctgtctcgca cgggtcaccg gcaccgcag gcgccgtccc cggccgtcgt cgctcctcaa	60
cttcgcccg gcgccgttac gccgcgagac ggccgcgtcga gcctgaaacg catggcggtg	120
ttggacacgc cgaccgggtt cttcggcagc gaggcgatac aggatcccta cccgctgtat	180
gagcgcatgc acgccgaggc gccgatgcac cggatcggcg actccgtctt ctacgccgtg	240
tgccgggtgg acgcgggtca cgaggccatc gagcgggtcg aggacttctc gtccaacctc	300
accgcgacga tgggtcttcca cgaggacggc accgtcaccg cgttcgacat gggcgcgccg	360
ggcgccccga tgcacgcgct ggccaccgcc gacgaccccg tgcacgcggt gcaccgcaag	420
atcctgctgc cgcaccttcc ggcaagcga atccggatca tcgaggagtt cgccaccag	480
accgcggacc ggctgtggga cgagaacctg tccgacggcc ggatcgagtg gatgagcgcc	540
atcgccaacc gggtgcccac gatggtggtc tgccggctgc tgggtctgcc cgatgacgat	600
gtcgacaaa tcacccggct cggctacgcc accaccacac tgctggacgg gatcgctgcc	660
cccgaacagc tcgagcaggc cggcatggcc gccatcgaac tgtccggcta cgtgctggag	720
catttcgaaa aggcaagcga gaagccgga tccagcctga tggccgatct tgcggcgcg	780
tgcccgccg gcgaactcga gcagctccg gcgctgggaa tcatgctcac gctgttcagc	840
gcggccggcg aatcgaccgc gtcgctgctg ggcagcgcg cctggatcct ggccgaccgg	900
ccggcgatcc aacggcagct tcgcgaaaac ccggagttgc taagcacttt catcgaagag	960
acgtgcgct tcgaggcgcc gtttcggggt cactaccgcc acgtgtggcg cgacaccag	1020
ctgggcggga tcgagctgcc cgaggcgca cactgctgc tgatgtgggg agcggccaac	1080
cgcgacccga cacatttcaa ggatcccaac gagtttcggc tcgacccgcg ggccgccaag	1140
agccacctga gcttcggcaa gggcgctccac ttctgcgtcg gtgccgcgct ggcccggtcg	1200

-continued

gaagcccaca tcgtgttgcg ccggtgctc gagcgaccca gctggatcga cgcgaccgac	1260
gtcggggatt ggctgcccag catcctggtc cggcgccggg aacggctggg gttggccgtg	1320
cgtga	1326

<210> SEQ ID NO 30
 <211> LENGTH: 435
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 30

ctacggccgc ggttgacgt cgacgctcgg cggccgcgac tgcggctcct gcggttctc	60
gcgcaccgag cgccgcacga tgctgaacgc ggcgccgccc ccggcgagaa cggcgacggc	120
gacaccggcg atgaccacg ggcgttgcc ccgtcgctgc gcgcgccgag cgtcctgcag	180
cgctgggggc aggcgggtga ccacctctg agcggcgggc agttcctggg cgagggctc	240
ctggggccgc gcgaggtctc gggccaggcg cccctcccg taccggcgcc gcagctgctc	300
ggcgtgcac cgccgggact gcacgctcag gccggcgact ccccggtga ggtccagcgg	360
tccagtgtc gagtaggtca ggccgcgggc cagtcgctct cgccgggtca gccggtcttc	420
cgcttagcg cgcat	435

<210> SEQ ID NO 31
 <211> LENGTH: 549
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 31

gtggcagact ctgatgccgt gactaacagc ccctttcaga ctgctaccgc cagctgcac	60
accaaccgag gtgacatcaa ggtgcacctg ttgggaaacc acgcgcccaa gaccgtggcc	120
aaattcgtcg gctggccca gggcaccaag gactactcga ccagaacgc gtcgggcggc	180
tctcggggc cgttctacga cggcgcggtg ttccaccggg tgatccggg cttcatgatc	240
cagggcgggc acccgaccg caccggccgc ggcggtccgg ggtacaagtt cgcgcagag	300
ttccaccccg agctgcagtt cgaccggccg tactgctgg cgatggccaa cgcggggccg	360
ggcaccaacg gctcgcagtt cctcatcacc gtggacaaga ccccgacct gaaccggcgg	420
cacaccatct tcggcgaggt cgtcgaccg gagtcgcaga aggtcgtcga cgcgatctcg	480
acgaccagca ccgacggcaa cgaccgccg tccgagccgg tggatgatga gtcgatcacc	540
atctcgtag	549

<210> SEQ ID NO 32
 <211> LENGTH: 426
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 32

tcagcgctgt cgcgggcccg cgtaaccggc ggcggtgagc gcgtcgagca cctggagcgg	60
gtccgccccg agatcccagc gggacagcac caccaggtct ccgtcggcgc ttccacctc	120
gagcaaccgc accttgccgc cgtagcggcg gaactcgatg atccggatga tcttgatgtc	180
ggggcgttgt aatacctcgc tccgatacca gccgcggacg gccagccctg cgggtgtgat	240
tgccagcttc gggcgcgcgcc gccacgacat ccccgcaaac acgagcagac ccgcccggc	300
aacgccagcc agaacacgcc ccggcggggc tgtgaccacg gtcacagccg cgatagccat	360
cagaacgccc gcaactccac aaccagcgat tcccgccgga tgcggctccc accgtgtttg	420

-continued

ctgcat 426

<210> SEQ ID NO 33
 <211> LENGTH: 282
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 33

tcagtgccag cgcacgtga gcaacaaacc ggtgatcatg aaagcgaacg cgattgcgta	60
gttccacggc cccaattggg ccatccaatt gagggcggtc ggcgcctggc tgcccacgcg	120
ggccagttgg aagaccatca gccacaccaa tccgatcagc atcagcccga tgaacagcgc	180
gacgaaccac acgctggacg gtcccacctt gaccttcacc ggcgtgcggc tgaccgcgct	240
gacgggtgaag tcgttctttt tgcggacctt ggacttgggc at	282

<210> SEQ ID NO 34
 <211> LENGTH: 744
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 34

atggtccaga cgcgcagtc gccatggcgc ttcggcgctc cgctggctctg cctgctggcc	60
ggattgctgc tcgcgcac gcacggcggtg tccggcggtc ccgagatacg ccgcagcgac	120
gcaccccggc tgggtggacct ggtgcgcgag acgcaggcgt cgggtgaaccg gctcagtgcc	180
cagcgcgaac agctcgccgc caagatcgac gccgcgcacg gccgctctgc ggatgccgcg	240
ctggccgcga tgctgcgcgc ctccgcgcag ctgcgcggcg aggcgggat gagcccggtg	300
caaggggcgc gcctgggtgt caccctgcag gatgcgcagc gcgacgcaa cggccgcttc	360
ccccgcgacg cgtgcgccga cgacctggtg gtgcaccagc aggacatcca ggccgtgctc	420
aacgcactgt ggagcgcgcg cgcgcaggcg atccagatgc aggaccagcg gatcatcgcc	480
acgtcggtgc cccgctgcgt cggcaacacg ctgctgctca acgggcgcac ctacagcccg	540
ccctacacga tcaccgcgat cggcaacgcc gccgccatgc aggcgcgcct ggccgcggct	600
ccccgtgtga cgctgtacaa gcagtagcgc gtccggttcg gcctgggcta tcaggaagag	660
gtccgctccg atgtgcaggt ggtcgccac ttcgagcccg accggctgca ttccgcccag	720
cccaacggcc cgatcggtta cttaa	744

<210> SEQ ID NO 35
 <211> LENGTH: 687
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 35

atgcggatcc tgggtgtcga caactacgac agcttcgtgt tcaacctggt gcagtacctc	60
gggcaactgg gtgtggacgc cgacgtctgg cgcaacgacg acagccggtc cgccgaccac	120
aacgcgcagg cccgcgggtt cgacgggggtg ctgctcagtc ccgggcgggg caccgccgag	180
cgcgcggggc cttccatcgc gatgggtgcgc gcctgcgcgc ccgagcacac cccgctgctg	240
ggggctctgc tgggccacca ggccatcggc gtggcattcg gcgccaccgt ggaccgcgct	300
ccggaactgc tgcacggcaa gaccagcagc gtgcgccaca cgaatgccgg tgtgctgcaa	360
gggcttccgg accccttcac ggccaccggc taccactcgc tgaccatcct gcccagatcg	420
ctgcgcggcg tgctcgaggt caccgctcac accgacagcg gtgtgatcat ggcgctgcgc	480

-continued

cacacgcagt taccgatcca cggcggtccag ttccaccgag agtcgatcct gaccgagggg	540
ggtcatcgca tgctggcgaa ctggtccacc gaatgctggc ggggtcgcca cgacacctg	600
gtgctggcgc tggaaaaaga cgtccacgcc gcggtgagac cgtatctccc ggccgatccg	660
gccgctactg accgaacttc agcgtga	687

<210> SEQ ID NO 36

<211> LENGTH: 1881

<212> TYPE: DNA

<213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 36

ctactgaccg aacttcagcg tgatgatccc gtcccggttg accccggccc cggccggcgg	60
gttctgatac accaccgggt gggactgggc gccgcccggc tcgacgtcgg gacccttgtc	120
caggatcccc gtccagccca gcgcccgtag ccgcggttcg gcgtcggccc aaaacatccc	180
ggacaggtcg ggcgatgatga actgattgcc cttggacacc tgcagttcga tcaccgaatc	240
caccggaacc gtttgtccct tgggctgggt ggtgcccgat acctcgccgg ccgggcccgg	300
gctgtccacc tgcacctggg tgatcttggg gaagccgtag acggtcaggc tcttctgcgc	360
gatgtcgacg gtttgcctcg cgatgtcggg cacctgcttg gtctccggcc ccgagccgac	420
gatgaggggt atgacgttgg tgatcgccga cgtctggttg gccggcgggt tgggtccgat	480
caccttgccc aacaactcgg gtgtggacgg cgagttcgcc tgcttgaact tgctgaaccc	540
ggcggccttg agcttggtga ccgctcgga gtagctcagc gacgagacgt cgggcacctc	600
gcgtgctccc ggcccgggtg agacgttgat ggtgatctcg tcgcccgcgc tgaccgacgc	660
gttgccgccc gggctcgtgc cgatgacgtg gtccggcggg atggccaggt ccgcttctcg	720
cagggtgcgg gtcttgaagc ccgcttctcg cagcggcgca atggcatcgg cggacacctg	780
cccgccgacg tcgggcacct ggacgtcgcg gccgcccggc ccgaagggtg tgaacgcgat	840
gacgacgatg atggtcagga ccgcccgcgc gccaccgcc acgatccaac ggcccaccga	900
acccaccgtg ccgtctctgc cgctgtccgc gagcaactgg cgaggcagcg gatcggtgcg	960
tgacggaccc gccgccccgg caccggacga caacagcgag ctgcgctcgg cgtcggtgag	1020
aacctttggc gcttccggct tttcgccgtt gtgcacccgg accagatcgg cgcgcatttc	1080
gaccgcggtc tgatagcggt tgtccgggtt tttggccagc gccttgagca cgaaggcgtc	1140
gaggtcggcg gagatgcctt cgtgccgtg cgacggcggc accgggtctt cgcggacatg	1200
ctggtaggcc accgccaccg gcgagtcgcc ggtgaaaggc ggctcgccag tgaggatttc	1260
gtacagcagc cagcccagcg agtagacgtc cgagcgcgcg tcgacggcgt cgcgcggggc	1320
ctgctcgggc gagaggtact gtgcggttcc gatcaccgcc gcggtctggg tgacgtgtgt	1380
gccgctgtcg gcgattgcgc gggcgatgcc gaagtccatc accttgacgg cattggtggt	1440
gctgatcatg atgttcgccc gcttgacgtc gcggtggatg atgccgttct ggtggctgaa	1500
gttcagcgcc tggcaggcgt ccgcatgatg ctcatggcc gcccgcgcg gcagcgcccc	1560
gtcgggtgtc acgatgtcgc gcagcgtcac gccgtcgacg tactccatga cgatgtaggg	1620
cagcggcccc gagggcgctc ccgcctcgcc ggtgtcgtag acggcgacga tggacgggtg	1680
gttgagcgcg gccgctgttt gcgcctcgcg ccggaagcgc aggtagaaac tgggatcgcg	1740
ggccaggtcc gcgcgcagca ccttgaccgc gacgtcgcgg tgcagccgga cgtcacgggc	1800
caggtgaacc tcggacatgc ccgcaagcc gaggatctcg ccgagttcgt agcggtcgga	1860
caggtgttgc ggggtgttca t	1881

-continued

<210> SEQ ID NO 37
 <211> LENGTH: 510
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 37

```
tcattgcgat gtccagtgtc gggcggtgtc gtcgcggaat ccagacgccc caattgtgccc 60
gtcatccggc cagtcacgcc gcaggcccggt gtgggccccg gagccgctcg gcgttttgggt 120
cgccggcgcc gccgaggcgg gcgggggttc ggtgtcggtg accgtcgggg gctgttgctg 180
ctgggtccgg cgaggattga tgacgatgag caccgcgatg atgatcgcca gcgcgcccag 240
caccggcgcc gccacacgca gcgcccgtg gccggacgag aaggtgcgcc gggccggggg 300
cgggcggtgg ccgcgggtgg ccggccgggt ccgcggcgcc gcggcggttc gcccgaggga 360
cacggctgcc gcccggtggg tggggctgga cggaatggcc gccggtgacg cccgcccggg 420
cgccggggac tggctcgccc gcggcgcccg gcgaccggcg cgcaccggcg cgacggcgctc 480
ggcgaacggc ccccgcgtgc ggtagcgcat 510
```

<210> SEQ ID NO 38
 <211> LENGTH: 762
 <212> TYPE: DNA
 <213> ORGANISM: Mycobacterium avium

<400> SEQUENCE: 38

```
tcacaggggt atctcgatga gctcgcgcac gttgggtggc agctccgagg gcagcgcgcg 60
cgagggctcc ttgatgtgtt tcatcgccac cgtcagcgcc ccattcccgc tgaacggccg 120
cttggccgaa accacctcgt agccaacaac tcccagcgag tacacgtcgc tggcggggggt 180
ggcgctcgtg cccagcgccct gctcgggcgc gatgtattgg gcggtgccca tcaccattcc 240
ggtctgggtc accggcgcgg cgtcgacggc cttggcgata ccgaagtcgg tgatcttcac 300
ctgccccgct ggggtgatca ggatgttgcc cggtttgacg tcgcggtgca ccaggccggc 360
ggcgctcgcg acctcgaggg cgcggccgggt ctgctcgagc atgtccagcg cgtgcccgag 420
cgacagccgg ccgggtccgt tgagcaccga gttcagcgcc tcaccgttga ccagttccat 480
caccagatac gcggtgcggc cctcgccgct cagctggctc tcgcccagtg cgtgcacagc 540
cgcgatgccc ggggtggtga gcacgcgct ggtgcgcgcc tcggcgcgga accgctcgat 600
gaactcgggg tcttgggaga actcctgttt gagcaccttg accgcgagcg gccggccccg 660
ccggttgctg accgcctccc agacctgacc catgccgccg gtggcgatca ggcgctgcag 720
gcggtacctg ccagacagcg tcacaccaac tcgcgggctc at 762
```

What is claimed is:

1. A composition comprising an isolated mutant *M. paratuberculosis* bacterium, wherein the mutant bacterium comprises a disruption of function of a gene selected from the group consisting of *gcpE* (SEQ ID No:7), *pstA* (SEQ ID No:8), *kdpC* (SEQ ID No:9), *papA2* (SEQ ID No:10), *impA* (SEQ ID No:11), *umaA1* (SEQ ID No:12), *fabG2_2* (SEQ ID No:13), *aceAB* (SEQ ID No:14), *mbtH2* (SEQ ID No:15), *lppP* (SEQ ID No:16), *map0834c* (SEQ ID No:17), *cspB* (SEQ ID No:18), *lipN* (SEQ ID No:19), and *map1634* (SEQ ID No:20) genes of *M. paratuberculosis*.

2. The composition of claim 1 further comprising an adjuvant.

3. A composition comprising an isolated mutant *M. paratuberculosis* bacterium, wherein the mutant bacterium com-

prises a disruption of function of at least one gene from a genomic island selected from the group consisting of MAP-1 (SEQ ID No:21), MAP-2 (SEQ ID No:22), MAP-3 (SEQ ID No:23), MAP-4 (SEQ ID No:24), MAP-5 (SEQ ID No:25), MAP-6 (SEQ ID No:26), MAP-7 (SEQ ID No:27), MAP-8 (SEQ ID No:28), MAP-9 (SEQ ID No:29), MAP-10 (SEQ ID No:30), MAP-11 (SEQ ID No:31), MAP-12 (SEQ ID No:32), MAP-13 (SEQ ID No:33), MAP-14 (SEQ ID No:34), MAP-15 (SEQ ID No:35), MAP-16 (SEQ ID No:36), MAP-17 (SEQ ID No:37), and MAP-18 (SEQ ID No:38) genomic islands of *M. paratuberculosis*.

4. The composition of claim 3 further comprising an adjuvant.

5. The composition of claim 1 wherein the mutant *M. paratuberculosis* bacterium comprises a disruption of func-

87

tion of a gene selected from the group consisting of pstA (SEQ ID No: 8), papA2 (SEQ ID No:10), umaA1 (SEQ ID No:12), and fabG2_2 (SEQ ID No:13).

6. A composition comprising:

a) a eukaryotic expression vector comprising a nucleotide sequence encoding an antigen selected from the group comprising gcpE (SEQ ID No:7), pstA (SEQ ID No:8), kdpC (SEQ ID No:9), papA2 (SEQ ID No:10), impA (SEQ ID No:11), umaA1 (SEQ ID No:12), fabG2_2 (SEQ ID No:13), aceAB (SEQ ID No:14), mbtH2 (SEQ ID No:15), IpqP (SEQ ID No:16), map0834c (SEQ ID No:17), cspB (SEQ ID No:18), lipN (SEQ ID No:19), and map1634 (SEQ ID No:20) genes of *M. paratuberculosis*; and

b) a pharmaceutically acceptable carrier.

7. The composition of claim **6** further comprising an adjuvant.

88

8. A composition comprising:

a) a eukaryotic expression vector comprising a nucleotide sequence encoding an antigen selected from the group comprising MAP-1 (SEQ ID No:21), MAP-2 (SEQ ID No:22), MAP-3 (SEQ ID No:23), MAP-4 (SEQ ID No:24), MAP-5 (SEQ ID No:25), MAP-6 (SEQ ID No:26), MAP-7 (SEQ ID No:27), MAP-8 (SEQ ID No:28), MAP-9 (SEQ ID No:29), MAP-10 (SEQ ID No:30), MAP-11 (SEQ ID No:31), MAP-12 (SEQ ID No:32), MAP-13 (SEQ ID No:33), MAP-14 (SEQ ID No:34), MAP-15 (SEQ ID No:35), MAP-16 (SEQ ID No:36), MAP-17 (SEQ ID No:37), and MAP-18 (SEQ ID No:38) genomic islands of *M. paratuberculosis*; and

b) a pharmaceutically acceptable carrier.

9. The composition of claim **8** further comprising an adjuvant.

* * * * *