



US009254318B2

(12) **United States Patent**
Kawaoka et al.

(10) **Patent No.:** **US 9,254,318 B2**
(45) **Date of Patent:** **Feb. 9, 2016**

(54) **HIGH TITER RECOMBINANT INFLUENZA VIRUSES FOR VACCINES**

(75) Inventors: **Yoshihiro Kawaoka**, Middleton, WI (US); **Taisuke Horimoto**, Madison, WI (US); **Shin Murakami**, Tokyo (JP)

(73) Assignee: **Wisconsin Alumni Research Foundation**, Madison, WI (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/729,557**

(22) Filed: **Mar. 29, 2007**

(65) **Prior Publication Data**

US 2007/0231348 A1 Oct. 4, 2007

(51) **Int. Cl.**

A61K 39/12 (2006.01)

A61K 39/145 (2006.01)

C12N 7/00 (2006.01)

C12N 15/85 (2006.01)

A61K 39/00 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 39/145** (2013.01); **A61K 39/12** (2013.01); **C12N 7/00** (2013.01); **C12N 15/85** (2013.01); **A61K 2039/525** (2013.01); **A61K 2039/5254** (2013.01); **C12N 2760/16051** (2013.01); **C12N 2760/16134** (2013.01); **C12N 2760/16151** (2013.01); **C12N 2800/107** (2013.01)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,544,785	B1	4/2003	Palese et al.	
7,037,707	B2 *	5/2006	Webster et al.	435/235.1
7,226,774	B2	6/2007	Kawaoka	
7,507,411	B2 *	3/2009	Zhou et al.	424/186.1
8,012,736	B2	9/2011	Hoffman et al.	
8,309,099	B2	11/2012	Hoffmann	
8,475,806	B2	7/2013	Kawaoka	
2002/0164770	A1	11/2002	Hoffmann	
2003/0035814	A1	2/2003	Kawaoka et al.	
2007/0231348	A1 *	10/2007	Kawaoka et al.	424/209.1
2012/0020997	A1	1/2012	Hoffman et al.	

FOREIGN PATENT DOCUMENTS

AU	2012204138	B2	5/2014
EP	1201760	A1	5/2002
EP	2010557	B1	2/2014
JP	2004-500842	A	1/2004
JP	2005-523698	A	8/2005
JP	2005-245302	A	9/2005
JP	2005-535288	A	11/2005
JP	2014131516	A	7/2014
WO	WO-00/60050	A2	10/2000
WO	WO-01/83794	A2	11/2001

WO	WO-03/068923	A2	8/2003
WO	WO-03/091401	A2	11/2003
WO	WO-2004/112831	A2	12/2004
WO	WO-2004/112831	A2	7/2005
WO	WO-2005/062820	A2	7/2005
WO	WO-2007/126810	A2	11/2007

OTHER PUBLICATIONS

Voeten et al., Characterization of high-growth reassortant influenza A virus generated in MDCK cells cultured in serum-free medium, 1999, Vaccine, vol. 17, pp. 1942-1950.*

Kiseleva et al., Role of individual genes of the A/Leningrad/134/17/57 (H2N2) cold-adapted donor strain in manifestation of the temperature-sensitive phenotype of reassortant influenza A viruses, 2004, International Congress Series, vol. 1263, pp. 547-550.*

Lee et al., Generation of reassortant influenza vaccines by reverse genetics that allows utilization of a DIVA (Differentiating Infected from Vaccinated Animals) strategy for the control of avian influenza, 2004, Vaccine, vol. 22, pp. 3175-3181.*

Muster et al., An influenza A virus containing influenza B virus 5' and 3' noncoding regions on the neuraminidase gene is attenuated in mice, 1991, PNAS, vol. 88, pp. 5177-7181.*

Verma et al., Gene therapy—promises, problems and prospects, 1997, Nature, vol. 389, pp. 239-242.*

Orkin et al., Report and Recommendations of the Panel to Assess the NIH Investment in Research on Gene Therapy, 1995, downloaded on Sep. 5, 2008 from <<http://www.nih.gov/news/panelrep.html>>.*

Matsuoka et al., Neuraminidase Stalk Length and Additional Glycosylation of the Hemagglutinin Influence the Virulence of Influenza H5N1 Viruses for Mice, 2009, Journal of Virology, vol. 83, No. 9, pp. 4704-4708.*

Lazarovits et al., Endocytosis of Chimeric Influenza Virus Hemagglutinin Proteins That Lack a Cytoplasmic Recognition Feature for Coated Pits, 1996, The Journal of Cell Biology, vol. 134, No. 2, pp. 339-348.*

Yannarell et al., Factors affecting the yield of cold-adapted influenza virus vaccine, 1997, Journal of Virological Methods, vol. 64, pp. 161-169.*

Results 1, NCBI Blast nucleotide search of SEQ ID No. 7, database "nr"; Result 1, NCBI Blast nucleotide search of SEQ ID No. 8, database "Jul. 23, 2006", (Jul. 23, 2006), 8 pgs.

Result 17, NCBI Blast nucleotide search of SEQ ID No. 2, database "nr", (Jul. 18, 2006), 3 pgs.

Result 7, NCBI Blast nucleotide search of SEQ ID No. 1, database "nr", (Jul. 18, 2006), 3 pgs.

Result 1, NCBI Blast nucleotide search of SEQ ID No. 3, database "nr"; Result 4, NCBI Blast nucleotide search of SEQ ID No. 4, database "nr", (Jul. 22, 2006), 11 pgs.

(Continued)

Primary Examiner — Benjamin P Blumel

(74) *Attorney, Agent, or Firm* — Schwegman Lundberg & Woessner, P.A.

(57) **ABSTRACT**

The invention provides a composition useful to prepare high titer influenza viruses, e.g., in the absence of helper virus, which includes at least five internal genes from an influenza virus isolate that replicates to high titers in embryonated chicken eggs or MDCK cells.

32 Claims, 17 Drawing Sheets

(56)

References Cited

OTHER PUBLICATIONS

Result 2, NCBI Blast nucleotide search of SEQ ID No. 5, database "nr"; Result 4, NCBI Blast nucleotide search of SEQ ID No. 6, database "nr", (Jul. 22, 2006), 6 pgs.

PCT Application No. PCT/US2004/016680, International Search Report mailed Feb. 2, 2005, 7 pgs.

"PCT Application No. PCT/US2007/007562, International Search Report mailed Jan. 14, 2008", 8 pgs.

"PCT Application No. PCT/US2007/007562, Written Opinion mailed Jan. 14, 2008", 9 pgs.

De, B. P., et al., "Reverse Genetics of Negative Strand RNA Viruses", *Indian Journal of Biochemistry & Biophysics*, 31, (1994), 367-375.

Dreher, T. W., et al., "Mutational Analysis of the Sequence and Structural Requirements in Brome Mosaic Virus RNA for Minus Strand Promoter Activity", *Journal of Molecular Biology*, 201(1), (1988), 31-40.

Elliott, R. M., et al., "Rescue of Infectious Bunyavirus Entirely From Cloned cDNA", *10th International Conference on Negative Strand Virus*, (Abstract No. 96), (1997), 1 pg.

Fodor, E., "Rescue of Influenza A Virus from Recombinant DNA", *Journal of Virology*, 73(11), (1999), 9679-9682

Hoffmann, E., et al., "Ambisense Approach for the Generation of Influenza A Virus: vRNA and mRNA Synthesis from One Template", *Virology*, 267, (2000), 310-317.

Hoffmann, E., et al., "A DNA Transfection System for Generation of Influenza A Virus from Eight Plasmids", *Proceedings of the National Academy of Sciences of USA, National Academy of Science*, 97(11), (2000), 6108-6113.

Hoffmann, E., et al., "Eight-plasmid System for Rapid Generation of Influenza Virus Vaccines", *Vaccine, Butterworth Scientific Guildford*, 20(25-56), (2002), 3165-3170.

Hoffmann, E., et al., "Rescue of Influenza B Virus from Eight Plasmids", *Proceedings of the National Academy of Sciences of USA, National Academy of Science*, 99(17), (2002), 11411-11416.

Holmes, E. C., et al., "Whole-Genome Analysis of Human Influenza A Virus Reveals Multiple Persistent Lineages and Reassortment Among Recent H3N2 Viruses", *PLoS Biology*, 3(9), (2005), 1579-1589.

Horimoto, T., et al., "Generation of Influenza A Viruses with Chimeric (Type A/B) Hemagglutinins", *Journal of Virology*, 77(14), (2003), 8031-8038.

Horimoto, T., et al., "The Development and Characterization of H5 Influenza Virus Vaccines Derived from a 2003 Human Isolate", *Vaccine*, 24(17), (2006), 3669-3676.

Katfinger, D., et al., "Attenuated Influenza Viruses as a Vector for Mucosal Immunization Against HIV-1", *Vaccines*, 97, Cold Spring Harbor, (1997), 315-319.

Lamb, R. A., et al., "Chapter 20—*Paramyxoviridae*: The Viruses and Their Replication", *In: Fundamental Virology*, Fields, B. N., et al., Editors, Lippincott-Raven (2nd Edition), (1996), 577-647.

Neumann, G., et al., "Generation of influenza A Virus from Cloned cDNAs—Historical Perspective and Outlook for the New Millennium", *Reviews in Medical Virology*, 12(1), (2002), 13-30.

Neumann, G., et al., "Generation of Influenza A Viruses Entirely from Cloned cDNAs", *Proc. Natl. Acad. Sci. USA*, 96(16), (1999), 9345-9350.

Neumann, G., et al., "Plasmid-Driven Formation of Influenza Virus-like Particles", *Journal of Virology, The American Society For Microbiology*, 74(1), (2000), 547-551.

Odagiri, T., et al., "Nucleotide Sequence of the PA Gene of Influenza A/WSN/33 (H1N1)", *Nucleic Acids Research*, 18(3), Department of Virology, (1990), p. 654.

Schickli, J. H., et al., "Plasmid-only Rescue of Influenza A Virus Vaccine Candidates", *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 356(1416), (2001), 1965-1973.

Smeenk, et al., "Mutations in the Hemagglutinin and Matrix Genes of a Virulent Influenza Virus Variant, A/FM/1/47-MA, Control Different Stages in Pathogenesis", *Virus Research* 44, (1996), 79-95.

Subbarao, K., et al., "Evaluation of a Genetically Modified Reassortant H5N1 Influenza A Virus Vaccine Candidate Generated by Plasmid-based Reverse Genetics", *Virology*, vol. 30(1), (2003), 192-200.

"European Application Serial No. 07754132.4, Office Action mailed Apr. 28, 2009", 4 pgs.

"Australian Application Serial No. 2007245192, Office Action mailed Aug. 25, 2011", 2 pgs.

"Chinese Application Serial No. 200780020095.1, First Office Action mailed Jun. 24, 2011", (w/ English Translation), 13 pgs.

"European Application Serial No. 07754132.4, Office Action mailed Sep. 5, 2011", 5 pgs.

"European Application Serial No. 07754132.4, Response filed Feb. 5, 2010 to Office Action mailed Apr. 28, 2009", 15 pgs.

"Australian Application Serial No. 2007245192, Response filed Feb. 28, 2012 to Office Action mailed Aug. 25, 2011", 22 pgs.

"Chinese Application Serial No. 200780020095.1, Office Action mailed May 3, 2012", With English Translation, 10 pgs.

"Chinese Application Serial No. 200780020095.1, Response filed Sep. 17, 2012 to Office Action mailed May 3, 2012", (w/ English Translation of Claims), 17 pgs.

"Chinese Application Serial No. 200780020095.1, Response filed Nov. 8, 2011 to Office Action mailed Jun. 24, 2011", (w/ English Translation of Amended Claims), 20 pgs.

"European Application Serial No. 07754132.4, Office Action mailed Nov. 2, 2012", 4 pgs.

"European Application Serial No. 07754132.4, Response filed Mar. 15, 2012 to Office Action mailed Sep. 5, 2011", 21 pgs.

"Japanese Application Serial No. 2009-502945, Office Action mailed Oct. 23, 2012", (w/ English Translation), 16 pgs.

Chen, H, et al., "Generation and evaluation of a high-growth reassortant H9N2 influenza A virus as a pandemic vaccine candidate", *Vaccine*, 21(17-18), (May 16, 2003), 1974-1979.

"Australian Application Serial No. 2012204138, First Examiner Report mailed Jul. 16, 2013", 4 pgs.

"Canadian Application Serial No. 2,647,985, Office Action mailed May 15, 2013", 3 pgs.

"Chinese Application Serial No. 200780020095.1, Decision on Rejection mailed Jul. 22, 2013", (w/ English Translation), 11 pgs.

"Chinese Application Serial No. 200780020095.1, Response filed Jun. 9, 2013 to Office Action mailed Jun. 29, 2013", (w/ English Translation of Claims), 10 pgs.

"Australian Application Serial No. 2012204138, Response filed Dec. 24, 2013 to First Examiner Report mailed Jul. 16, 2013", 21 pgs.

"Canadian Application Serial No. 2,647,985, Response filed Sep. 30, 2013 Office Action mailed May 15, 2013", 20 pgs.

"Chinese Application Serial No. 200780020095.1, Response filed Nov. 5, 2013 to Decision on Rejection mailed Jul. 22, 2013", (w/ English Translation of Claims), 12 pgs.

"European Application Serial No. 07754132.4, Response filed Jun. 26, 2013", 8 pgs.

"Japanese Application Serial No. 2009-502945, Examiners Decision of Final Refusal mailed Nov. 12, 2013", (w/ English Translation), 8 pgs.

"PCT Application Serial No. PCT/US2007/007562, International Preliminary Report on Patentability mailed Oct. 9, 2008", 5 pgs.

"Chinese Application Serial No. 200780020095.1, Office Action mailed Jan. 29, 2013", (w/ English Translation), 10 pgs.

"European Application Serial No. 07754132.4, Response filed May 10, 2013 to Office Action mailed Nov. 2, 2012", 14 pgs.

"Japanese Application Serial No. 2009-502945, Response filed Apr. 10, 2013 to Office Action mailed Oct. 23, 2012", (w/ English Translation of Claims), 18 pgs.

"Chinese Application Serial No. 200780020095.1, Office Action mailed Mar. 5, 2015", (w/ English Translation), 12 pgs.

"Chinese Application Serial No. 200780020095.1, Response filed Jun. 23, 2015 to Office Action mailed Mar. 5, 2015", (w/ English Translation of Claims), 16 pgs.

"Japanese Application Serial No. 2014-049025, Office Action mailed Jun. 16, 2015", (w/ English Translation), 6 pgs.

Baez, M., et al., "Complete nucleotide sequence of the influenza A/PR/8/34 virus NS gene and comparison with the NS genes of the

(56)



References Cited


OTHER PUBLICATIONS


A/Udorn/72 and A/FPV/Rostock/34 strains", *Nucleic Acids Research*, 23(8), (1980), 5845-5858.


Winter, G., et al., "The use of synthetic oligodeoxynucleotide primers in cloning and sequencing segment 8 of influenza virus (A/PR/8/34)", *Nucleic Acids Research*, 9(2), (1981), 237-245.

* cited by examiner



*FIG. 2*

*FIG. 3*

*FIG. 4*





FIG. 5

*FIG. 6*

GROWTH OF REASSORTANT H5N1 VIRUSES POSSESSING PR8(UW)
OR PR8(CAMBRIDGE) INTERNAL GENES IN CHICKEN EMBRYONATED EGGS

	PA	PB1	PB2	HA	NP	NA	M	NS	LOG ₁₀ EID ₅₀ /ml
PR8(UW)/1194	R	R	R	G	R	G	R	R	9.07 ± 0.37
PR8(UW)/1194-CamPA	B	R	R	G	R	G	R	R	8.88 ± 0.25
PR8(UW)/1194-CamPB1	R	B	R	G	R	G	R	R	9.08 ± 0.38
PR8(UW)/1194-CamPB2	R	R	B	G	R	G	R	R	9.05 ± 0.40
PR8(UW)/1194-CamNP	R	R	R	G	B	G	R	R	9.00 ± 0.20
PR8(UW)/1194-CamPB12	R	B	B	G	R	G	R	R	8.75 ± 0.25
PR8(UW)/1194-CamP3	B	B	B	G	R	G	R	R	8.56 ± 0.13*
PR8(UW)/1194-CamP3NP	B	B	B	G	B	G	R	R	8.19 ± 0.31*
NIBRG-14	B	B	B	G	B	G	B	B	8.32 ± 0.20*

R	B	G
PR8(UW)	PR8(CAMBRIDGE)	A/VIETNAM/1194/2004

FIG. 7

THE EFFECT OF THE M AND NS GENES ON THE GROWTH OF
VIRUSES IN CHICKEN EMBRYONATED EGGS

	PA	PB1	PB2	HA	NP	NA	M	NS	$\times 10^8$ PFU/ml
PR8(UW)/1194	R	R	R	G	R	G	R	R	3.39 ± 1.42
PR8(UW)/1194-CamM	R	R	R	G	R	G	B	R	2.94 ± 0.28
PR8(UW)/1194-CamNS	R	R	R	G	R	G	R	B	$6.25 \pm 1.44^*$
NIBRG-14	B	B	B	G	B	G	B	B	0.73 ± 0.43
	R			B			G		
	PR8(UW)			PR8(CAMBRIDGE)			A/VIETNAM/1194/2004		

FIG. 8

Growth of PR8(UW)/1194 and NIBRG-14 virus in MDCK cells




FIG. 9

Identification of a gene segment responsible for the better growth of PR8(UW)/1194 relative to NIBRG-14 in MDCK cells




FIG. 10

Identification of the amino acid in PB2 responsible for the high growth rate of the vaccine seed virus in MDCK cells





FIG. 11

GROWTH RATES IN MDCK CELL OF REASSORTANTS WITH DIFFERENT HA, NA, AND NS GENES

	PA	PB1	PB2	HA	NP	NA	M	NS	x10 ⁸ PFU/ml
PR8(UW)/1194	R	R	R	G1	R	G1	R	R	0.39 ± 0.21
PR8(UW)/1203	R	R	R	G2	R	G2	R	R	0.40 ± 0.20
PR8(UW)/1203/03FILL	R	R	R	G2	R	Y	R	R	1.26 ± 0.47*
PR8(UW)/1203/HK213	R	R	R	G2	R	BR	R	R	1.10 ± 0.37*
PR8(UW)/1203/03FILL-CamNS	R	R	R	G2	R	Y	R	B	2.33 ± 0.27**
PR8(UW)/1203/HK213-CamNS	R	R	R	G2	R	BR	R	B	2.35 ± 0.21**
	R			B			33°C, 36 hpi, moi=0.01		
	PR8(UW)			PR8(CAMBRIDGE)					
	G1			G2					
	A/VIETNAM/1194/2004			A/VIETNAM/1203/2004 (1203)					
	Y			BR					
	VN1203FILL (1203 WITH LONGER STALK)			A/HONG KONG/213/2003					

FIG. 12

Growth in MDCK cells of the H5N1 vaccine seed virus containing an heterologous NS segment



THE GENOTYPE OF AN H5N1 VACCINE SEED VIRUS WITH
HIGH GROWTH CAPACITY IN CHICKEN EMBRYONATED EGGS OR MDCK CELLS




FIG. 14

PR8(Cambridge)

PB2

AGCGAAAGCAGGTCAATTATATTCATATGGAAGAATAAAGAACTAAGAAATCTAATGTCGCGAGTCTCGCACCCGCGAGATA
CTCACAAAACACCCTGGACCATATGGCCATATCAAGAAGTACACATCAGGAAGACAGGAGAAGAACCCAGCACTTAGGATG
AAATGGATGATGGCAATGAAATATCCAATTACAGCAGACAAGAGGATAACCGAAATGATTCTGAGAGAAATGAGCAAGGACAA
ACTTTATGGAGTAAATGAATGATGCCGATCAGACCGAGTGATGGTATCACCTCTGGCTGTGACATGGTGGAATAGGAATGGA
CCAATGACAAATACAGTTCAATTATCCAAAACTACAAAACTTATTTTGAAGAGTGAAGAGGCTAAAGCATGGAACCTTTGGC
CCTGTCCATTTTAGAAACCAAGTCAAAATACGTCCGAGAGTTGACATAAATCCTGGTCAATGCAGATCTCAGTGCCAAAGGAGGCA
CAGGATGTAATCATGGAAGTTGTTTTCCCTAACGAAGTGGGAGCCAGGATACTAACATCGGAATCGCAACTAACGATAACCAAA
GAGAAGAAAGAAAGAACTCCAGGATTGCAAAATTTCTCCTTTGATGGTTGCATACATGTTGGAGAGAGAACTGGTCCGCAAAACG
AGATTCTCCAGTGGCTGGTGGAACAAGCAGTGTGTACATTGAAGTGTTCATTTGACTCAAGGAACATGCTGGGAACAGATG
TATACTCCAGGAGGGGAAGTGAAGAATGATGATGTTGATCAAGCTTGATTATTGCTGCTAGGAACATAGTGAGAAGAGCTGCA
GTATCAGCAGACCCACTAGCATCTTTATTTGGAGATGTGCCACAGCACAGATTGGTGGAAATTAGGATGGTAGACATCCTTAAG
CAGAACCCAAACAGAAGAGCAAGCCGTGGATATATGCAAGGCTGCAATGGGACTGAGAATTAGCTCATCCTTCAGTTTTGGTGGA
TTCACATTTAAGAGAACAAGCGGATCATCAGTCAAGAGAGAGGAAGAGGTGCTTACGGGCAATCTTCAACATTGAAGATAAGA
GTGCATGGGGATCTGAAGAGTTCACAATGTTGGGAGAAGAGCAACAGCCATCTCAGAAAAGCAACCCAGGAGATTGATTCAG
CTGATAGTGAGTGGGAGAGACGAACAGTCGATTGCCGAAGCAATAATTGTGGCCATGGTATTTTCAAGAGAGGATTGTATGATA
AAAGCAGTTAGAGGTGATCTGAATTTCTGCAATAGGGCGAATCAGCGACTGAATCCTATGCATCAACTTTTAAGACATTTTCAG
AAGGATGCCAAAGTGCTTTTTCAAAATTTGGGGAGTTGAACCTATCGACAATGTGATGGGAATGATTGGGATTTGCCGACATG
ACTCCAAGCATCGAGATGTCAATGAGAGGAGTGAGAATCAGCAAAATGGGTGTAGATGAGTACTCCAGCAGGAGAGGGTATGTG
GTGAGCATTGACCGGTTCTTGAGAGTCAAGGACCAACGAGGAATGTACTACTGTCTCCGAGGAGGTGAGTGAACACAGGGA
ACAGAGAACTGACAATAACTTACTCATCTGTCATGATGTGGGAGATTAAATGGTCTGAATCAGTGTGGTCAATACCTATCAA
TGGATCATCAGAACTGGGAACTGTTAAATTCAGTGGTCCAGAACCTACAATGCTATACAATAAAATGGAATTTGAACCA
TTTCAGTCTTTAGTACCTAAGGCCATTAGAGGCCAATACAGTGGGTTGTGAAGAACTCTGTTCCAACAAATGAGGGATGTGCTT
GGGACATTTGATACCGCACAGATAATAAACTTCTTCCCTTCGACGCCCTCCACCAAGCAAGTAGAATGCGAGTTCTCCTCA
TTTACTGTGAATGTGAGGGGATCAGGAATGAGAATACTTGTGAAGGGGCAATTCTCCTGTATTCACTACAACAAGGCCACGAAG
AGACTCACAGTTCTCGGAAAGGATGCTGGCATTAAACCGAAGACCCAGATGAAGGCACAGCTGGAGTGGAGTCCGCTGTTCTG
AGGGGATTCCTCATTCTGGGCAAGAAAGACAGGAGATATGGGCCAGCTTAAGCATCAATGAACTGAGCAACCTTGGCAAGGA
GAGAAGGCTAATGTGCTAATTGGGCAAGGAGCTGTGTTGGTGAATGAAACGAAACGGGACTCTAGCATACTTACTGACAGC
CAGACAGCGACCAAAAGAAATTCGGATGGCCATCAATTAGTGTGCAATAGTTTAAAAACGACCTTGTCTCTACT

SEQ ID NO. 33

PB1

AGCGAAAGCAGGCAAAACCATTTGAATGGATGTCAATCCGACCTTACTTTTCTTAAAAAGTGCCAGCACAAAATGCTATAAGCACA
ACTTTCCCTTATACCGGAGACCCTCCTTACAGCCATGGGACAGGAACAGGATACACCATGGATACTGTCAACAGGACACATCAG
TACTCAGAAAAGGGAAGATGGACAACAAACACCGAACTGGAGCACCGCAACTCAACCCGATTGATGGGCCACTGCCAGAAGAC
AATGAACCAAGTGTTTATGCCCAACAGATTGTGATTGGAAGCAATGGCTTTCTTGGGAATCCCATCCTGGTATTTTGGAA
AACTCGTGTATTGAAACGATGGAGGTTGTTGAGCAAAACAGAGTAGACAAGCTGACACAAGGCCGACAGACCTATGACTGGACT
TTAAATAGAAACAGCCTGTGCAACAGCATTGGCCAAACACATAGAAGTGTTCAGATCAAAATGGCCTCACGGCCAATGAGTCA
GGAAGGCTCATAGACTTCTTAAAGGATGTAATGGAGTCAATGAAAAAGAAAGAAATGGGGATCACAACCTATTTTCAGAGAAAG
AGACGGGTGAGAGACAATATGACTAAGAAAATGATAACACAGAGAACAATAGGTAAAAGGAAACAGAGATTGAACAAAAGGGGT
TATCTAATTAGAGCATTGACCTGAACACAATGACCAAGATGCTGAGAGAGGGGAAGCTAAAACGGAGAGCAATTGCAACCCCA
GGGATGCAAAATAGGGGGTTTGTATACTTTGTTGAGACACTGGCAAGGAGTATATGTGAGAACTTGAACAATCAGGGTTGCCA
GTTGGAGGCAATGAGAAGAAAGCAAGTTGGCAATGTTGTGAAGGAAGATGATACCAATTCTCAGGACACCGAATTTCTTTTC
ACCATCACTGGAGATAACACCAAAATGGAACGAAAATCAGAATCCTCGGATGTTTTTGGCCATGATCACATATATGACCAGAAAT
CAGCCCGAATGGTTCAAGAAATGTTCTAAGTATTGCTCCAATAATGTTCTCAACAAAATGGCGAGACTGGGAAAAGGGTATATG
TTTGAGAGCAAGAGTATGAACTTAGAACTCAAAATACCTGCAGAAATGCTAGCAAGCATTGATTTGAAATATTTCAATGATTCA
ACAAGAAAGAAAGATTGAAAAATCCGACCGCTCTTAATAGAGGGGACTGCATCATTGAGCCCTGGAATGATGATGGGCATGTTT
AATATGTTAAGCACTGTATTAGGCGTCTCCATCCTGAATCTTGGACAAAAGAGATACCCAAGACTACTTACTGGTGGGATGGT
CTTCAATCCTCTGACGATTTTGTCTGATTGTGAATGCACCAATCATGAAGGGATTCAAGCCGGAGTCGACAGGTTTATTCGA
ACCTGTAAGCTACTTGAATCAATATGAGCAAGAAAAAGTCTTACATAAACAGAACAGGTACATTTGAATTCACAAGTTTTTTC
TATCGTTATGGGTTTGTGCCAATTTAGCATGGAGCTTCCAGTTTTGGGGTGTCTGGGATCAACGAGTCAGCGGACATGAGT
ATTGGAGTTACTGTCTCAAAAACAATATGATAACAATGATCTTGGTCCAGCAACAGCTCAAATGGCCCTTCAGTTGTTTATC
AAAGATTACAGGTACACGTACCGATGCCATAGAGGTGACACACAATAACAACCCGAAGATCATTGAAATAAAGAACTGTGG
GAGCAAAACCGTTCCAAAGCTGGACTGCTGGTCTCCGACGGAGGCCAAATTTATACAACATTAGAAATCTCCACATTCCTGAA
GTCTGCGCTAAATGGGAATTTGATGGATGAGGATTACAGGGGCGTTTATGCAACCCCACTGAACCCATTGTGCAACAAAGAA
ATTGAATCAATGAACAAATGAGTGATGATGCCAGCATGGTCCAGCCAAAACATGGAGTATGATGCTGTTGCAACAACACAC
TCCTGGATCCCCAAAGAAATCGATCCATCTTGAATACAAGTCAAGAGGAGTACTTGAAGTGAACAAATGTACCAAGAGGTTC
TGCAATTTATTTGAAAAATCTTCCCAGCAGTTTATACAGAAGACAGTCGGGATATCCAGTATGGTGGAGGCTATGGTTCC
AGAGCCCGAATTTGTCACGGATTGTTTTGCAATCTGGAAGGATAAGAAAGAAAGATTCACTGAGATCATGAAGATCTGTTCC
ACCATTGAAGAGCTCAGACGGCAAAAATAGTGAATTTAGCTGTCTTCATGAAAAATGCCCTGTTTCTACT

SEQ ID NO. 34

FIG. 15A

PR8(Cambridge)

PA

AGCGAAAGCAGGTAAGTATCAAAATGGAAGATTTTGTGCGACAATGCTTCAATCCGATGATTGTCGAGCTTGGCGAAAAACA
ATGAAAGAGTATGGGGAGGACCTGAAATCGAAACAAACAAATTTGCGACCAATATGCACTCACTTGGAAAGTATGCTTCATGTAT
TCAGATTTCCACTTCATCAATGAGCAAGGCGAGTCAATATCGTAGAATCTTGGTATCCTAATGCACCTTTGAAGCACAGATTT
GAAATAATCGAGGGAAGAGATCGCAATGGCCTGGACAGTAGTAACAGTATTTGCAACACTACAGGGGCTGAGAAACCAAG
TTTCTACCAGATTTGTATGATTACAAGGAAAATAGATTCATCGAAATTTGGAGTAACAAGGAGAGAAGTTCACATATACTATCTG
GAAAAGGCCAATAAAATTAATCTGAGAAAAACACATCCACATTTTCTCGTTCACTGGGGAAGAAATGGCCACAAGGGCCGAC
TACACTCTCGATGAAGAAAGCAGGGCTAGGATCAAAACAGGCTATTCAACATAAGACAAGAAATGGCCAGCAGAGGCCTCTGG
GATTCTTTCTGTCAGTCCGAGAGAGGAGAGAGACAATTGAAGAAAGGTTTGAATCACAGGAACAATGGCAAGCTTGCCGAC
CAAAGTCTCCCGCGAATCTTCCAGCCTTGAAAATTTTAGAGCCTATGTGGATGGATTGCAACCGAAGCGCTACATTGAGGGC
AAGCTGTCTCAATGTCCAAGAAAGTAAATGCTAGAAATGAACCTTTTGAAGAAACACACCAGACCTTAGACTTCCGAAT
GGGCTCTCTGTTCTCAGCGGTCCAAATTTCTGCTGATGGATGCTTAAATTAAGCAATTGAGGACCAAGTCAAGAGGAGAG
GGAATACCGCTATATGATGCAATCAATGCAATGAGAACATTTCTTGGATGGAAGGAACCAATGTTGTTAAACCAACAGAAAAG
GGAATAAATCAAAATTATCTTCTGTCTGGAAGCAAGTACTGGCAGAACTGCAGGACATTGAGAATGAGGAGAAAATTTCAAAG
ACTAAAATATGAAAAAACAAAGTCAAGTAAAGTGGGCACTTGGTGAGAACATGGCACCAGAAAAGGTAGACTTTGACGACTGT
AAAGATGTAGGTGATTTGAAGCAATATGATAGTATGATGAACCAAGATTTGAGGTGCTTGAAGTTGGATTGAGATGAGGCAAC
AAGGCATGCGAACTGACAGATTCAAGCTGGATAGAGCTTGTATGAGATTGGAGAAGATGTGGCTCCAATTGAACACATTGCAAGC
ATGAGAAGGAATTTATTTACATCAGAGGTGTCTCACTGAGAGCCACAGAAATACATAATGAAGGGGGTGTACATCAATCTGCC
TTACTTAATGCATCTTGTGCGCAATGGATGATTTCAAATTAATTTCCAATGATAAGCAAGTGTAGAATGAAGGAGGAAGGCGA
AAGACCAACTTGTATGTTTCTCATATAAAGGAAGATCCCACTTAAGGAATGACACCGAGCTGGTAACTTTGTGAGCATGGAG
TTTTCTCTCACTGACCCAGACTTGAACCAACAAATGGGAGAAAGTACTGTGTTCTTGAGATAGGAGATATGCTTCTAAGAAAT
GCCATAGGCCAGGTTTCAAGGCCCATGTTCTTGTATGTGAGGACAAATGGAACCTCAAAAATTAATAATGAATGGGGAATGGAG
ATGAGGCGTTGTCTCTCCAGTCACTTCAACAAATTTGAGAGTATGATTGAAGCTGAGTCTCTGTCAAAGAGAAAGACATGACC
AAAGAGTCTTTGAGAACAAATCAGAAACATGGCCCATTTGGAGAGTCTCCCAAGGAGTGGAGGAAAGTTCATTGGGAAGGTC
TGCAGGACTTTATAGCAAAGTCGGTATTTAACAGCTTGTATGCATCTCAACACTAGAAGGATTTTCACTGAATCAAGAAAA
CTGCTTCTTATCGTTCAAGGCTCTTAGGACAATCTGGAACCTGGGACCTTTGATCTTGGGGGCTATATGAAGCAATGAGGAG
TGCCTAATTAATGATCCCTGGGTTTGTCTTAATGCTTCTTGGTTCAACTCTTCTTACACATGCATTGAGTTAGTTGTGGCAG
TGCTACTATTTGCTATCATCTGTCCAAAAAGTACCTTGTCTACT

SEQ ID NO. 35

NP

AGCAAAAGCAGGTAGATAATCACTCACTGAGTGACATCAAAATCATGGCGTCCCAAGGCACCAACCGGCTTACGAACAGATG
GAGACTGATGGAGAAGCCAGAAATGCCACTGAAATCAGAGCATCCGTCGGAAAAATGATTGGTGGAAATGGACGATTCTACATC
CAAATGTGCACAGAACTTAAACTCAGTGATTATGAGGGACGGTTGATCCAAACAGCTTAACAATAGAGAGAATGGTGCTCTCT
GCTTTTGACGAAAGGAGAAATAAATACCTGGAAGAACATCCCACTGCGGGGAAAGATCCTAAGAAAACTGGAGGACCTATATAC
AGAAGAGTAAACGGAAAGTGGATGAGAGAACTCATCTTTATGACAAAGAAAGAAATGAAGCGAATCTGGCGCCAAAGCTAATAAT
GGTGACGATGCAACGGCTGGTCTGACTCACATGATGATCTGGCATTCCAATTTGAATGATGCAACTTATCAGAGGACAAGGGCT
CTTGTTCGACCCGGAATGGATCCAGGATGTGCTCTCTGATGCAAGGTTCAACTCTCCCTAGGAGGTCTGGAGCCGAGGTGCT
GCAGTCAAAGGAGTTGGAACAAATGGTGTGGAATTTGGTCAGGATGATCAACCGTGGGATCAATGATCGGAACCTTCTGGAGGGT
GAGAATGGACGAAAAACAAGAAATGCTTATGAAAGAAATGTGCAACTTCAAAAGGAAATTTCAAATGCTGCACAAAAAGCA
ATGATGGATCAAGTGAGAGAGAGCCGGAACCCAGGGAATGCTGAGTTGGAAGATCTCACTTTCTAGCACGGTCTGCACTCATA
TTGAGAGGGTCTGTTGCTCACAAGTCTGCTGCTGCTGCTGTTGATGGACCTGCGGTAGCCAGTGGGTACGACTTTGAAAGA
GAGGGATACTCTAGTCGGAATAGACCTTTCAAGTCTGCTTCAAAACAGCCAAAGTGTACAGCCTAATCAGACCAATGAGAAT
CCAGCACACAAGAGTCACTGGTGTGGATGGCATGCCATTCTGCCGCAATTTGAAGATCTAAGAGTATTGAGCTTCATCAAGGG
ACGAAGTGGTCCCAAGAGGGAAGCTTTCCAATAGAGGAGTTCAAATGCTTCCAATGAAATATGGAGACTATGGAATCAAGT
ACACTTGAAGTGAAGAGCAGGTAAGGACCAAGGAGGAGGAAACCAATCAACAGAGGGGATCTGCGGGCCAA
ATCAGCATACAACCTACGTTCTCAGTACAGAGAAATCTCCCTTTGACAGAACCAACCGTTATGGCAGCATTTCACTGGGAATACA
GAGGGGAGAACATCTGACATGAGGACCGAAATCATAAGGATGATGGAAAGTGAAGACCAAGATGTGTCTTCCAGGGGCGG
GGAGTCTTCGAGCTCTCGACGAAAAAGCAGCGAGCCGATCGTGCTTCTTGTGATGAGTAATGAAGGATCTTATTTCTTC
GGAGACAATGCAGAGGAGTACGACAATTAAAGAAAAATACCTTGTCTTACT

SEQ ID NO. 36

M

AGCAAAAGCAGGTAGATATTGAAGATGAGTCTTCTAACCGAGGTGCAAAACGTACGTTCTCTATCATCCCGTCAGGCCCTCT
CAAAGCCGAGATCGCACAGAGACTTGAAGATGCTTTGAGGGAAGAAACACCGATCTTGAAGTTCTCATGGAATGGCTAAAGAC
AAGACCAATCTGTCACTCTGACTAAGGGGATTTAGGATTTGTGTTACGCTCACCGTGCCAGTGAGCGAGGACTGCAACG
TAGACGCTTTGTCCAAATGCCCTTAATGGGAACGGGGATCCAAATAACATGGACAAAGCAGTTAACTGTATAGGAAGCTCAA
GAGGGAGATAACATTCATGGGGCCAAAGAAATCTCACTCAATTATTCTGCTGGTGCATTTGCCAGTTGTATGGGCTCATATA
CAACAGGATGGGGCTGTGACCACTGAAGTGGCATTGTGCTGATGTGCAACCTGTGAACAGATTGCTGACTCCAGCATCG

FIG. 15B

PR8 (Cambridge)

GTCTCATAGGCAATGGTGACAACAACCAACCCACTAATCAGACATGAGAACAGAATGGTTTTAGCCAGCACTACAGCTAAGGC
TATGGAGCAATGGCTGGATCGAGTGAGCAAGCAGCAGAGGCCATGGAGGTTGCTAGTCAGGCTAGGCAATGGTGCAAGCGAT
GAGAACCATTGGGACTCATCCTAGCTCCAGTGCTGGTCTGAAAAATGATCTTCTTGAAAAATTTGCAGGCTATCAGAAACGAAT
GGGGGTGCAGATGCAACGGTTCAAGTGATCCTCTCGCTATTGCCGCAAAATCATTGGGATCTTGCACTTGATATTGTGGATTCT
TTGATCGTCTTTTTTCAAATGCATTTACCGTCGCTTTAATAACGGAAGTAAAGGAGGGCCTTCTACGGAAGGAGTGCCAAAGT
CTATGAGGGAAGAATATCGAAAGGAACAGCAGAGTGCTGTGGATGCTGACGATGGTCATTTTGTGAGCATAGAGCTGGAGTAA
AAACTACCTTGTTTCTACT

SEQ ID NO. 37

NS

AGCAAAAGCAGGGTGACAAAGACATAATGGATCCAAACACTGTGTCAAGCTTTCAGGTAGATTGCTTTCTTTGGCATGTCCGCA
AACGAGTTGCAGACCAAGAAGTGGTGATGCCCCATTCTTGATCGGCTTCGCCGAGATCAGAAATCCCTAAGAGGAAGGGGCA
GCACTCTTGGTCTGGACATCGAGACAGCCACACGTGCTGGAAAGCAGATAGTGGAGCGGATTCTGAAAGAAGAATCCGATGAGG
CACTTAAATGACCATGGCCTCTGTACCTGCGTCGCGTTACCTAACCGACATGACTCTTGAGGAAATGTCAAGGGAATGGTCCA
TGCTCATACCAAGCAGAAAGTGGCAGGCCCTCTTTGTATCAGAATGGACCAGGCGATCATGGATAAAAAACATCATACTGAAAG
CGAATTTCAGTGTGATTTTGGACCGGCTGGAGACTCTAATATTGCTAAGGGCTTTCACCGAAGAGGGAGCAATTGTTGGCGAAA
TTTCACCATTTGCCCTTCTCTCCAGGACATACTGCTGAGGATGTCAAAAATGCAGTTGGAGTCCTCATCGGAGGACTTGAATGGA
ATGATAACACAGTTCGAGTCTCTGAAACTCTACAGAGATTGCTTTGGAGAAGCAGTAATGAGAATGGGAGACCTCCACTCACTC
CAAAACAGAAACGAGAAATGGCGGGAACAATTAGGTCAGAAGTTTGAAGAAATAAGATGGTTGATTGAAGAAGTGAGACACAAA
CTGAAGGTAACAGAGAATAGTTTTGAGCAATAACATTTATGCAAGCCTTACATCTATTGCTTGAAGTGGAGCAAGAGATAAGA
ACTTTCTCATTTGAGCTTATTTAATAATAAAAAACACCCTTGTTTCTACT

SEQ ID NO. 38

FIG. 15C

1

HIGH TITER RECOMBINANT INFLUENZA VIRUSES FOR VACCINES

GOVERNMENT SUPPORT

This invention was made with government support under Grant Number AI044386 from the National Institutes of Health. The United States Government has certain rights in the invention.

BACKGROUND

Negative-sense RNA viruses are classified into seven families (*Rhabdoviridae*, *Paramyxoviridae*, *Filoviridae*, *Bornaviridae*, *Orthomyxoviridae*, *Bunyaviridae*, and *Arenaviridae*) which include common human pathogens, such as respiratory syncytial virus, influenza virus, measles virus, and Ebola virus, as well as animal viruses with major economic impact on the poultry and cattle industries (e.g., Newcastle disease virus and Rinderpest virus). The first four families are characterized by nonsegmented genomes, while the latter three have genomes comprised of six-to-eight, three, or two negative-sense RNA segments, respectively. The common feature of negative-sense RNA viruses is the negative polarity of their RNA genome; i.e., the viral RNA (vRNA) is complementary to mRNA and therefore is not infectious by itself. In order to initiate viral transcription and replication, the vRNA has to be transcribed into a plus-sense mRNA or cRNA, respectively, by the viral polymerase complex and the nucleoprotein; for influenza A viruses, the viral polymerase complex is comprised of the three polymerase proteins PB2, PB1, and PA. During viral replication, cRNA serves as a template for the synthesis of new vRNA molecules. For all negative-stranded RNA viruses, non-coding regions at both the 5' and 3' termini of the vRNA and cRNA are critical for transcription and replication of the viral genome. Unlike cellular or viral mRNA transcripts, both cRNA and vRNA are neither capped at the 5' end nor polyadenylated at the very 3' end.

The basic functions of many viral proteins have been elucidated biochemically and/or in the context of viral infection. However, reverse genetics systems have dramatically increased our knowledge of negative-stranded segmented and non-segmented RNA viruses with respect to their viral replication and pathogenicity, as well as to the development of live attenuated virus vaccines. Reverse genetics, as the term is used in molecular virology, is defined as the generation of virus possessing a genome derived from cloned cDNAs (for a review, see Neumann et al., 2002).

In order to initiate viral replication of negative-stranded RNA viruses, vRNA(s) or cRNA(s) must be coexpressed with the polymerase complex and the nucleoprotein. Rabies virus was the first non-segmented negative-sense RNA virus which was generated entirely from cloned cDNA: Schnell et al. (1994) generated recombinant rabies virus by cotransfection of a cDNA construct encoding the full-length cRNA and protein expression constructs for the L, P, and N proteins, all under control of the T7 RNA polymerase promoter. Infection with recombinant vaccinia virus, which provided T7 RNA polymerase, resulted in the generation of infectious rabies virus. In this T7 polymerase system, the primary transcription of the full length cRNA under control of the T7 RNA polymerase resulted in a non-capped cRNA transcript. However, three guanine nucleotides, which form the optimal initiation sequence for T7 RNA polymerase, were attached to the 5' end. In order to create an authentic 3' end of the cRNA transcript which is essential for a productive infective cycle,

2

the hepatitis delta ribozyme (HDVRz) sequence was used for exact autocatalytic cleavage at the 3' end of the cRNA transcript.

Since the initial report by Schnell et al. (1994), reverse genetics systems using similar techniques led to the generation of many non-segmented negative strand RNA viruses (Conzelmann, 1996; Conzelmann, 1998; Conzelmann et al., 1996; Marriott et al., 1999; Munoz et al., 2000; Nagai, 1999; Neumann et al., 2002; Roberts et al., 1998; Rose, 1996). Refinements of the original rescue procedure included the expression of T7 RNA polymerase from stably transfected cell lines (Radecke et al., 1996) or from protein expression plasmids (Lawson et al., 1995), or heat shock procedures to increase rescue efficiencies (Parks et al., 1999). Based on the T7 polymerase system, Bridgen and Elliott (1996) created Bunyamwera virus (family Bunyaviridae) from cloned cDNAs and demonstrated the feasibility of artificially generating a segmented negative-sense RNA virus by the T7 polymerase system.

In 1999, a plasmid-based reverse genetics technique was generated based on the cellular RNA polymerase I for the generation of segmented influenza A virus entirely from cloned cDNAs (Fodor et al., 1999; Neumann and Kawaoka, 1999). RNA polymerase I, a nucleolar enzyme, synthesizes ribosomal RNA which, like influenza virus RNA, does not contain 5' cap or 3' polyA structures. The RNA polymerase I transcription of a construct containing an influenza viral cDNA, flanked by RNA polymerase I promoter and terminator sequences, resulted in influenza vRNA synthesis (Fodor et al., 1999; Neumann and Kawaoka, 1999; Neumann and Kawaoka, 2001; Pekosz et al., 1999). The system was highly efficient, producing more than 10^8 infectious virus particles per ml of supernatant of plasmid-transfected cells 48 hours post-transfection.

What is needed is a method to prepare high titer orthomyxoviruses such as influenza A virus, entirely from cloned cDNAs.

SUMMARY OF THE INVENTION

The invention provides a composition comprising a plurality of influenza virus vectors of the invention, e.g., those useful to prepare reassortant viruses including 7:1 reassortants, 6:1:1 reassortants, 5:1:2 reassortants, and 5:1:1:1 reassortants. In one embodiment of the invention, the composition includes vectors for vRNA production selected from a vector comprising a promoter operably linked to an influenza virus PA cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus PB1 cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus PB2 cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus HA cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus NP cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus NA cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus M cDNA linked to a transcription termination sequence, and a vector comprising a promoter operably linked to an influenza virus NS cDNA linked to a transcription termination sequence. The composition also includes vectors for viral protein production selected from a vector encoding influenza virus PA, a vector encoding influenza virus PB1, a vector encoding influenza virus PB2, and a vector encoding influenza virus NP, and

optionally one or more vectors encoding NP, NS, M, e.g., M1 and M2, HA or NA. Preferably, the vectors encoding viral proteins further comprise a transcription termination sequence.

In one embodiment, the cDNAs for PB1, PB2, PA, NP, M, and NS, and optionally NA, have sequences for PB1, PB2, PA, NP, M, and NS, and optionally NA, from an influenza virus that replicates to high titers in embryonated eggs, and the cDNA for HA has sequences from a different strain of influenza virus (from a heterologous influenza virus isolate with the same or a different HA subtype, i.e., a heterologous HA). For HA from pathogenic H5N1 viruses which do not grow to high titers in embryonated eggs, the cDNA for at least NA has sequences from a N1 influenza virus that replicates to high titers in embryonated eggs.

In one embodiment, the cDNAs for PB1, PB2, PA, NP, M, and NS include a nucleic acid molecule corresponding to a sequence (polynucleotide) encoding at least one of the proteins of a high titer, e.g., titers greater than 10^8 EID₅₀/mL, e.g., 10^9 EID₅₀/mL, 10^{10} EID₅₀/mL, or more, influenza virus. Reassortants within the scope of the invention that have high titers in embryonated eggs have titers of at least about 10^9 EID₅₀/mL for 5:1:1:1 reassortants (with NS K55), 5:1:2 reassortants (with NS K55) and 6:1:1 reassortants (with NS K55) and at least 4×10^8 PFU/mL for 5:1:1:1 reassortants (with NS K55E) or 5:1:2 reassortants (with NS K55E). Reassortants within the scope of the invention that have high titers in MDCK cells have titers of at least 0.75×10^8 PFU/mL, e.g., at least 2.0×10^8 PFU/mL, for 5:1:1:1 or 6:1:1.

In one embodiment, the invention includes a composition comprising a plurality of influenza virus vectors for a 5:1:2 or a 6:1:1 reassortant. The composition includes a vector comprising a promoter operably linked to an influenza virus PA cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus PB1 cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus PB2 cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus HA cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus NP cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus NA cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus M cDNA linked to a transcription termination sequence, and a vector comprising a promoter operably linked to an influenza virus NS cDNA linked to a transcription termination sequence. The cDNAs for PB1, PB2, PA, NP, and M have sequences that are from one or more influenza viruses that replicate to high titers in embryonated eggs, wherein the cDNA for NS is from the one or more influenza viruses that replicate to high titers in embryonated eggs, and the cDNA for NA is from the one or more influenza viruses that replicate to high titers in embryonated eggs or has sequences for a heterologous NA. The cDNA for HA has sequences for a heterologous HA, which is heterologous to at least the viral gene segments for PB1, PB2, PA, NP, and M. In one embodiment, the cDNA for NS has a Glu at position 55. The composition also includes a vector comprising a promoter operably linked to a DNA segment encoding influenza virus PA, a vector comprising a promoter operably linked to a DNA segment encoding influenza virus PB1, a vector comprising a promoter operably linked to a DNA segment encoding influenza virus PB2, and a vector comprising a promoter operably linked to a DNA segment encoding influenza virus NP, and optionally a vector comprising

ing a promoter operably linked to a DNA segment encoding influenza virus HA, a vector comprising a promoter operably linked to a DNA segment encoding influenza virus NA, a vector comprising a promoter operably linked to a DNA segment encoding influenza virus M1, a vector comprising a promoter operably linked to a DNA segment encoding influenza virus M2, or a vector comprising a promoter operably linked to a DNA segment encoding influenza virus NS2. In one embodiment, the cDNAs for PB1, PB2, PA, NP, M, and NS include a nucleic acid molecule corresponding to a sequence (polynucleotide) encoding at least one of the proteins of a high titer, e.g., titers greater than 10^8 EID₅₀/mL, e.g., 10^9 EID₅₀/mL, 10^{10} EID₅₀/mL, or more, influenza virus.

In one embodiment, a composition comprising a plurality of influenza virus vectors for a 5:1:1:1 or 6:1:1 reassortant. The composition includes comprising a vector comprising a promoter operably linked to an influenza virus PA cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus PB1 cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus PB2 cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus HA cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus NP cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus NA cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus M cDNA linked to a transcription termination sequence, and a vector comprising a promoter operably linked to an influenza virus NS cDNA linked to a transcription termination sequence. The cDNAs for PB1, PB2, PA, NP, and M have sequences from one or more influenza viruses that replicate to high titers in MDCK cells, wherein the cDNA for NS is from the one or more influenza viruses that replicate to high titers in MDCK cells, wherein the cDNA for NA may have sequences for a heterologous NA, and wherein the cDNA for HA has sequences for a heterologous HA. The composition also includes a vector comprising a promoter operably linked to a DNA segment encoding influenza virus PA, a vector comprising a promoter operably linked to a DNA segment encoding influenza virus PB1, a vector comprising a promoter operably linked to a DNA segment encoding influenza virus PB2, and a vector comprising a promoter operably linked to a DNA segment encoding influenza virus NP, and optionally a vector comprising a promoter operably linked to a DNA segment encoding influenza virus HA, a vector comprising a promoter operably linked to a DNA segment encoding influenza virus NA, a vector comprising a promoter operably linked to a DNA segment encoding influenza virus M1, a vector comprising a promoter operably linked to a DNA segment encoding influenza virus M2, or a vector comprising a promoter operably linked to a DNA segment encoding influenza virus NS2. In one embodiment, the cDNAs for PB1, PB2, PA, NP, M, and NS include a nucleic acid molecule corresponding to a sequence (polynucleotide) encoding at least one of the proteins of a high titer, e.g., titers greater than 10^8 EID₅₀/mL, e.g., 10^9 EID₅₀/mL, 10^{10} EID₅₀/mL, or more, influenza virus.

As described herein, recombinant (6:2 reassortant) viruses grow less well in eggs than does the wild-type PR8 strain, even though they possess the same PR8 "internal" genes (i.e., those other than the HA and NA). Since vigorous growth in eggs is an essential property of vaccine seed viruses used in the production of inactivated vaccines, H5N1 vaccine candidates were generated that grow as well as the PR8 donor strain

5

in eggs. It was found that HA-NA balance and PB1 function are important growth determinants. With this knowledge, a series of H5N1 viruses was produced with altered HA-NA combinations, with the PR8 background, to assess their growth in eggs against more conventional 6:2 reassortants, including the WHO-recommended NIBRG-14 virus. A 7:1 reassortant virus and one of the 6:2 reassortants showed enhanced growth in eggs. Thus, for vaccine viruses that generally produce low titers in eggs, replacement of at least the NA of the vaccine virus with the NA of an influenza virus that grows well in eggs, or replacement of all but the HA and NA, or all but the HA, of the vaccine virus, with the other viral gene segments from an influenza virus that grows to high titers in eggs, can result in significantly higher viral titers. The titers of the reassortant viruses of the invention may be 2-fold, 3-fold, or greater, e.g., 7-fold or greater, than the corresponding nonreassortant vaccine virus. As also described herein, the internal genes responsible for the high growth rate of reassortants in eggs having genes from two different PR8 virus isolates was determined. The highest viral titers were those where the majority of internal genes were from PR8HG (PR8(UW)). In particular, 5:1:2 reassortants (PR8(UW) PB1, PB2, PA, NP and M; PR8(Cam) NS; and H5N1 HA and NA) and 6:1:1 reassortants (PR8(UW) NA, PB1, PB2, PA, NP and M; PR8(Cam) NS; and H5 HA) had high titers in eggs.

As also described herein, the viral genes responsible for a high growth rate in MDCK cells, cells likely to be approved as a source of vaccine virus, was assessed. The highest growth rate in MDCK cells was found with PB2 from PR8(UW), NS from PR8(Cam) or NS K55E from PR8(UW), and a NA with a long stalk, e.g., a stalk greater than 20 amino acids but less than about 100 amino acids, e.g., greater than about 40 and up to about 80 amino acids in length. Thus 5:1:1:1 and 6:1:1 reassortants with PR8(UW) PA, PB1, PB2, NP and M, and NS K55E from PR8(UW) or PR8(Cam), NA from PR8(UW) or a heterologous NA source, and a heterologous HA, grew to the highest titers in MDCK cells.

In one embodiment, the nucleic acid molecule corresponds to a sequence encoding PB1, PB2, PA, NP, M, and NS, and optionally NA, having substantially the same activity as a corresponding polypeptide encoded by one of SEQ ID NOs: 1-6 or 8. As used herein, "substantially the same activity" includes an activity that is about 0.1%, 1%, 10%, 30%, 50%, 90%, e.g., up to 100% or more, or detectable protein level that is about 80%, 90% or more, the activity or protein level, respectively, of the corresponding full-length polypeptide. In one embodiment, the nucleic acid molecule corresponds to a sequence encoding a polypeptide which is substantially the same as, e.g., having at least 80%, e.g., 90%, 92%, 95%, 97% or 99%, contiguous amino acid sequence identity to, a polypeptide encoded by one of SEQ ID NOs: 1-6 or 8. In one embodiment, the isolated and/or purified nucleic acid molecule comprises a nucleotide sequence which is substantially the same as, e.g., having at least 50%, e.g., 60%, 70%, 80% or 90% or more contiguous nucleic acid sequence identity to, one of SEQ ID NOs: 1-6, 8, or 33 to 38 and, in one embodiment, also encodes a polypeptide having at least 80%, e.g., 90%, 92%, 95%, 97% or 99%, contiguous amino acid sequence identity to a polypeptide encoded by one of SEQ ID NOs: 1-6, 8, or 33 to 38. In one embodiment, the isolated and/or purified nucleic acid molecule encodes a polypeptide with one or more, for instance, 2, 5, 10, 15, 20 or more, conservative amino acid substitutions, e.g., conservative substitutions of up to 10% or 20% of the residues, relative to a polypeptide encoded by one of SEQ ID NOs: 1-6 or 8. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For

6

example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine and tryptophan; a group of amino acids having basic side chains is lysine, arginine and histidine; and a group of amino acids having sulfur-containing side chain is cysteine and methionine. In one embodiment, conservative amino acid substitution groups are: valine-leucine-isoleucine; phenylalanine-tyrosine; lysine-arginine; alanine-valine; glutamic-aspartic; and asparagine-glutamine. In one embodiment, the isolated and/or purified nucleic acid molecule encodes a polypeptide with one or more, for instance, 2, 3 or 4, nonconservative amino acid substitutions, relative to a polypeptide encoded by one of SEQ ID NOs: 1-6 or 33-38. For instance, a K55E NS and a S360Y PB2 substitution are nonconservative substitutions.

In another embodiment, the nucleic acid molecule having PB1, PB2, PA, NP, M, and NS, and optionally NA, sequences, or the complement thereof, hybridizes to one of SEQ ID NOs: 1-6, 8, or 33 to 38, the complement thereof, under low stringency, moderate stringency or stringent conditions. For example, the following conditions may be employed: 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50° C. with washing in 2X SSC, 0.1% SDS at 50° C. (low stringency), more desirably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50° C. with washing in 1X SSC, 0.1% SDS at 50° C. (moderate stringency), more desirably still in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50° C. with washing in 0.5X SSC, 0.1% SDS at 50° C. (stringent), preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50° C. with washing in 0.1X SSC, 0.1% SDS at 50° C. (more stringent), more preferably in 7% sodium dodecyl sulfate (SDS), 0.5 M NaPO₄, 1 mM EDTA at 50° C. with washing in 0.1X SSC, 0.1% SDS at 65° C. (very stringent). In one embodiment, the nucleic acid molecule encodes a polypeptide which is substantially the same as, e.g., having at least 50%, e.g., 60%, 70%, 80% or 90% or more contiguous nucleic acid sequence identity to, one of SEQ ID NOs: 1-6, or 33 to 38, and preferably has substantially the same activity as a corresponding full-length polypeptide encoded by one of SEQ ID NOs: 1-6, 8 or 33 to 28. Those nucleic acid molecules, or nucleic acid molecules from other Ni strains that grow well in eggs, may be employed with nucleic acid for any HA, e.g., H5.

Thus, nucleic acid molecule may be employed to express influenza proteins, to prepare chimeric genes, e.g., with other viral genes including other influenza virus genes, and/or to prepare recombinant virus. Thus, the invention also provides isolated polypeptides, recombinant virus, and host cells contacted with the nucleic acid molecules or recombinant virus of the invention.

The invention also provides a plurality of the following isolated and/or purified vectors: a vector comprising a promoter operably linked to an influenza virus PA cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus PB1 cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus PB2 cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus HA cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus NP cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to

an influenza virus NA cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus M cDNA linked to a transcription termination sequence, a vector comprising a promoter operably linked to an influenza virus NS cDNA linked to a transcription termination sequence, wherein at least one vector comprises sequences corresponding to those encoding PB1, PB2, PA, NP, M, NS, and optionally NA, or a portion thereof, having substantially the same activity as a corresponding polypeptide encoded by one of SEQ ID NOs: 1-6 or 8, e.g., a sequence encoding a polypeptide with at least 80% amino acid identity to a polypeptide encoded by one of SEQ ID NOs: 1-6, 8 or 33 to 38. Optionally, two vectors may be employed in place of the vector comprising a promoter operably linked to an influenza virus M cDNA linked to a transcription termination sequence, e.g., a vector comprising a promoter operably linked to an influenza virus M1 cDNA linked to a transcription termination sequence and a vector comprising a promoter operably linked to an influenza virus M2 cDNA linked to a transcription termination sequence.

The invention includes the use of isolated and purified vectors or plasmids, which express or encode influenza virus proteins, or express or encode influenza vRNA, both native and recombinant vRNA. Preferably, the vectors comprise influenza cDNA, e.g., influenza A (e.g., any influenza A gene including any of the 15 HA or 9 NA subtypes), B or C DNA (see Chapters 45 and 46 of Fields *Virology* (Fields et al. (eds.), Lippincott-Raven Publ., Philadelphia, Pa. (1996), which are specifically incorporated by reference herein), although it is envisioned that the gene(s) of any organism may be employed in the vectors or methods of the invention. The cDNA may be in the sense or antisense orientation relative to the promoter. Thus, a vector of the invention may encode an influenza virus protein (sense) or vRNA (antisense). Any suitable promoter or transcription termination sequence may be employed to express a protein or peptide, e.g., a viral protein or peptide, a protein or peptide of a nonviral pathogen, or a therapeutic protein or peptide.

A composition of the invention may also comprise a gene or open reading frame of interest, e.g., a foreign gene encoding an immunogenic peptide or protein useful as a vaccine. Thus, another embodiment of the invention comprises a composition of the invention as described above in which one of the vectors is replaced with, or the composition further comprises, a vector comprising a promoter linked to 5' influenza virus sequences optionally including 5' influenza virus coding sequences or a portion thereof, linked to a desired nucleic acid sequence, e.g., a desired cDNA, linked to 3' influenza virus sequences optionally including 3' influenza virus coding sequences or a portion thereof, linked to a transcription termination sequence. Preferably, the desired nucleic acid sequence such as a cDNA is in an antisense orientation. The introduction of such a composition to a host cell permissive for influenza virus replication results in recombinant virus comprising vRNA corresponding to sequences of the vector. The promoter in such a vector for vRNA production may be a RNA polymerase I promoter, a RNA polymerase II promoter, a RNA polymerase III promoter, a T7 promoter, and a T3 promoter, and optionally the vector comprises a transcription termination sequence such as a RNA polymerase I transcription termination sequence, a RNA polymerase II transcription termination sequence, a RNA polymerase III transcription termination sequence, or a ribozyme. In one embodiment, the vector comprising the desired nucleic acid sequence comprises a cDNA of interest. The cDNA of interest, whether in a vector for vRNA or protein production, may encode an immunogenic epitope, such as an epitope useful in a cancer

therapy or vaccine, or a peptide or polypeptide useful in gene therapy. When preparing virus, the vector or plasmid comprising the gene or cDNA of interest may substitute for a vector or plasmid for an influenza viral gene or may be in addition to vectors or plasmids for all influenza viral genes.

A plurality of the vectors of the invention may be physically linked or each vector may be present on an individual plasmid or other, e.g., linear, nucleic acid delivery vehicle.

The promoter or transcription termination sequence in a vRNA or virus protein expression vector may be the same or different relative to the promoter or any other vector. Preferably, the vector or plasmid which expresses influenza vRNA comprises a promoter suitable for expression in at least one particular host cell, e.g., avian or mammalian host cells such as canine, feline, equine, bovine, ovine, or primate cells including human cells, or preferably, for expression in more than one host.

In one embodiment, one or more vectors for vRNA production comprise a promoter including, but not limited to, a RNA polymerase I promoter, e.g., a human RNA polymerase I promoter, a RNA polymerase II promoter, a RNA polymerase III promoter, a T7 promoter, or a T3 promoter. Preferred transcription termination sequences for the vRNA vectors include, but are not limited to, a RNA polymerase I transcription termination sequence, a RNA polymerase II transcription termination sequence, a RNA polymerase III transcription termination sequence, or a ribozyme. Ribozymes within the scope of the invention include, but are not limited to, tetrahymena ribozymes, RNase P, hammerhead ribozymes, hairpin ribozymes, hepatitis ribozyme, as well as synthetic ribozymes.

In one embodiment, at least one vector for vRNA comprises a RNA polymerase II promoter linked to a ribozyme sequence linked to viral coding sequences linked to another ribozyme sequences, optionally linked to a RNA polymerase II transcription termination sequence. In one embodiment, at least 2 and preferably more, e.g., 3, 4, 5, 6, 7 or 8, vectors for vRNA production comprise a RNA polymerase II promoter, a first ribozyme sequence, which is 5' to a sequence corresponding to viral sequences including viral coding sequences, which is 5' to a second ribozyme sequence, which is 5' to a transcription termination sequence. Each RNA polymerase II promoter in each vRNA vector may be the same or different as the RNA polymerase II promoter in any other vRNA vector. Similarly, each ribozyme sequence in each vRNA vector may be the same or different as the ribozyme sequences in any other vRNA vector. In one embodiment, the ribozyme sequences in a single vector are not the same.

The invention also provides a method to prepare influenza virus. The method comprises contacting a cell with a plurality of the vectors of the invention, e.g., sequentially or simultaneously, for example, employing a composition of the invention, in an amount effective to yield infectious influenza virus. The invention also includes isolating virus from a cell contacted with the composition. Thus, the invention further provides isolated virus, as well as a host cell contacted with the composition or virus of the invention. In another embodiment, the invention includes contacting the cell with one or more vectors, either vRNA or protein production vectors, prior to other vectors, either vRNA or protein production vectors.

The methods of producing virus described herein, which do not require helper virus infection, are useful in viral mutagenesis studies, and in the production of vaccines (e.g., for AIDS, influenza, hepatitis B, hepatitis C, rhinovirus, filoviruses, malaria, herpes, and foot and mouth disease) and gene therapy vectors (e.g., for cancer, AIDS, adenosine

deaminase, muscular dystrophy, omithine transcarbamylase deficiency and central nervous system tumors). Thus, a virus for use in medical therapy (e.g., for a vaccine or gene therapy) is provided.

The invention also provides a method to immunize an individual against a pathogen, e.g., a bacteria, virus, or parasite, or a malignant tumor. The method comprises administering to the individual an amount of at least one isolated virus of the invention, optionally in combination with an adjuvant, effective to immunize the individual. The virus comprises vRNA comprising a polypeptide encoded by the pathogen or a tumor-specific polypeptide.

Also provided is a method to augment or increase the expression of an endogenous protein in a mammal having an indication or disease characterized by a decreased amount or a lack of the endogenous protein. The method comprises administering to the mammal an amount of an isolated virus of the invention effective to augment or increase the amount of the endogenous protein in the mammal. Preferably, the mammal is a human.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1. Titer of various influenza viruses.

FIG. 2. Schematic diagram of the Ni NAs used to generate H5N1/PR8 reassortant viruses by reverse genetics. VN1203fill contains a 20 amino acid (aa) insertion derived from the N1 of the H5N1 precursor strain, GsGd96. VN1203fill.N2 contains, in addition to 20 aa from GsGd96 NA, a 14-aa insertion from N2 NA, resulting in a 34-aa insertion into the stalk of VN1203 NA. VN1202fill.N2N9 contains, in addition to 20 aa from GsGd96 NA and 14 aa from N2 NA, a 14-aa insertion from N9 NA, resulting in a 48-aa insertion into the stalk of VN1203. The predicted total length of the stalk region of each NA is given beneath each molecule.

FIG. 3. Growth of H5N1/PR8 reassortant viruses in chicken embryonated eggs. The titers of the reassortant viruses containing avirulent-form VN1203 HA and either homologous NA (VN1203) or heterologous NAs (VN1203fill, VN1203fill.N2, HK213, or PR8) with a PR8 background were compared by plaque titration with MDCK cells. The titer of wild-type (egg-adapted) PR8 also is included for comparison. The data are reported as mean titers and standard deviations for 3 eggs inoculated with each virus.

FIG. 4. Growth kinetics of H5N1 reassortant viruses in chicken embryonated eggs. We inoculated eggs with the same amounts (10^4 EID₅₀) of viruses containing PR8 NA (PR8), VN1203 NA (VN1203), or VN1203fill NA (VN1203fill). Mean HA titers and standard deviations for 3 eggs inoculated with each virus were determined at the indicated time points.

FIG. 5. Virus elution from chicken erythrocytes. Twofold dilutions of each virus (HA titers of 1:1024) containing VN1203 NA with a different stalk length, or PR8 NA, were incubated with chicken erythrocytes in a microtiter plate at 4° C. for 1 hour. The plate was then stored at 37° C. and reductions in the HA titer were recorded for 8 hours.

FIG. 6. Growth comparison of H5N1/PR8 reassortant viruses in chicken embryonated eggs. Viral titers of the 6:2 and 7:1 reassortant viruses, including the WHO-recommended NIBRG-14 strain (a VN1194/PR8 6:2 reassortant virus) were compared by plaque titration with MDCK cells. Mean titers and standard deviations of 3 eggs inoculated with each virus are shown. Thus, replacing just the NA of H5N1 viruses with the NA of PR8 may improve titers in eggs.

FIG. 7. Growth of reassortant H5N1 viruses possessing PR8(UW) or PR8(Cambridge) internal genes in chicken

embryonated eggs. Asterisks indicate a significant ($p < 0.05$, Student t-test) reduction in infectivity compared to PR8 (UW)/1194.

FIG. 8. The effect of the M and NS genes on the growth of viruses in chicken embryonated eggs. The asterisk indicates a significant ($p < 0.05$, Student t-test) increase in infectivity compared to PR8(UW)/1194.

FIG. 9. Growth of PR8(UW)/1194 and NIBRG-14 virus in MDCK cells.

FIG. 10. Identification of a gene segment responsible for the enhanced growth of PR8(UW)/1194 relative to NIBRG-14 in MDCK cells.

FIG. 11. Identification of the amino acid in PB2 responsible for the high growth rate of the vaccine seed virus in MDCK cells.

FIG. 12. Growth rates in MDCK cells of reassortants with different HA, NA, and NS genes. The asterisk indicates significantly better virus growth compared to that of PR8(UW)/1194. Double asterisks indicate significantly better growth rates compared to viruses expressing PR8(UW) NS.

FIG. 13. Growth in MDCK cells of the H5N1 vaccine seed virus containing a heterologous NS segment.

FIG. 14. Schematic of the genotype of an H5N1 vaccine seed virus with high growth capacity in chicken embryonated eggs or MDCK cells.

FIG. 15. Nucleotide sequence for PR8(Cambridge) genes (SEQ ID NOs:33-38).

DETAILED DESCRIPTION OF THE INVENTION

Definitions

As used herein, the terms “isolated and/or purified” refer to in vitro preparation, isolation and/or purification of a vector, plasmid or virus of the invention, so that it is not associated with in vivo substances, or is substantially purified from in vitro substances. An isolated virus preparation is generally obtained by in vitro culture and propagation, and/or via passage in eggs, and is substantially free from other infectious agents.

As used herein, “substantially free” means below the level of detection for a particular infectious agent using standard detection methods for that agent.

A “recombinant” virus is one which has been manipulated in vitro, e.g., using recombinant DNA techniques, to introduce changes to the viral genome. Reassortant viruses can be prepared by recombinant or nonrecombinant techniques.

As used herein, the term “recombinant nucleic acid” or “recombinant DNA sequence or segment” refers to a nucleic acid, e.g., to DNA, that has been derived or isolated from a source, that may be subsequently chemically altered in vitro, so that its sequence is not naturally occurring, or corresponds to naturally occurring sequences that are not positioned as they would be positioned in the native genome. An example of DNA “derived” from a source, would be a DNA sequence that is identified as a useful fragment, and which is then chemically synthesized in essentially pure form. An example of such DNA “isolated” from a source would be a useful DNA sequence that is excised or removed from said source by chemical means, e.g., by the use of restriction endonucleases, so that it can be further manipulated, e.g., amplified, for use in the invention, by the methodology of genetic engineering.

As used herein, a “heterologous” influenza virus gene or gene segment is from an influenza virus source that is different than a majority of the other influenza viral genes or gene segments in a reassortant influenza virus.

Influenza Virus Replication

Influenza A viruses possess a genome of eight single-stranded negative-sense viral RNAs (vRNAs) that encode a total of ten proteins. The influenza virus life cycle begins with binding of the hemagglutinin (HA) to sialic acid-containing receptors on the surface of the host cell, followed by receptor-mediated endocytosis. The low pH in late endosomes triggers a conformational shift in the HA, thereby exposing the N-terminus of the HA2 subunit (the so-called fusion peptide). The fusion peptide initiates the fusion of the viral and endosomal membrane, and the matrix protein (M1) and RNP complexes are released into the cytoplasm. RNPs consist of the nucleoprotein (NP), which encapsidates vRNA, and the viral polymerase complex, which is formed by the PA, PB1, and PB2 proteins. RNPs are transported into the nucleus, where transcription and replication take place. The RNA polymerase complex catalyzes three different reactions: synthesis of an mRNA with a 5' cap and 3' polyA structure, of a full-length complementary RNA (cRNA), and of genomic vRNA using the cDNA as a template. Newly synthesized vRNAs, NP, and polymerase proteins are then assembled into RNPs, exported from the nucleus, and transported to the plasma membrane, where budding of progeny virus particles occurs. The neuraminidase (NA) protein plays a crucial role late in infection by removing sialic acid from sialyloligosaccharides, thus releasing newly assembled virions from the cell surface and preventing the self aggregation of virus particles. Although virus assembly involves protein-protein and protein-vRNA interactions, the nature of these interactions is largely unknown.

Although influenza B and C viruses are structurally and functionally similar to influenza A virus, there are some differences. For example, influenza B virus does not have a M2 protein. Similarly, influenza C virus does not have a M2 protein.

Cell Lines and Influenza Viruses That Can Be Used in the Present Invention

According to the present invention, any cell which supports efficient replication of influenza virus can be employed in the invention, including mutant cells which express reduced or decreased levels of one or more sialic acids which are receptors for influenza virus. Viruses obtained by the methods can be made into a reassortant virus.

Preferably, the cells are WHO certified, or certifiable, continuous cell lines. The requirements for certifying such cell lines include characterization with respect to at least one of genealogy, growth characteristics, immunological markers, virus susceptibility, tumorigenicity and storage conditions, as well as by testing in animals, eggs, and cell culture. Such characterization is used to confirm that the cells are free from detectable adventitious agents. In some countries, karyology may also be required. In addition, tumorigenicity is preferably tested in cells that are at the same passage level as those used for vaccine production. The virus is preferably purified by a process that has been shown to give consistent results, before being inactivated or attenuated for vaccine production (see, e.g., World Health Organization, 1982).

It is preferred to establish a complete characterization of the cell lines to be used, so that appropriate tests for purity of the final product can be included. Data that can be used for the characterization of a cell to be used in the present invention includes (a) information on its origin, derivation, and passage history; (b) information on its growth and morphological characteristics; (c) results of tests of adventitious agents; (d) distinguishing features, such as biochemical, immunological, and cytogenetic patterns which allow the cells to be clearly recognized among other cell lines; and (e) results of tests for

tumorigenicity. Preferably, the passage level, or population doubling, of the host cell used is as low as possible.

It is preferred that the virus produced in the cell is highly purified prior to vaccine or gene therapy formulation. Generally, the purification procedures will result in the extensive removal of cellular DNA, other cellular components, and adventitious agents. Procedures that extensively degrade or denature DNA can also be used. See, e.g., Mizrahi, 1990.

Vaccines

A vaccine of the invention may comprise immunogenic proteins including glycoproteins of any pathogen, e.g., an immunogenic protein from one or more bacteria, viruses, yeast or fungi. Thus, in one embodiment, the influenza viruses of the invention may be vaccine vectors for influenza virus or other viral pathogens including but not limited to lentiviruses such as HIV, hepatitis B virus, hepatitis C virus, herpes viruses such as CMV or HSV or foot and mouth disease virus.

A complete virion vaccine is concentrated by ultrafiltration and then purified by zonal centrifugation or by chromatography. It is inactivated before or after purification using formalin or beta-propiolactone, for instance.

A subunit vaccine comprises purified glycoproteins. Such a vaccine may be prepared as follows: using viral suspensions fragmented by treatment with detergent, the surface antigens are purified, by ultracentrifugation for example. The subunit vaccines thus contain mainly HA protein, and also NA. The detergent used may be cationic detergent for example, such as hexadecyl trimethyl ammonium bromide (Bachmeyer, 1975), an anionic detergent such as ammonium deoxycholate (Laver & Webster, 1976); or a nonionic detergent such as that commercialized under the name TRITON X100. The hemagglutinin may also be isolated after treatment of the virions with a protease such as bromelain, then purified by a method such as that described by Grand and Skehel (1972).

A split vaccine comprises virions which have been subjected to treatment with agents that dissolve lipids. A split vaccine can be prepared as follows: an aqueous suspension of the purified virus obtained as above, inactivated or not, is treated, under stirring, by lipid solvents such as ethyl ether or chloroform, associated with detergents. The dissolution of the viral envelope lipids results in fragmentation of the viral particles. The aqueous phase is recuperated containing the split vaccine, constituted mainly of hemagglutinin and neuraminidase with their original lipid environment removed, and the core or its degradation products. Then the residual infectious particles are inactivated if this has not already been done.

Inactivated Vaccines. Inactivated influenza virus vaccines of the invention are provided by inactivating replicated virus of the invention using known methods, such as, but not limited to, formalin or β -propiolactone treatment. Inactivated vaccine types that can be used in the invention can include whole-virus (WV) vaccines or subvirion (SV) (split) vaccines. The WV vaccine contains intact, inactivated virus, while the SV vaccine contains purified virus disrupted with detergents that solubilize the lipid-containing viral envelope, followed by chemical inactivation of residual virus.

In addition, vaccines that can be used include those containing the isolated HA and NA surface proteins, which are referred to as surface antigen or subunit vaccines. In general, the responses to SV and surface antigen (i.e., purified HA or NA) vaccines are similar. An experimental inactivated WV vaccine containing an NA antigen immunologically related to the epidemic virus and an unrelated HA appears to be less

effective than conventional vaccines (Ogra et al., 1977). Inactivated vaccines containing both relevant surface antigens are preferred.

Live Attenuated Virus Vaccines. Live, attenuated influenza virus vaccines, can also be used for preventing or treating influenza virus infection, according to known method steps. Attenuation is preferably achieved in a single step by transfer of attenuated genes from an attenuated donor virus to a replicated isolate or reassorted virus according to known methods (see, e.g., Murphy, 1993). Since resistance to influenza A virus is mediated by the development of an immune response to the HA and NA glycoproteins, the genes coding for these surface antigens must come from the reassorted viruses or high growth clinical isolates. The attenuated genes are derived from the attenuated parent. In this approach, genes that confer attenuation preferably do not code for the HA and NA glycoproteins. Otherwise, these genes could not be transferred to reassortants bearing the surface antigens of the clinical virus isolate.

Many donor viruses have been evaluated for their ability to reproducibly attenuate influenza viruses. As a non-limiting example, the A/Ann Arbor(AA)/6/60 (H2N2) cold adapted (ca) donor virus can be used for attenuated vaccine production (see, e.g., Edwards, 1994; Murphy, 1993). Additionally, live, attenuated reassortant virus vaccines can be generated by mating the ca donor virus with a virulent replicated virus of the invention. Reassortant progeny are then selected at 25° C., (restrictive for replication of virulent virus), in the presence of an H2N2 antiserum, which inhibits replication of the viruses bearing the surface antigens of the attenuated A/AA/6/60 (H2N2) ca donor virus.

A large series of H1N1 and H3N2 reassortants have been evaluated in humans and found to be satisfactorily: (a) infectious, (b) attenuated for seronegative children and immunologically primed adults, (c) immunogenic and (d) genetically stable. The immunogenicity of the ca reassortants parallels their level of replication. Thus, the acquisition of the six transferable genes of the ca donor virus by new wild-type viruses has reproducibly attenuated these viruses for use in vaccinating susceptible adults and children.

Other attenuating mutations can be introduced into influenza virus genes by site-directed mutagenesis to rescue infectious viruses bearing these mutant genes. Attenuating mutations can be introduced into non-coding regions of the genome, as well as into coding regions. Such attenuating mutations can also be introduced into genes other than the HA or NA, e.g., the PB2 polymerase gene (Subbarao et al., 1993). Thus, new donor viruses can also be generated bearing attenuating mutations introduced by site-directed mutagenesis, and such new donor viruses can be used in the reduction of live attenuated reassortants H1N1 and H3N2 vaccine candidates in a manner analogous to that described above for the A/AA/6/60 ca donor virus. Similarly, other known and suitable attenuated donor strains can be reassorted with influenza virus of the invention to obtain attenuated vaccines suitable for use in the vaccination of mammals (Enami et al., 1990; Muster et al., 1991; Subbarao et al., 1993).

It is preferred that such attenuated viruses maintain the genes from the virus that encode antigenic determinants substantially similar to those of the original clinical isolates. This is because the purpose of the attenuated vaccine is to provide substantially the same antigenicity as the original clinical isolate of the virus, while at the same time lacking infectivity to the degree that the vaccine causes minimal change of inducing a serious pathogenic condition in the vaccinated mammal.

The virus can thus be attenuated or inactivated, formulated and administered, according to known methods, as a vaccine to induce an immune response in an animal, e.g., a mammal. Methods are well-known in the art for determining whether such attenuated or inactivated vaccines have maintained similar antigenicity to that of the clinical isolate or high growth strain derived therefrom. Such known methods include the use of antisera or antibodies to eliminate viruses expressing antigenic determinants of the donor virus; chemical selection (e.g., amantadine or rimantidine); HA and NA activity and inhibition; and DNA screening (such as probe hybridization or PCR) to confirm that donor genes encoding the antigenic determinants (e.g., HA or NA genes) are not present in the attenuated viruses. See, e.g., Robertson et al., 1988; Kilbourne, 1969; Aymard-Henry et al., 1985; Robertson et al., 1992.

Pharmaceutical Compositions

Pharmaceutical compositions of the present invention, suitable for inoculation or for parenteral or oral administration, comprise attenuated or inactivated influenza viruses, optionally further comprising sterile aqueous or non-aqueous solutions, suspensions, and emulsions. The compositions can further comprise auxiliary agents or excipients, as known in the art. See, e.g., Berkow et al., 1987; *Avery's Drug Treatment*, 1987; Osol, 1980; Katzung, 1992. The composition of the invention is generally presented in the form of individual doses (unit doses).

Conventional vaccines generally contain about 0.1 to 200 µg, preferably 10 to 15 µg, of hemagglutinin from each of the strains entering into their composition. The vaccine forming the main constituent of the vaccine composition of the invention may comprise a virus of type A, B or C, or any combination thereof, for example, at least two of the three types, at least two of different subtypes, at least two of the same type, at least two of the same subtype, or a different isolate(s) or reassortant(s). Human influenza virus type A includes H1N1, H2N2 and H3N2 subtypes.

Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and/or emulsions, which may contain auxiliary agents or excipients known in the art. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl oleate. Carriers or occlusive dressings can be used to increase skin permeability and enhance antigen absorption. Liquid dosage forms for oral administration may generally comprise a liposome solution containing the liquid dosage form. Suitable forms for suspending liposomes include emulsions, suspensions, solutions, syrups, and elixirs containing inert diluents commonly used in the art, such as purified water. Besides the inert diluents, such compositions can also include adjuvants, wetting agents, emulsifying and suspending agents, or sweetening, flavoring, or perfuming agents. See, e.g., Berkow et al., 1992; Avery's, 1987; Osol, 1980; and Katzung, 1992.

When a composition of the present invention is used for administration to an individual, it can further comprise salts, buffers, adjuvants, or other substances which are desirable for improving the efficacy of the composition. For vaccines, adjuvants, substances which can augment a specific immune response, can be used. Normally, the adjuvant and the composition are mixed prior to presentation to the immune system, or presented separately, but into the same site of the organism being immunized. Examples of materials suitable for use in vaccine compositions are provided in Osol (1980).

Heterogeneity in a vaccine may be provided by mixing replicated influenza viruses for at least two influenza virus strains, such as 2-50 strains or any range or value therein.

Influenza A or B virus strains having a modern antigenic composition are preferred. According to the present invention, vaccines can be provided for variations in a single strain of an influenza virus, using techniques known in the art.

A pharmaceutical composition according to the present invention may further or additionally comprise at least one chemotherapeutic compound, for example, for gene therapy, immunosuppressants, anti-inflammatory agents or immune enhancers, and for vaccines, chemotherapeutics including, but not limited to, gamma globulin, amantadine, guanidine, hydroxybenzimidazole, interferon- α , interferon- β , interferon- γ , tumor necrosis factor- α , thiosemicarbazones, methisazone, rifampin, ribavirin, a pyrimidine analog, a purine analog, foscarnet, phosphonoacetic acid, acyclovir, dideoxynucleosides, a protease inhibitor, or ganciclovir. See, e.g., Katzung (1992), and the references cited therein on pages 798-800 and 680-681, respectively.

The composition can also contain variable but small quantities of endotoxin-free formaldehyde, and preservatives, which have been found safe and not contributing to undesirable effects in the organism to which the composition is administered.

Pharmaceutical Purposes

The administration of the composition (or the antisera that it elicits) may be for either a "prophylactic" or "therapeutic" purpose. When provided prophylactically, the compositions of the invention which are vaccines, are provided before any symptom of a pathogen infection becomes manifest. The prophylactic administration of the composition serves to prevent or attenuate any subsequent infection. When provided prophylactically, the gene therapy compositions of the invention, are provided before any symptom of a disease becomes manifest. The prophylactic administration of the composition serves to prevent or attenuate one or more symptoms associated with the disease.

When provided therapeutically, an attenuated or inactivated viral vaccine is provided upon the detection of a symptom of actual infection. The therapeutic administration of the compound(s) serves to attenuate any actual infection. See, e.g., Berkow et al., 1992; Avery, 1987; and Katzung, 1992. When provided therapeutically, a gene therapy composition is provided upon the detection of a symptom or indication of the disease. The therapeutic administration of the compound(s) serves to attenuate a symptom or indication of that disease.

Thus, an attenuated or inactivated vaccine composition of the present invention may thus be provided either before the onset of infection (so as to prevent or attenuate an anticipated infection) or after the initiation of an actual infection. Similarly, for gene therapy, the composition may be provided before any symptom of a disorder or disease is manifested or after one or more symptoms are detected.

A composition is said to be "pharmacologically acceptable" if its administration can be tolerated by a recipient patient. Such an agent is said to be administered in a "therapeutically effective amount" if the amount administered is physiologically significant. A composition of the present invention is physiologically significant if its presence results in a detectable change in the physiology of a recipient patient, e.g., enhances at least one primary or secondary humoral or cellular immune response against at least one strain of an infectious influenza virus.

The "protection" provided need not be absolute, i.e., the influenza infection need not be totally prevented or eradicated, if there is a statistically significant improvement compared with a control population or set of patients. Protection

may be limited to mitigating the severity or rapidity of onset of symptoms of the influenza virus infection.

Pharmaceutical Administration

A composition of the present invention may confer resistance to one or more pathogens, e.g., one or more influenza virus strains, by either passive immunization or active immunization. In active immunization, an inactivated or attenuated live vaccine composition is administered prophylactically to a host (e.g., a mammal), and the host's immune response to the administration protects against infection and/or disease. For passive immunization, the elicited antisera can be recovered and administered to a recipient suspected of having an infection caused by at least one influenza virus strain. A gene therapy composition of the present invention may yield prophylactic or therapeutic levels of the desired gene product by active immunization.

In one embodiment, the vaccine is provided to a mammalian female (at or prior to pregnancy or parturition), under conditions of time and amount sufficient to cause the production of an immune response which serves to protect both the female and the fetus or newborn (via passive incorporation of the antibodies across the placenta or in the mother's milk).

The present invention thus includes methods for preventing or attenuating a disorder or disease, e.g., an infection by at least one strain of pathogen. As used herein, a vaccine is said to prevent or attenuate a disease if its administration results either in the total or partial attenuation (i.e., suppression) of a symptom or condition of the disease, or in the total or partial immunity of the individual to the disease. As used herein, a gene therapy composition is said to prevent or attenuate a disease if its administration results either in the total or partial attenuation (i.e., suppression) of a symptom or condition of the disease, or in the total or partial immunity of the individual to the disease.

At least one inactivated or attenuated influenza virus, or composition thereof, of the present invention may be administered by any means that achieve the intended purposes, using a pharmaceutical composition as previously described.

For example, administration of such a composition may be by various parenteral routes such as subcutaneous, intravenous, intradermal, intramuscular, intraperitoneal, intranasal, oral or transdermal routes. Parenteral administration can be by bolus injection or by gradual perfusion over time. A preferred mode of using a pharmaceutical composition of the present invention is by intramuscular or subcutaneous application. See, e.g., Berkow et al., 1992; Avery, 1987; and Katzung, 1992.

A typical regimen for preventing, suppressing, or treating an influenza virus related pathology, comprises administration of an effective amount of a vaccine composition as described herein, administered as a single treatment, or repeated as enhancing or booster dosages, over a period up to and including between one week and about 24 months, or any range or value therein.

According to the present invention, an "effective amount" of a composition is one that is sufficient to achieve a desired biological effect. It is understood that the effective dosage will be dependent upon the age, sex, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect wanted. The ranges of effective doses provided below are not intended to limit the invention and represent preferred dose ranges. However, the most preferred dosage will be tailored to the individual subject, as is understood and determinable by one of skill in the art. See, e.g., Berkow et al., 1992; Avery's, 1987; and Katzung, 1992.

17

The dosage of an attenuated virus vaccine for a mammalian (e.g., human) or avian adult organism can be from about 10^3 - 10^7 plaque forming units (PFU)/kg, or any range or value therein. The dose of inactivated vaccine can range from about 0.1 to 200, e.g., 50 μ g of hemagglutinin protein. However, the dosage should be a safe and effective amount as determined by conventional methods, using existing vaccines as a starting point.

The dosage of immunoreactive HA in each dose of replicated virus vaccine can be standardized to contain a suitable amount, e.g., 1-50 μ g or any range or value therein, or the amount recommended by the U.S. Public Health Service (PHS), which is usually 15 μ g, per component for older children 3 years of age, and 7.5 μ g per component for older children <3 years of age. The quantity of NA can also be standardized, however, this glycoprotein can be labile during the processor purification and storage (Kendal et al., 1980). Each 0.5-ml dose of vaccine preferably contains approximately 1-50 billion virus particles, and preferably 10 billion particles.

The invention will be further described by the following nonlimiting examples.

EXAMPLE 1

To develop a reverse genetics system for influenza A/Puerto Rico/8/34, viral RNA was extracted from the allantoic fluid of A/Puerto Rico/8/34 (H1N1), Madison high grower variant (PR8HG), using RNeasy Mini kit (Qiagen) according to the manufacturer's protocol. cDNA was synthesized using MMLV-RTase (Promega) and Uni12 primer. The cDNAs were amplified overnight by PCR using the following:

PB1:

Ba PB1-1 and PB1-1735R (front fragment) and PB1-903 and Ba-PB1-2341R (rear fragment)

Ba-PB1-1

(SEQ ID NO:9)

CACACACGGTCTCCGGGAGCGAAAGCAGGCA

173PB1-1735R

(SEQ ID NO:28)

GGGTTTGATTTGTGTGTACAC

233PB1-903

(SEQ ID NO:10)

CCAGGACACTGAAATTTCTTTCAC

Ba-PB1-2341R

(SEQ ID NO:11)

CACACAGGTCTCCTATTAGTAGAAACAAGGCATTT

PB2:

Ba PB2-1 and B2 1260R (front fragment) and WSN PB2 seq-2 and Ba-PB2-2341R (rear fragment)

Ba-PB2-1

(SEQ ID NO:12)

CACACAGGTCTCCGGGAGCGAAAGCAGGTC

B2 1260R

(SEQ ID NO:13)

CACACACGTCTCCATCATAAATCCTCTTG

WSN PB2 seq-2

(SEQ ID NO:14)

CTCCTCTGATGGTGGCATACT

Ba-PB2-2341R

(SEQ ID NO:15)

CACACAGGTCTCCTATTAGTAGAAACAAGGTCGTTT

18

-continued

PA:

Bm-PA-1

(SEQ ID NO:16)

CACACACGTCTCCGGGAGCGAAAGCAGGTAC

5

Bm-PA-2233R

(SEQ ID NO:17)

CACACACGTCTCCTATTAGTAGAAACAAGGTACTT

HA:

10 Bm-HA-1:

(SEQ ID NO:18)

CACACACGTCTCCGGGAGCAAAAGCAGGGG

Bm-NS-890R:

(SEQ ID NO:19)

15

CACACACGTCTCCTATTAGTAGAAACAAGGGTGTGTTT

NP

Bm-NP-1

(SEQ ID NO:20)

CACACACGTCTCCGGGAGCAAAAGCAGGGTA

20

Bm-NP-1565R

(SEQ ID NO:21)

CACACACGTCTCCTATTAGTAGAAACAAGGGTATTTTT

NA:

25 Ba-NA-1:

(SEQ ID NO:22)

CACACAGGTCTCCGGGAGCAAAAGCAGGAGT

Ba-NA-1413R:

(SEQ ID NO:23)

CACACAGGTCTGGTATTAGTAGAAACAAGGAGTTTTTT

30

M:

Bm-M-1

(SEQ ID NO:24)

CACACACGTCTCCGGGAGCAAAAGCAGGTAG

35

Bm-M-1027R

(SEQ ID NO:25)

CACACACGTCTCCTATTAGTAGAAACAAGGTAGTTTTT

NS:

Bm-NS-1

(SEQ ID NO:26)

40

CACACACGTCTCCGGGAGCAAAAGCAGGGTG

Bm-NS-890R

(SEQ ID NO:27)

CACACACGTCTCCTATTAGTAGAAACAAGGGTGTGTTT

45

DNA polymerase: pfu Native DNA polymerase (Stratagene)

The PCR products were separated by gel electrophoresis and extracted from the agarose gel using a gel extraction kit (Qiagen). The extracted genes were ligated into pT7Blue blunt vector (Novagen) using a Takara ligation kit ver. II (Takara). After 5 hours, the ligated genes were transformed into JM109 (PB2, M, and NS genes) or DH5alpha (PA, PB1, and NP). Six colonies for each gene were cultured in TB for 8 hours. The plasmids were extracted from the bacteria culture, and four clones per gene were sequenced.

The PA, NP, M, and NS genes in pT7Blue were excised by Bsm BI enzyme (New England Biolabs). The PB1 gene was excised by Bsa I (New England Biolabs). The excised genes were ligated overnight with pPollR vector which contains the human RNA polymerase I promoter and the mouse RNA polymerase I terminator which had been digested with Bsm BI. The front fragment of the PB2 gene in pT7Blue was excised by Bsr GI (New England Biolabs) and Bam HI (Roche), and the rear fragment was excised by Bsr GI (New England Biolabs) and Spe I (Roche). The excised fragments were mixed and digested by Bsa I. After 6 hours, the digested

genes were purified using a PCR purification kit (Qiagen) and ligated overnight between the Bsm BI sites of the pPolIR vector.

The ligated PB1, PA, NP, M, and NS-pPolIR genes were used to transform JM109 (M and NS genes) or DH5alpha (PB1, PA and NP genes) overnight. The colonies of transformed bacteria were cultured in LB overnight. The ligated PB2-pPolIR was used to transform JM109 overnight.

The plasmids were extracted from the bacterial cultures and gene inserts were confirmed by enzyme digestion. The colonies of bacteria transformed by PB2-PolIR were cultured in LB for 8 hours. The plasmids were then extracted and the gene insertion was confirmed by enzyme digestion. All pPol constructs were sequenced to ensure that they did not contain unwanted mutations.

The pPolIR constructs for PR8HG were transfected into 293T human embryonic kidney cells with A/WSN/33(WSN)-HA and NA, A/Hong Kong/483/97(HK)-HAavir and NA, or A/Kawasaki/01(Kawasaki)-HA and NA PolI constructs and four protein-expression constructs for the polymerase proteins and NP of A/WSN/33. The supernatants from transfected 293T cells were serially diluted (undiluted to 10^{-7}) and infected into the allantoic cavities of 9-day-old embryonated chicken eggs. The allantoic fluids of the infected eggs were harvested and their virus titers tested by HA assay (Table 1).

TABLE 1

Virus possessing PR8 genes together	HA titer (HAU/ml) of allantoic fluid from eggs inoculated with 293T supernatants diluted at:							
with the following HA and NA genes	un- dilut- ed	10 ⁻¹	10 ⁻²	10 ⁻³	10 ⁻⁴	10 ⁻⁵	10 ⁻⁶	10 ⁻⁷
WSN-HA NA	<1	<1	200	<1	<1	<1	<1	<1
HK-HAavir NA	100	<1	<1	<1	<1	<1	<1	<1
Kawasaki-HA NA	<1	<1	<1	<1	<1	<1	<1	<1

HA-positive samples (virus with WSN-HA NA at 10^{-2} and virus with HK-HAavir NA at undiluted) were diluted serially from 10^{-2} to 10^{-8} and 100 ul of each dilution was infected into embryonated chicken eggs. The allantoic fluids of the infected eggs were harvested and their virus titers tested by HA assay (Table 2). The 50% egg infectious dose (EID_{50}) of A/Puerto Rico/8/34 (H1N1) prepared from plasmids was $10^{10.33}$ /ml, and the HA titer was 1:3200.

A recombinant virus having the HA and NA genes from A/Hong Kong/213/2003 (H5N1) and the remainder of the type A influenza virus genes from PR8HG was prepared. The titer of the recombinant virus was $10^{10.67}$ EID_{50} /ml, and the HA titer was 1:1600

TABLE 2

Virus possessing PR8 genes together with the following HA and NA genes	HA titer (HAU/ml) in each dilution						
	10-2	10-3	10-4	10-5	10-6	10-7	10-8
WSN-HA NA	160	40	40	320	40	640	<1
HK-HAavir NA	400	800	400	400	400	800	<1

Sequences of PR8 genes:

PA	(SEQ ID NO:1)
5 AGCGAAAGCA GGTACTGATC CAAAATGGAA GATTTTGTGC	
GACAATGCTT	
CAATCCGATG ATTGTCGAGC TTGCGGAAAA AACAAATGAAA	
10 GAGTATGGGG	
AGGACCTGAA AATCGAAACA AACAAATTTG CAGCAATATG	
CACTCACTTG	
15 GAAGTATGCT TCATGTATTG AGATTTTCAC TTCATCAATG	
AGCAAGGCGA	
GTCAATAATC GTAGAACTTG GTGATCCAAA TGCATTTTGG	
20 AAGCACAGAT	
TTGAAATAAT CGAGGGAAGA GATCGCACAA TGGCCTGGAC	
AGTAGTAAAC	
25 AGTATTTGCA ACACTACAGG GGCTGAGAAA CCAAAGTTTC	
TACCAGATTT	
GTATGATTAC AAGGAGAATA GATTCATCGA AATTGGAGTA	
ACAAGGAGAG	
30 AAGTTCACAT ATACTATCTG GAAAAGGCCA ATAAAATTAA	
ATCTGAGAAA	
ACACACATCC ACATTTTCTC GTTCACTGGG GAAGAAATGG	
35 CCACAAAGGC	
AGACTACACT CTCGATGAAG AAAGCAGGGC TAGGATCAAA	
ACCAGACTAT	
40 TCACCATAAG ACAAGAAATG GCCAGCAGAG GCCTCTGGGA	
TTCTTTTCGT	
CAGTCCGAGA GAGGAGAAGA GACAATTGAA GAAAGGTTTG	
45 AAATCACAGG	
AACAATGCGC AAGCTTGCCG ACCAAAGTCT CCCGCCGAAC	
TTCTCCAGCC	
50 TTGAAAATTT TAGAGCCTAT GTGGATGGAT TCGAACCGAA	
CGGCTACATT	
GAGGGCAAGC TGTCTCAAAT GTCCAAAGAA GTAAATGCTA	
55 GAATTGAACC	
TTTTTTGAAA ACAACACCAC GACCACTTAG ACTTCCGAAT	
GGGCCTCCCT	
60 GTTCTCAGCG GTCCAAATTC CTGCTGATGG ATGCCTTAAA	
ATTAAGCATT	
GAGGACCCAA GTCATGAAGG AGAGGGAATA CCGCTATATG	
65 ATGCAATCAA	

21

-continued

ATGCATGAGA ACATTCTTTG GATGGAAGGA ACCCAATGTT
 GTTAAACCAC
 ACGAAAAGGG AATAAATCCA AATTATCTTC TGTCATGGAA
 GCAAGTACTG
 GCAGAACTGC AGGACATTGA GAATGAGGAG AAAATTCCAA
 AGACTAAAAA
 TATGAAGAAA ACAAGTCAGC TAAAGTGGGC ACTTGGTGAG
 AACATGGCAC
 CAGAAAAGGT AGACTTTGAC GACTGTAAAG ATGTAGGTGA
 TTTGAAGCAA
 TATGATAGTG ATGAACCAGA ATTGAGGTCG CTTGCAAGTT
 GGATTGAGAA
 TGAGTTTAAC AAGGCATGCG AACTGACAGA TTCAAGCTGG
 ATAGAGCTCG
 ATGAGATTGG AGAAGATGTG GCTCCAATTG AACACATTGC
 AAGCATGAGA
 AGGAATTATT TCACATCAGA GGTGTCTCAC TGCAGAGCCA
 CAGAATACAT
 AATGAAGGGA GTGTACATCA ATACTGCCTT GCTTAATGCA
 TCTTGTGCAG
 CAATGGATGA TTTCCAATTA ATTCCAATGA TAAGCAAGTG
 TAGAACTAAG
 GAGGGAAGGC GAAAGACCAA CTTGTATGGT TTCATCATAA
 AAGGAAGATC
 CCACTTAAGG AATGACACCG ACGTGGTAAA CTTTGTGAGC
 ATGGAGTTTT
 CTCTCACTGA CCCAAGACTT GAACCAATA AATGGGAGAA
 GTACTGTGTT
 CTTGAGATAG GAGATATGCT TATAAGAAGT GCCATAGGCC
 AGGTTTCAAG
 GCCCATGTTT TTGTATGTGA GAACAAATGG AACCTCAAAA
 ATTAAATGA
 AATGGGGAAT GGAGATGAGG CGTTGCCTCC TCCAGTCACT
 TCAACAAATT
 GAGAGTATGA TTGAAGCTGA GTCCTCTGTC AAAGAGAAAG
 ACATGACCAA
 AGAGTTCTTT GAGAACAAAT CAGAAACATG GCCCATTGGA
 GAGTCCCCCA
 AAGGAGTGGA GGAAAGTTCC ATTGGGAAGG TCTGCAGGAC
 TTTATTAGCA
 AAGTCGGTAT TCAACAGCTT GTATGCATCT CCACAAC TAG

22

-continued

AAGGATTTTC
 5 AGCTGAATCA AGAAACTGC TTCTTATCGT TCAGGCTCTT
 AGGGACAACC
 TGGAACCTGG GACCTTTGAT CTTGGGGGGC TATATGAAGC
 AATTGAGGAG
 10 TGCCCTGATTA ATGATCCCTG GGTTTTGCTT AATGCTTCTT
 GGTTCAACTC
 CTTCTTACA CATGCATTGA GTTAGTTGTG GCAGTGCTAC
 15 TATTTGCTAT
 CCATACTGTC CAAAAAAGTA CCTTGTTTCT ACT
 PB1
 20 AGCGAAAGCA GGCAAACCAT TTGAATGGAT GTCAATCCGA
 CTTACTTTT CTTAAAAGTG CCAGCACAAA ATGCTATAAG
 CACAACTTTC
 25 CCTTATACTG GAGACCCTCC TTACAGCCAT GGGACAGGAA
 CAGGATACAC
 CATGGATACT GTCAACAGGA CACATCAGTA CTCAGAAAAG
 30 GGAAGATGGA
 CAACAAACAC CGAAACTGGA GCACCGCAAC TCAACCCGAT
 TGATGGGCCA
 35 CTGCCAGAAG ACAATGAACC AAGTGGTTAT GCCCAAACAG
 ATTGTGTATT
 GGAGGCGATG GCTTTCCTTG AGGAATCCCA TCCTGGTATT
 40 TTTGAAACT
 CGTGTATTGA AACGATGGAG GTTGTTTCAGC AAACACGAGT
 AGACAAGCTG
 45 ACACAAGGCC GACAGACCTA TGACTGGACT CTAATAGAA
 ACCAACCTGC
 TGCAACAGCA TTGGCCAACA CAATAGAAGT GTTCAGATCA
 AATGGCCTCA
 50 CGGCCAATGA GTCTGGAAGG CTCATAGACT TCCTTAAGGA
 TGTAATGGAG
 TCAATGAACA AAGAAGAAAT GGGGATCACA ACTCATTTTC
 55 AGAGAAAGAG
 ACGGGTGAGA GACAATATGA CTAAGAAAAT GATAACACAG
 AGACAATGG
 60 GTAAAAAGAA GCAGAGATTG AACAAAAGGA GTTATCTAAT
 TAGAGCATTG
 ACCCTGAACA CAATGACCAA AGATGCTGAG AGAGGGAAGC
 65 TAAAACGGAG

(SEQ ID NO:2)

23

-continued

AGCAATTGCA ACCCCAGGGA TGCAATAAG GGGTTTGTA
 TACTTTGTTG
 AGACACTGGC AAGGAGTATA TGTGAGAAAC TTGAACAATC
 AGGGTTGCCA
 GTTGGAGGCA ATGAGAAGAA AGCAAAGTTG GCAAATGTTG
 TAAGGAAGAT
 GATGACCAAT TCTCAGGACA CCGAACTTTC TTTCACCATC
 ACTGGAGATA
 ACACCAAATG GAACGAAAAT CAGAATCCTC GGATGTTTTT
 GGCCATGATC
 ACATATATGA CCAGAAATCA GCCCGAATGG TTCAGAAATG
 TTCTAAGTAT
 TGCTCCAATA ATGTTCTCAA ACAAATGGC GAGACTGGGA
 AAAGGGTATA
 TGTTTGAGAG CAAGAGTATG AAAGTTAGAA CTCAAATACC
 TGCAGAAATG
 CTAGCAAGCA TCGATTTGAA ATATTTCAAT GATTCAACAA
 GAAAGAAGAT
 TGAAAAATC CGACCGCTCT TAATAGAGGG GACTGCATCA
 TTGAGCCCTG
 GAATGATGAT GGGCATGTTC AATATGTAA GCACTGTATT
 AGGCGTCTCC
 ATCCTGAATC TTGGACAAAA GAGATACACC AAGACTACTT
 ACTGGTGGGA
 TGGTCTTCAA TCCTCTGACG ATTTTGCTCT GATTGTGAAT
 GCACCCAATC
 ATGAAGGGAT TCAAGCCGGA GTCGACAGGT TTTATCGAAC
 CTGTAAGCTA
 CTTGGAATCA ATATGAGCAA GAAAAAGTCT TACATAAACA
 GAACAGGTAC
 ATTTGAATTC ACAAGTTTTT TCTATCGTTA TGGGTTTGT
 GCCAATTTCA
 GCATGGAGCT TCCCAGTTTT GGGGTGTCTG GGATCAACGA
 GTCAGCGGAC
 ATGAGTATTG GAGTTACTGT CATCAAAAC AATATGATAA
 ACAATGATCT
 TGGTCCAGCA ACAGCTCAAA TGGCCCTTCA GTTGTTCATC
 AAAGATTACA
 GGTACACGTA CCGATGCCAT ATAGGTGACA CACAAATACA
 AATCCGAAGA
 TCATTTGAAA TAAAGAACT GTGGGAGCAA ACCCGTTCCA

24

-continued

AAGCTGGACT
 5 GCTGGTCTCC GACGGAGGCC CAAATTTATA CAACATTAGA
 AATCTCCACA
 TTCCTGAAGT CTGCCTAAAA TGGGAATTGA TGGATGAGGA
 TTACCAGGGG
 10 CGTTTATGCA ACCCACTGAA CCCATTTGTC AGCCATAAAG
 AAATTGAATC
 AATGAACAAT GCAGTGATGA TGCCAGCACA TGGTCCAGCC
 15 AAAACATGG
 AGTATGATGC TGTTGCAACA ACACACTCCT GGATCCCCAA
 AAGAAATCGA
 20 TCCATCTTGA ATACAAGTCA AAGAGGAGTA CTGAGGATG
 AACAAATGTA
 CCAAAGGTGC TGCAATTTAT TTGAAAAAT CTCCCCAGC
 25 AGTTCATACA
 GAAGACCACT CGGGATATCC AGTATGGTGG AGGCTATGGT
 TTCCAGAGCC
 30 CGAATTGATG CACGGATTGA TTTCGAATCT GGAAGATAA
 AGAAAGAAGA
 GTTCACTGAG ATCATGAAGA TCTGTTCCAC CATTGAAGAG
 35 CTCAGACGGC
 AAAAATAGTG AATTTAGCTT GTCCTTCATG AAAAATGCC
 TTGTTTCTAC T
 PB2
 40 (SEQ ID NO:3)
 AGCGAAAGCA GGTCAATTAT ATTCAATATG GAAAGAATAA
 AAGAACTACG
 AAATCTAATG TCGCAGTCTC GCACCCGCGA GATACTCACA
 45 AAAACCACCG
 TGGACCATAT GGCCATAATC AAGAAGTACA CATCAGGAAG
 ACAGGAGAAG
 50 AATCCAGCAC TTAGGATGAA ATGGATGATG GCAATGAAAT
 ATCCAATTAC
 AGCAGACAAG AGGATAACGG AAATGATTCC TGAGAGAAAT
 55 GAGCAAGGAC
 AAATTTTATG GAGTAAATG AATGATGCCG GATCAGACCG
 AGTGATGGTA
 60 TCACCTCTGG CTGTGACATG GTGGAATAGG AATGGACCAA
 TAACAAATAC
 AGTTCATTAT CCAAAAATCT ACAAACCTTA TTTTGAAAGA
 65 GTCGAAAGGC

25

-continued

TAAAGCATGG AACCTTTGGC CCTGTCCATT TTAGAAACCA
 AGTCAAAATA
 CGTCGGAGAG TTGACATAAA TCCTGGTCAT GCAGATCTCA
 GTGCCAAGGA
 GGCACAGGAT GTAATCATGG AAGTTGTTTT CCCTAACGAA
 GTGGGAGCCA
 GGATACTAAC ATCGGAATCG CAACTAACGA TAACCAAAGA
 GAAGAAAGAA
 GAACTCCAGG ATTGCAAAAT TTCTCCTTG ATGGTTGCAT
 ACATGTTGGA
 GAGAGAACTG GTCCGCAAAA CGAGATTCCT CCCAGTGGCT
 GGTGGAACAA
 GCAGTGTGTA CATTGAAGTG TTGCATTGA CTCAGGAAC
 ATGCTGGGAA
 CAGATGTATA CTCCAGGAGG GGAAGTGAGG AATGATGATG
 TTGATCAAAG
 CTTGATTATT GCTGCTAGGA ACATAGTGAG AAGAGCTGCA
 GTATCAGCAG
 ATCCACTAGC ATCTTTATTG GAGATGTGCC ACAGCACACA
 GATTGGTGGA
 ATTAGGATGG TAGACATCCT TAGGCAGAAC CCAACAGAAG
 AGCAAGCCGT
 GGATATATGC AAGGCTGCAA TGGGACTGAG AATTAGCTCA
 TCCTTCAGTT
 TTGGTGGATT CACATTTAAG AGAACAAGCG GATCATCAGT
 CAAGAGAGAG
 GAAGAGGTGC TTACGGGCAA TCTCAAACA TTGAAGATAA
 GAGTGCATGA
 GGGATATGAA GAGTTCACAA TGGTTGGGAG AAGAGCAACA
 GCCATACTCA
 GAAAAGCAAC CAGGAGATTG ATTCAGCTGA TAGTGAGTGG
 GAGAGACGAA
 CAGTCGATTG CCGAAGCAAT AATTGTGGCC ATGGTATTTT
 CACAAGAGGA
 TTGTATGATA AAAGCAGTCA GAGGTGATCT GAATTCGTC
 AATAGGGCGA
 ATCAACGATT GAATCCTATG CATCAACTTT TAAGACATTT
 TCAGAAGGAT
 GCGAAAGTGC TTTTTCAAAA TTGGGGAGTT GAACCTATCG
 ACAATGTGAT
 GGGAATGATT GGGATATTGC CCGACATGAC TCCAAGCATC

26

-continued

GAGATGTCAA
 5 TGAGAGGAGT GAGAATCAGC AAAATGGGTG TAGATGAGTA
 CTCCAGCACG
 GAGAGGGTAG TGGTGAGCAT TGACCGTTTT TTGAGAATCC
 GGGACCAACG
 10 AGGAAATGTA CTACTGTCTC CCGAGGAGGT CAGTGAAACA
 CAGGGAACAG
 AGAAACTGAC AATAACTTAC TCATCGTCAA TGATGTGGGA
 15 GATTAATGGT
 CCTGAATCAG TGTTGGTCAA TACCTATCAA TGGATCATCA
 GAAACTGGGA
 20 AACTGTTAAA ATTCAGTGGT CCCAGAACCC TACAATGCTA
 TACAATAAAA
 TGGAAATTGA ACCATTTTCAG TCTTTAGTAC CTAAGGCCAT
 25 TAGAGGCCAA
 TACAGTGGGT TTGTAAGAAC TCTGTTCCAA CAAATGAGGG
 ATGTGCTTGG
 30 GACATTTGAT ACCGCACAGA TAATAAACT TCTTCCCTTC
 GCAGCCGCTC
 CACCAAAGCA AAGTAGAATG CAGTTCTCCT CATTTACTGT
 35 GAATGTGAGG
 GGATCAGGAA TGAGAATACT TGTAAGGGGC AATTCTCCTG
 TATTCAACTA
 40 TAACAAGGCC ACGAAGAGAC TCACAGTTCT CGGAAAGGAT
 GCTGGCACTT
 TAACTGAAGA CCCAGATGAA GGCACAGCTG GAGTGGAGTC
 CGCTGTTCTG
 45 AGGGGATTCC TCATTCTGGG CAAAGAAGAC AAGAGATATG
 GGCCAGCACT
 AAGCATCAAT GAACTGAGCA ACCTTGCGAA AGGAGAGAAG
 50 GCTAATGTGC
 TAATTGGGCA AGGAGACGTG GTGTTGGTAA TGAAACGGAA
 ACGGGACTCT
 55 AGCATACTTA CTGACAGCCA GACAGCGACC AAAAGAATTC
 GGATGGCCAT
 CAATTAGTGT CGAATAGTTT AAAACGACC TTGTTTCTAC
 60 T
 NP
 (SEQ ID NO: 4)
 AGCAAAAGCA GGGTAGATAA TCACTCACTG AGTGACATCA
 65 AAATCATGGC GTCTCAAGGC ACCAAACGAT CTTACGAACA

27

-continued

GATGGAGACT
 GATGGAGAAC GCCAGAATGC CACTGAAATC AGAGCATCCG
 TCGGAAAAAT
 GATTGGTGGA ATTGGACGAT TCTACATCCA AATGTGCACC
 GAACTCAAAC
 TCAGTGATTA TGAGGGACGG TTGATCCAAA ACAGCTTAAC
 AATAGAGAGA
 ATGGTGCTCT CTGCTTTTGA CGAAAGGAGA AATAAATACC
 TTGAAGAACA
 TCCCAGTGC GGGAAAGATC CTAAGAAAAC TGGAGGACCT
 ATATACAGGA
 GAGTAAACGG AAAGTGGATG AGAGAACTCA TCCTTTATGA
 CAAAGAAGAA
 ATAAGGCGAA TCTGGCGCCA AGCTAATAAT GGTGACGATG
 CAACGGCTGG
 TCTGACTCAC ATGATGATCT GGCATTCCAA TTTGAATGAT
 GCAACTTATC
 AGAGGACAAG AGCTCTTGTT CGCACCGGAA TGGATCCCAG
 GATGTGCTCT
 CTGATGCAAG GTTCAACTCT CCCTAGGAGG TCTGGAGCCG
 CAGGTGCTGC
 AGTCAAAGGA GTTGAACAA TGGTGATGGA ATTGGTCAGA
 ATGATCAAAC
 GTGGGATCAA TGATCGGAAC TTCTGGAGGG GTGAGAATGG
 ACGAAAAACA
 AGAATTGCTT ATGAAAGAAT GTGCAACATT CTCAAAGGGA
 AATTTCAAAC
 TGCTGCACAA AAAGCAATGA TGGATCAAGT GAGAGAGAGC
 CGGAACCCAG
 GGAATGCTGA GTTCGAAGAT CTCACCTTTC TAGCACGGTC
 TGCACTCATA
 TTGAGAGGGT CGGTGCTCA CAAGTCCTGC CTGCCTGCCT
 GTGTGTATGG
 ACCTGCCGTA GCCAGTGGGT ACGACTTTGA AAGGGAGGGA
 TACTCTCTAG
 TCGGAATAGA CCCTTTCAGA CTGCTTCAAA ACAGCCAAGT
 GTACAGCCTA
 ATCAGACCAA ATGAGAATCC AGCACACAAG AGTCAACTGG
 TGTGGATGGC
 ATGCCATTCT GCCGCATTG AAGATCTAAG AGTATTAAGC
 TTCATCAAAG

28

-continued

GGACGAAGGT GCTCCCAAGA GGGAAGCTTT CCACTAGAGG
 AGTTCAAATT
 5 GCTTCCAATG AAAATATGGA GACTATGGAA TCAAGTACAC
 TTGAACTGAG
 AAGCAGGTAC TGGGCCATAA GGACCAGAAG TGGAGGAAAC
 10 ACCAATCAAC
 AGAGGGCATC TGCGGGCCAA ATCAGCATAC AACCTACGTT
 CTCAGTACAG
 15 AGAAATCTCC CTTTGTACAG AACAACCATT ATGGCAGCAT
 TCAATGGGAA
 TACAGAGGGG AGAACATCTG ACATGAGGAC CGAAATCATA
 20 AGGATGATGG
 AAAGTGCAAG ACCAGAAGAT GTGTCTTTC AGGGGCGGGG
 AGTCTTCGAG
 25 CTCTCGGACG AAAAGGCAGC GAGCCCGATC GTGCCTTCCT
 TTGACATGAG
 TAATGAAGGA TCTTATTTCT TCGGAGACAA TGCAGAGGAG
 30 TACGACAATT
 AAAGAAAAAT ACCCTTGTTT CTACT
 M
 (SEQ ID NO:5)
 35 AGCAAAAGCA GGTAGATATT GAAAGATGAG TCTTCTAACC
 GAGGTCGAAA
 CGTACGTACT CTCTATCATC CCGTCAGGCC CCCTCAAAGC
 40 CGAGATCGCA
 CAGAGACTTG AAGATGTCTT TGCAGGGAAG AACACCGATC
 TTGAGGTTCT
 45 CATGGAATGG CTAAGACAA GACCAATCCT GTCACCTCTG
 ACTAAGGGGA
 TTTTAGGATT TGTGTTACG CTCACCGTGC CCAGTGAGCG
 AGGACTGCAG
 50 CGTAGACGCT TTGTCCAAA TGCCCTTAAT GGAACGGGG
 ATCCAAATAA
 CATGGACAAA GCAGTTAAAC TGTATAGGAA GCTCAAGAGG
 55 GAGATAACAT
 TCCATGGGGC CAAAGAAATC TCACTCAGTT ATTCTGCTGG
 TGCACTTGCC
 60 AGTTGTATGG GCCTCATATA CAACAGGATG GGGGCTGTGA
 CCACTGAAGT
 GGCATTGGC CTGGTATGTG CAACCTGTGA ACAGATTGCT
 65 GACTCCAGC

29

-continued

ATCGGTCTCA TAGGCAAATG GTGACAACAA CCAATCCACT
 AATCAGACAT
 GAGAACAGAA TGGTTTTAGC CAGCACTACA GCTAAGGCTA
 TGGAGCAAAT
 GGCTGGATCG AGTGAGCAAG CAGCAGAGGC CATGGAGGTT
 GCTAGTCAGG
 CTAGACAAAT GGTGCAAGCG ATGAGAACCA TTGGGACTCA
 TCCTAGCTCC
 AGTGCTGGTC TGAAAAATGA TCTTCTTGAA AATTTGCAGG
 CCTATCAGAA
 ACGAATGGG GTGCAGATGC AACGTTCAA GTGATCCTCT
 CACTATTGCC
 GCAAATATCA TTGGGATCTT GCACTTGACA TTGTGGATTC
 TTGATCGTCT
 TTTTTTCAAA TGCATTTACC GTCGCTTTAA ATACGGACTG
 AAAGGAGGGC
 CTTCTACGGA AGGAGTGCCA AAGTCTATGA GGAAGAATA
 TCGAAAGGAA
 CAGCAGAGTG CTGTGGATGC TGACGATGGT CATTTTGTCA
 GCATAGAGCT
 GGAGTAAAAA ACTACCTTGT TTCTACT
 NS
 (SEQ ID NO: 6)
 AGCAAAGCA GGGTGACAAA AACATAATGG ATCCAAACAC
 TGTGTCAAGC
 TTTCAGGTAG ATTGCTTTCT TTGGCATGTC CGCAAACGAG
 TTGCAGACCA
 AGAACTAGGC GATGCCCCAT TCCTTGATCG GCTTCGCCGA
 GATCAGAAAT
 CCCTAAGAGG AAGGGGCAGT ACTCTCGTC TGGACATCAA
 GACAGCCACA
 CGTGCTGGAA AGCAGATAGT GGAGCGGATT CTGAAAGAAG
 AATCCGATGA
 GGCACTTAAA ATGACCATGG CCTCTGTACC TGCCTCGCGT
 TACCTAACTG
 ACATGACTCT TGAGGAAATG TCAAGGGACT GGTCCATGCT
 CATACCCAAG
 CAGAAAGTGG CAGGCCCTCT TTGTATCAGA ATGGACCAGG
 CGATCATGGA
 TAAGAATC ATACTGAAAG CGAACTTCAG TGTGATTTTT
 GACCGGCTGG
 GACTCTAAT ATTGCTAAG GCTTTCACCG AAGAGGGAGC

30

-continued

AATTGTTGGC
 5 GAAATTTTAC CATTGCCTTC TCTTCCAGGA CATACTGCTG
 AGGATGTCAA
 AAATGCAGTT GGAGTCCTCA TCGGAGGACT TGAATGGAAT
 GATAACACAG
 10 TTCGAGTCTC TGAACTCTA CAGAGATTCTG CTGGAGAAG
 CAGTAATGAG
 AATGGGAGAC CTCCACTCAC TCCAAAACAG AAACGAGAAA
 15 TGGCGGGAAC
 AATTAGGTCA GAAGTTTGAA GAAATAAGAT GGTTGATTGA
 AGAAGTGAGA
 20 CACAACTGA AGATAACAGA GAATAGTTTT GAGCAAATAA
 CATTTATGCA
 AGCCTTACAT CTATTGCTTG AAGTGGAGCA AGAGATAAGA
 25 ACTTCTCTCGT
 TTCAGCTTAT TTAGTACTAA AAAACACCCT TGTTCCTACT
 HA
 (SEQ ID NO: 7)
 30 AGCAAAAGCAGGGGAAAATAAAAACAACCAAATGAGGCAACCTACTGG
 TCCTGTTATGTGCACTTGCACTGCAAT
 GCAGACACAATATGTATAGGCTACCATGCGAACAATTCAACCGACACTGT
 35 TGACACAGTACTCGAGAAGAATGTGACAGT
 GACACACTCTGTTAACCTGCTCGAAGACAGCCACAACGAAAACTATGTA
 GATTAAGGAATAGCCCCACTACAATTGG
 40 GGAAATGTAACATCGCCGGATGGCTCTTGGGAAACCCAGAATGCGACCCA
 CTGCTTCCAGTGAGATCATGGTCCTACATT
 GTAGAAACACCAAACCTCTGAGAATGGAATATGTTATCCAGGAGATTTTCA
 CGACTATGAGGAGCTGAGGGAGCAATTGAG
 45 CTCAGTGTATCATTCGAAAGATTCGAAATATTTCCCAAAGAAAGCTCAT
 GGCCCAACCACAAACAAACGAGTAACGG
 CAGCATGCTCCCATGAGGGGAAAAGCAGTTTTTACAGAAATTTGCTATGG
 50 CTGACGGAGAAGGAGGGCTCATACCCAAAG
 CTGAAAAATTTCTTATGTGAACAAAAAGGGAAGAAGTCCTTGTACTGTG
 GGGTATTTCATCCCCGCTAACAGTAAGGA
 55 ACAACAGAATCTCTATCAGAATGAAATGCTTATGTCTGTAGTACTT
 CAAATTATAACAGGAGATTTACCCCGGAAA
 TAGCAGAAAGACCCAAAGTAAGAGATCAAGCTGGGAGGATGAACATTAC
 60 TGGACCTTGCTAAAACCCGGAGACACAATA
 ATATTTGAGGCAAATGGAAATCTAATAGCACCAATGTATGCTTTTCGCACT
 GAGTAGAGGCTTTGGGTCCGGCATCATCAC
 65 CTCAAACGCATCAATGCATGAGTGAACACGAAGTGTCAAACACCCCTGG

31

-continued

GAGCTATAAACAGCAGTCTCCCTTACCAGA
 ATATACACCCAGTCACAATAGGAGAGTGCCTCAAAATACGTGAGGAGTGCC
 AAATTGAGGATGGTTACAGGACTAAGGAAC
 ATTCCGTCCATTCAATCCAGAGGTCTATTTGGAGCCATTGCCGGTTTTAT
 TGAAGGGGATGGACTGGAATGATAGATGG
 ATGGTATGGTTATCATCATCAGAATGAACAGGATCAGGCTATGCAGCGG
 ATCAAAAAGCACACAAAATGCCATTAAACG
 GGATTACAAACAAGGTGAACACTGTTCGAGAAAATGAACATTCAATTC
 ACAGCTGTGGTAAAGAATTCAACAAATTA
 GAAAAAAGGATGGAATTTAAATAAAAAAGTTGATGATGGATTTCTGGA
 CATTTGGACATATAATGCAGAATTGTTAGT
 TCTACTGGAATAAGGACTCTGGATTCCATGACTCAAATGTGAAGA
 ATCTGTATGAGAAAGTAAAAAGCCAATTAA
 AGAATAATGCCAAAGAAATCGAAATGGATGTTTTGAGTTCTACCAAG
 TGTGACAATGAATGCATGGAAAGTGAAGA
 AATGGGACTTATGATTATCCCAAATATTGAGAAGAGTCAAAGTTGAACAG
 GGAAAAGGTAGATGGAGTGAATTTGGAATC
 AATGGGATCTATCAGATTCTGGCGATCTACTCAACTGTGCGCAGTTTAC
 TGGTGCTTTTGGTCTCCCTGGGGGCAATCA
 GTTTCTGGATGTGTTCTAATGGATCTTTGCAGTGCAGAATATGCATCTGA
 GATTAGAATTTACAGATATGAGGAAAAAC
 ACCCTTGTTTCTACT
 NA
 (SEQ ID NO:8)
 AGCAAAGCAGGGTTTAAATGAATCCAAATCAGAAAATAATAACCAATT
 GGATCAATCTGTCTGGTAGTCGGACTAATT
 AGCCTAATATTGCAATAGGGAATATAATCTCAATATGGATTAGCCATTC
 AATTCAAACCTGGAAGTCAAACCACTACTGG
 AATATGCAACCAACATCATTACCTATAAAAAATAGCACCTGGGTAAAGG
 ACACAACCTCAGTGATATTAACCGGCAATT
 CATCTCTTTGTCCATCCGTGGGTGGGCTATATACAGCAAAGACAATAGC
 ATAAGAATTGGTTCCAAAGGAGACGTTTTTT
 GTCATAAGAGAGCCCTTTATTTTCATGTTCTCACTTGAATGCAGGACCTT
 TTTTCTGACCCAAGTGCTTACTGAATGA
 CAAGCATTCAAGTGGGACTGTTAAGGACAGAAGCCCTTATAGGGCCTTAA
 TGAGCTGCCCTGTGCGTGAAGCTCCGTCCC
 CGTACAATTCAAGATTGAATCGGTTGCTTGGTCAGCAAGTGCATGTCAT
 GATGGCATGGGCTGGCTAACAAATCGGAATT
 TCAGGTCCAGATAATGGAGCAGTGGCTGTATTAAATACAACGGCATAAT
 AACTGAAACCATAAAAAGTTGGAGGAAGAA
 AATATTGAGGACACAAGAGTCTGAATGTGCCTGTGTAATGGTTCATGTT
 TTACTATAATGACTGATGGCCCGAGTATG

32

-continued

GGCTGGCCTCGTACAAAATTTTCAAGATCGAAAAGGGGAAGTTACTAAA
 TCAATAGAGTTGAATGCACCTAATTCTCAC
 5 TATGAGGAATGTTCTGTTCACCTGATACCGGCAAAGTATGTGTGTG
 CAGAGACAATTGGCATGGTTCGAACCGGCC
 ATGGGTGTCTTTTCGATCAAAACCTGGATTATCAAATAGGATACATCTGCA
 10 GTGGGGTTTTCGGTGACAACCCGCGTCCC
 AAGATGGAACAGGCAGCTGTGGTCCAGTGTATGTTGATGGAGCAAACGGA
 GTAAAGGGATTTTCATATAGGTATGGTAAT
 15 GGTGTTTGGATAGGAAGGACCAAAGTCACAGTTCAGACATGGGTTTGA
 GATGATTTGGGATCCTAATGGATGGACAGA
 GACTGATAGTAAGTTCTCTGTGAGGCAAGATGTTGTGGCAATGACTGATT
 20 GGTCAGGGTATAGCGGAAGTTTCGTTCAAC
 ATCCTGAGCTGACAGGGCTAGACTGTATGAGGCCGTGCTTCTGGGTTGAA
 TTAATCAGGGGACGACCTAAAGAAAAACA
 25 ATCTGGACTAGTGCAGCAGCATTCTCTTTTGTGGCGTGAATAGTGATAC
 TGATGATTGGTCTTGGCCAGACGGTGCTGA
 GTTGCCATTGAGCATTGACAAGTAGTCTGTTCAAAAACCTCCTGTTTCT
 30 ACT

EXAMPLE 2

35 Influenza virus A/Hong Kong/213/2003 (H5N1, HK213) replicates systemically in chickens, causing lethal infection. Furthermore, this virus is lethal to chicken embryos. Thus, although its surface proteins are highly related to the currently circulating pathogenic avian influenza viruses, HK213 cannot be used as a vaccine strain as attempts to grow it in
 40 embryonated chicken eggs result in the production of poor-quality allantoic fluid. Additionally, the use of this highly virulent virus in the production of vaccines is unsafe for vaccine workers. To test the feasibility of using AIPR/8/34 as a master vaccine strain, the cleavage site of the hemagglutinin (HA) gene of HK213 (containing multiple basic amino acids) was mutated from a virulent to an avirulent phenotype (from RERRRKKR (SEQ ID NO:29) to - - - TETR (SEQ ID NO:30)). A virus containing the mutated HA gene produced
 45 non-lethal, localized infection in chickens. Additionally, the mutated virus was non-lethal to chicken embryos. Thus, growth of the mutated virus in embryonated eggs yielded high-quality allantoic fluid, and in this attenuated form, the virus is safe for vaccine producers.

55 A recombinant virus containing the neuraminidase (NA) and mutated HA genes from HK213, and all the remaining genes from high-titer A/PR/8/34 (H1N1, HG-PR8) virus (Example 1), which grows 10 times better than other A/PR/8/34 PR8 strains in eggs (10^{10} EID₅₀/ml; HA titer: 1:8,000), was generated in embryonated chicken eggs. This recombinant virus, which expresses surface proteins related to the currently circulating pathogenic avian influenza virus, grew to high titers in embryonated chicken eggs (FIG. 1). Thus, replacement of the HA and NA genes of HG-PR8 with those
 65 of a currently circulating strain of influenza virus resulted in a vaccine strain that can be safely produced, and demonstrates the use of PR8-HG as a master vaccine strain.

In Hong Kong in 1997, a highly pathogenic H5N1 avian influenza virus was transmitted directly from birds to humans, causing 18 confirmed infections and 6 deaths (Subbarao et al., 1998; Claas et al., 1998). In 2004-6, the geographic distribution of H5N1 viruses expanded in Asia, spreading to several adjacent European countries and to Africa. Altogether, 96 people infected with the virus have died in Vietnam, Thailand, Cambodia, Indonesia, China, Turkey, and Iraq (Li et al., 2004; WHO). These fatal outbreaks and the continued threat of a pandemic have led to the development of H5N1 virus vaccines for use in humans. However, because pathogenic H5N1 viruses grow poorly in embryonated chicken eggs and pose serious biosafety concerns for vaccine producers, reverse genetics has been used to generate vaccine candidates (Subbarao et al., 2003; Webby et al., 2004; Stephanson et al., 2004; Wood & Robertson, 2004).

Recombinant (6:2 reassortant) viruses that possess modified avirulent-type hemagglutinin (HA) and neuraminidase (NA) genes, both derived from a pathogenic H5N1 strain, with all remaining genes from a donor virus that grows well in eggs, are among the candidates to be produced by this method. The World Health Organization (WHO) recommends A/Puerto Rico/8/34 (H1N1; PR8) as a donor virus, because of its safety in humans and vigorous growth in eggs (Wood & Robertson, 2004; Webby & Webster, 2003). Recently, it was shown that such recombinant viruses grow less well in eggs than does the wild-type PR8 strain, even though they possess the same PR8 "internal" genes (i.e., those other than the HA and NA) (Horimoto et al., 2006).

Since vigorous growth in eggs is an essential property of vaccine seed viruses used in the production of inactivated vaccines, as described below, H5N1 vaccine candidates were generated that grow as well as the PR8 donor strain in eggs. First, the molecular basis for the high growth of PR8 in eggs was determined by defining the genes responsible for this property using reassortment analysis between PR8 and a WSN strain that grows poorly in eggs. It was found that HA-NA balance and PB1 function are important growth determinants. With this knowledge, a series of H5N1 viruses was produced with altered HA-NA combinations, with the PR8 background, to assess their growth in eggs against more conventional 6:2 reassortants, including the WHO-recommended NIBRG-14 virus.

Methods

Cells and Viruses

293T human embryonic kidney cells were maintained in Dulbecco's modified Eagle's minimal essential medium (DMEM) with 10% fetal calf serum and antibiotics. Madin-Darby canine kidney (MDCK) cells were grown in MEM with 5% newborn calf serum and antibiotics. African green monkey Vero WCB cells, which had been established after biosafety tests for use in human vaccine production (Sugawara et al., 2002), were maintained in serum-free VP-SFM medium (GIBCO-BRL) with antibiotics. Cells were maintained at 37° C. in 5% CO₂. The A/Vietnam/1194/2004 and A/Vietnam/1203/2004 (H5N1; VN1194 and VN1203) strains, isolated from humans, were propagated in 10-day-old embryonated chicken eggs for 2 days at 37° C., after which time the allantoic fluids containing virus were harvested and used for further experiments. All experiments with these viruses were carried out in a Biosafety Level 3 containment laboratory. The WHO-recommended vaccine seed virus, NIBRG-14 (VN1194/PR8 6:2 reassortant virus), was kindly gifted by Drs. John Wood and Jim Robertson at the National Institute for Biological Standards and Control, UK.

Construction of Plasmids and Reverse Genetics

To generate reassortants of influenza A viruses, a plasmid-based reverse genetics (Neumann et al., 1999) was used. Viral RNA from VN1194 or VN1203 was extracted from allantoic fluid by using a commercial kit (ISOGEN LS, Nippon Gene) and was converted to cDNA by using reverse transcriptase (SuperScript III; GIBCO-BRL) and primers containing the consensus sequences of the 3' ends of the RNA segments for the H5 viruses. The full-length cDNAs were then PCR-amplified with ProofStart polymerase (QIAGEN) and H5 subtype-specific primer pairs, and cloned into a plasmid under control of the human polymerase I promoter and the mouse RNA polymerase I terminator (Poll plasmids), generating a Poll-VN1194/HA or a Poll-VN1203/HA construct containing the VN1194 or VN1203 HA gene, respectively. By inverse PCR using back-to-back primer pairs, followed by ligation, the HA cleavage site sequence of the wild-type VN1194 or VN1203 (RERRRKKR; SEQ ID NO:29) virus was altered to create the avirulent-type sequence (RETR; SEQ ID NO:31) as described in Horimoto et al. (2006), the disclosure of which is incorporated by reference herein. A Poll-VN1203NA containing the VN1203 NA gene was constructed by the RT-PCR procedure (described above) with N1-specific primers. A series of pPoll NA mutant plasmids were prepared by inverse PCR. Using the Poll-VN1203NA as a template, pPoll-NAfill was constructed, which encodes a mutant NA containing a 20-amino acid (aa) (CNQSIITY-ENNTWVNQTYVN; SEQ ID NO:32) insertion derived from A/goose/Guangdong/1/96 (H5N1; GsGd96) NA into the NA stalk between 48-Pro and 49-Ile. pPoll-NAfill.N2 and -NAfill.N2N9, in which N2 (12 aa) or N2+N9 (12+12 aa) sequences derived from the stalk region of each NA subtype were inserted into the NA stalk between 42-Asn and 43-Gln, were constructed as described in Castrucci et al. (1993). All of these constructs were sequenced to ensure the absence of unwanted mutations.

A previously produced series of Poll constructs, derived from A/WSN/33 (H5N1; WSN) and PR8 strains was used, for reverse genetics (Horimoto et al., 2006; Neumann et al., 1999). Additionally, Poll constructs containing NA genes derived from A/Hong Kong/213/03 (H5N1; HK213), and A/Kanagawa/173/2001 (H1N1; Kanagawa) were used in this study (Horimoto et al., 2006; Kobasa et al., 2004; Peiris et al., 2004).

Plasmids expressing WSN or PR8 NP, PA, PB1, or PB2 under control of the chicken β -actin promoter were used for all reverse genetics experiments (Horimoto et al., 2006; Neumann et al., 1999). Briefly, Poll plasmids and protein expression plasmids were mixed with a transfection reagent, Trans-IT 293T (Panvera), incubated at room temperature for 15 min, and then added to 293T cells. Transfected cells were incubated in Opti-MEM 1 (GIBCO-BRL) for 48 hours. For reverse genetics in Vero WCB cells, an electroporator (Amaxa) was used to transfect the plasmid mixtures according to the manufacturer's instructions. Sixteen hours after transfection, freshly prepared Vero WCB cells were added onto the transfected cells and TPCK-trypsin (1 μ g/ml) was added to the culture 6 hours later. Transfected cells were incubated in serum-free VP-SFM for a total of 4 days. Supernatants containing infectious viruses were harvested, biologically cloned by limiting dilution in embryonated eggs, and used in further experiments.

Properties of Viral Replication in Eggs

Virus was inoculated into the allantoic cavity of 10-day-old embryonated chicken eggs, and incubated at 37° C. for 48 hours. Virus in the allantoic fluids was then titrated by HA assay using either 0.5% chicken erythrocytes or 0.8% guinea

pig erythrocytes or in eggs to determine the median egg infectious dose (EID₅₀)/ml of virus. For some viruses, plaque titration was conducted with MDCK cells and TPCK-trypsin (1 µg/ml). The growth kinetics of some viruses was assessed in eggs after inoculating 10⁴ EID₅₀ of virus.

Virus Elution Assay from Chicken Erythrocytes

Fifty µl of twofold dilutions of virus containing the HA titers of 1:1024 were incubated with 50 µl of 0.5% chicken erythrocytes in a microtiter plate at 4° C. for 1 hour. The plate was then stored at 37° C., and the reduction of HA titers was recorded periodically. Phosphate-buffered saline with 6.8 mM CaCl₂ was used as a diluent.

Results

Molecular Basis for the High Growth Property of PR8 in Chicken Eggs

Although PR8 is recommended by WHO for use as a donor virus to generate reverse genetics-based H5 influenza vaccine, the molecular basis of its high growth property is not fully understood. The M gene was said to be responsible for the vigorous growth of PR8 in eggs (Subbarao et al., 2003), but this claim is apparently not found in the published original data (Kilbourne et al., 1969). Thus, a reassortment analysis was conducted using a WSN strain that grows poorly in eggs. Table 3 shows the compatibility between the HAs and NAs of PR8 versus the WSN strain in terms of viral growth in embryonated chicken eggs. All reassortant test viruses grew better than the wild-type WSN, but less well than the egg-adapted PR8, demonstrating that both surface glycoproteins and internal proteins are responsible for the high growth property of PR8.

TABLE 3

Compatibility between the HAs and NAs of PR8 versus WSN strains, assessed by viral growth in chicken embryonated eggs				
Gene constellation of reassortant			HA titer ^{b)}	
HA	NA	6 others ^{a)}	Chicken RBC	Guinea pig RBC
WSN	WSN	WSN	16/8	32/8
PR8	WSN	WSN	64/32	64/32
WSN	PR8	WSN	16/16	32/16
PR8	PR8	WSN	128/128	128/128
WSN	WSN	PR8	64/64	64/64
PR8	WSN	PR8	64/128	64/128
WSN	PR8	PR8	512/512	512/512
PR8	PR8	PR8	2048/2048	2048/2048

^{a)}Genes encoding the internal proteins PB1, PB2, PA, NP, M, and NS.

^{b)}Growth of each reassortant virus in chicken eggs, assessed in HA assays with 0.5% chicken RBC and 0.8% guinea pig RBC. HA titers from two independent experiments are shown.

Since the growth of a reassortant virus containing both of the PR8 glycoproteins and all six internal proteins derived from WSN was drastically reduced in eggs, as compared with that of PR8 (Tables 3 and 4), a series of reassortant viruses was produced to define the internal proteins responsible for this property. A single-gene reassortant virus containing the WSN PB1 and all remaining genes from PR8 grew poorly, at a level similar to that of a reassortant containing all of the WSN genes encoding internal proteins, whereas a reassortant containing the PR8 PB1 and WSN genes encoding all remaining internal proteins replicated to a high titer (Table 4). Thus, the PR8 PB1 likely possesses the optimal polymerase activity for viral genome replication in eggs, in contrast to a previous report implicating the M segment in this role (Subbarao et al., 2003). Table 4 Compatibility among genes encoding internal proteins of PR8 and WSN viruses, assessed by viral growth in chicken embryonated eggs

Gene constellation of reassortant ^{a)}									
HA	NA	PB2	PB1	PA	NP	M	NS	HA titer ^{b)}	
5	PR8	PR8	PR8	PR8	PR8	PR8	PR8	2048/2048/	1024
10	PR8	PR8	PR8	PR8	PR8	PR8	WSN	1024/1024/	1024
	PR8	PR8	PR8	PR8	PR8	WSN	PR8	2048/1024/	1024
15	PR8	PR8	PR8	PR8	PR8	WSN	WSN	1024/1024/512	1024/1024/512
	PR8	PR8	PR8	WSN	PR8	PR8	PR8	1024/512/256	128/64/64
20	PR8	PR8	WSN	PR8	PR8	PR8	PR8	1024/1024/	1024
	PR8	PR8	WSN	WSN	WSN	PR8	PR8	64/64/32	128/64/64
25	PR8	PR8	WSN	WSN	WSN	WSN	WSN	1024/512/512	

^{a)}Both the HA and NA genes were derived from PR8 in all reassortant viruses, while some of the genes encoding internal proteins were from the WSN strain.

^{b)}Growth rate of each reassortant virus in chicken eggs was assessed with HA assays in 0.5% chicken RBC. HA titers, obtained in three independent experiments, are shown.

Generation of H5N1 Vaccine Seed Candidates with Enhanced Growth Ability in Chicken Eggs

In an earlier study, the growth of WSN in eggs was shown to be enhanced by lengthening the NA stalk to increase NA function: the longer the stalk, the better the replication of the virus (Castrucci et al., 1993). This finding prompted the production of a series of H5N1 viruses comprising mutated or heterologous N1s with the PR8 background and compare their growth in eggs. The A/Vietnam/1203/2004 (H5N1; VN1203) NA contains a 20-amino acid (20-aa) deletion in its stalk region (hence, 24 aa in the stalk). Therefore, a mutant NA, VN1203fill, was constructed containing a 44-aa stalk like the H5N1 precursor virus A/goose/Guangdong/1/96 (H5N1) (Xu et al., 1999), as well as other NA mutants, VN1202fill.N2 and VN1203fill.N2N9 that contained longer stalks, 58- and 72-aa, respectively (FIG. 2). The heterologous N1 from A/Hong Kong/213/03 (H5N1; HK213) containing 44-aa in the stalk was also examined. The NAs from H1N1 strains such as PR8, A/Kanagawa/173/2001 (H1N1; Kanagawa), and WSN, all of which possess 24-aa in the stalk, were also tested. Using these NA constructs, a total of eight reassortant viruses was generated, seven 6:2 and one 7:1 with the modified avirulent-type VN1203 HA and PR8 background (Table 5). Another series of reassortant viruses was constructed with the modified avirulent-type A/Vietnam/1194/2004 (H5N1; VN1194) HA. By comparison with constructs containing the parental VN1203 NA, only the 7:1 reassortant virus and a 6:2 reassortant containing a combination of the modified VN1194 HA and VN1203fill NA, showed enhanced growth in eggs.

TABLE 5

Viral titers of H5N1/PR8 reassortant viruses in chicken embryonated eggs ^{a)}										
HA titer/Infectivity titer (log ₂ /log ₁₀ EID ₅₀ /ml)										
HA ^{b)}	Experiment ^{c)}	NA derived from ^{d)}								Wild-type PR8 ^{e)}
		VN1203	VN1203fill	VN1203fill.N2	VN1203fill.N2N9	HK213	PR8	Kanagawa	WSN	
VN1203	1	9.2 ± 0.4/	9.6 ± 0.5/	9.2 ± 0.5/	9.0 ± 0.0/	9.6 ± 0.5/	9.6 ± 0.5/	9.8 ± 0.4/	<1.0/	10.7 ± 0.6/
		8.9 ± 0.3 ^{f)}	8.8 ± 0.6	8.9 ± 0.4	8.8 ± 0.5	8.8 ± 0.1	9.5 ± 0.4	9.4 ± 0.2	ND	10.3 ± 0.4
	2	9.0 ± 1.0/	9.0 ± 0.0/	8.3 ± 0.6/	8.0 ± 0.0/	8.7 ± 0.6/	9.7 ± 0.6/	ND/ND	ND/ND	11.0 ± 0.0/
VN1194	1	9.4 ± 0.2	9.7 ± 0.2	8.6 ± 0.2	ND	8.5 ± 0.3	10.1 ± 0.2	9.0 ± 0.0/	<1.0/	10.3 ± 0.4
		8.7 ± 0.6/	9.3 ± 0.6/	9.3 ± 0.6/	9.0 ± 0.0/	ND/ND	9.3 ± 0.6/	9.0 ± 0.0/	<1.0/	10.7 ± 0.6/
	2	8.7 ± 0.2	9.3 ± 0.2	9.2 ± 0.2	8.6 ± 0.2		9.5 ± 0.3	8.8 ± 0.9	5.2 ± 0.2	10.1 ± 0.2

^{a)}Eggs (10-day-old) were inoculated with virus (10⁴EID₅₀), and incubated for 48 hours at 37° C.; viral titers in allantoic fluids were determined.

^{b)}Two H5 HA genes (VN1203 and VN1194) were used to generate reassortant viruses with a PR8 background. The HA cleavage sites of both VN1203 and VN1194 were modified to that of the avirulent-type H5 HA.

^{c)}Two independent experiments, each using 3 to 5 eggs, were performed for VN1203 constructs, while a single experiment was done for VN1194.

^{d)}A total of eight NA genes were used to generate reassortant viruses; three insertion mutant NAs (VN1203fill, VN1203fill.N2, and VN1203fill.N2N9) were prepared to assess the influence of NA stalk length on virus growth in eggs by comparison to parental VN1203 NA; the other NAs were derived from an H5N1 human isolate (HK213) or H1N1 viruses (PR8, Kanagawa, and WSN). Thus, all reassortant viruses except one containing PR8 NA (7:1 reassortant) are 6:2 reassortant viruses with a PR8 background.

^{e)}Growth of wild-type PR8 was also assessed as a control for each experiment.

^{f)}Growth of each reassortant virus in eggs was assessed by either HA or infectivity assay, and reported as mean ± s.d. of HA titer (log₂)/mean ± s.d. of infectivity titer (log₁₀EID₅₀/ml). Significantly enhanced HA and infectivity titers (p < 0.05, t-test), by comparison to those of standard viruses containing VN1203 HA and VN1203 NA or VN1194 HA and VN1203 NA, are shown in boldface type.

ND, not determined.

Further testing of selected reassortant viruses by a plaque assay of the stock viruses demonstrated a greater than 3-fold higher titer (p=0.003, Student t-test) for the reassortant virus containing PR8 NA compared with the virus containing parental VN1203 NA, although it did not grow as well as egg-adapted PR8 (FIG. 3). Assessment of the growth kinetics of reassortant viruses with the PR8, VN1203fill or VN1203 NA in eggs revealed a superior growth rate for the virus with PR8 NA (7:1 reassortant) (FIG. 4).

To determine the molecular basis of the high growth property observed in the 7:1 reassortant virus, the NA function of reassortant viruses was tested by an assay evaluating virus elution from chicken erythrocytes (FIG. 5). Reassortant viruses containing PR8 or VN1203fill NA were eluted from erythrocytes more rapidly than those with the parental VN1203 NA, indicating greater NA activity for PR8 or VN1203fill.NA. These results support the idea that high NA function enhances viral growth in eggs (Castrucci et al., 1993).

Growth Comparison of H5N1 Vaccine Seed Candidates Produced in this Study with the WHO-Recommended Vaccine Seed Virus, NIBRG-14, in Eggs

To validate the potential of candidate seed viruses in the production of H5N1 vaccines, their infectivity titers were compared with that of the WHO-provided NIBRG-14 virus under the same experimental conditions. The 7:1 reassortant viruses containing either VN1194 or VN1203-derived HAs and all the other genes from our PR8 strain showed significantly higher titers (p<0.05, Student t-test) than the NIBRG-14 virus in eggs, as assessed by EID₅₀ (Table 6) and plaque titration (FIG. 6). Interestingly, even the 6:2 reassortant virus containing both its HA and NA from the VN1194 virus grew significantly better (about 7-fold, p=0.047) than NIBRG-14 (also a VN1194PR8 6:2 reassortant virus) by plaque titration (FIG. 5). This difference in the growth of two 6:2 reassortant viruses possessing identical VN1194 HAs and NAs indicates that the PR8 strain used in this study would be superior to the one used to generate NIBRG-14 for supporting high viral growth during vaccine production in eggs.

TABLE 6

25	Growth comparison of H5N1/PR8 reassortant viruses generated in this study with the WHO-recommended vaccine seed virus (NIBRG-14) ^{a)}					
	Infectivity titer (log ₁₀ EID ₅₀ /ml)					
	Hours	Reassortants made in this study ^{b)}				NIBRG-14
30	Post-infection	VN1194/ VN1194	VN1194/ PR8	VN1203/ VN1203	VN1203/ PR8	VN1194/ VN1194
	48	8.7 ± 0.4	9.4 ± 0.2	9.1 ± 0.2	9.5 ± 0.3	8.2 ± 0.3
	60	8.3 ± 0.5	8.9 ± 0.5	8.6 ± 0.4	9.2 ± 0.3	7.4 ± 0.2
25						

^{a)}Growth of reassortant viruses was assessed by inoculating eggs (n = 3) with each virus, harvesting allantoic fluid at the indicated times, and determining the EID₅₀. The data are shown as mean ± s.d. of infectivity titers (log₁₀EID₅₀/ml). Significantly enhanced infectivity titers (p < 0.05, t-test), by comparison with those of NIBRG-14, are shown in boldface type.

^{b)}Categorized by the derivation of the HA/NA. The HA cleavage site of both VN1203 and VN1194 were modified to that of the avirulent-type H5 HA.

40 Discussion

Recombinant viruses possessing modified avirulent-type HA and NA genes, both derived from an H5N1 human isolate, and all remaining genes from the PR8 strain (6:2 reassortant) have been produced and used as seed viruses for inactivated influenza vaccines now being tested in human clinical trials (Wood & Robertson, 2004). Seed strains used in this way must grow well in embryonated eggs. Although egg-adapted PR8 meets this requirement, some 6:2 reassortant viruses, despite containing six internal genes from PR8, do not grow well in eggs (Tables 3 and 5). Here it is demonstrated that the growth of egg-adapted PR8 in chicken eggs is affected by the functional balance of the HA and NA surface glycoproteins.

It is likely that low yields of some 6:2 reassortant viruses with a PR8 background and surface glycoproteins from highly pathogenic avian viruses may result not only from an HA-NA functional imbalance for growth in eggs but also from genetic (and/or functional) incompatibility between the avian surface glycoprotein genes and the internal genes from PR8. Here it is shown that among the internal genes of PR8, PB1 is very important for its high growth in eggs. This information suggests another strategy for reverse genetics-based H5N1 vaccine production; that is, the PB8 PB1 gene alone may be sufficient to generate vigorously growing reassortants for vaccine seed viruses. Thus, by using genes that encode non-PB1 internal proteins from strains other than PR8, one might avoid genetic incompatibility between avian and PR8

viruses. Studies to dissect the molecular basis for the high growth property of PR8 PB1 in eggs would be of considerable interest. One could, for example, analyze the structural and functional differences between the PB1s or PB1-F2s of PR8 and WSN (which differ by 18 and 10 amino acids, respectively; Chen et al., 2004).

The 7:1 reassortant viruses produced in this study replicated significantly better (more than 20-fold by plaque titration) than the WHO-recommended 6:2 reassortant virus NIBRG-14. Even the 6:2 reassortant that was identical to the NIBRG-14 except for the PR8 strain of origin replicated 7-fold better than the recommended virus. These findings suggest that the PR8 strain used in this study may be a superior donor virus for the production of reverse genetics-based pandemic vaccines.

One could argue that the 7:1 reassortant viruses would induce a loss of protective immune response due to antigenic differences in the NA proteins (even though both PR8 and the highly pathogenic viruses contain N1 NAs) (Murphy et al., 1972; Kilbourne et al., 1968; Chen et al., 2000). However, since the HA is the major protective antigen in inactivated vaccines, the higher growth property conferred by the PR8 NA would likely offset the limited antigenic mismatch in this minor protective antigen. In the event of a pandemic caused by a highly pathogenic avian influenza virus, chicken eggs will be in short supply. It is proposed that under such conditions, 7:1 reassortant-based vaccine seed viruses possessing an enhanced growth property in eggs would offer an attractive option for the generation of reverse genetics-based H5 vaccine viruses.

EXAMPLE 4

To identify the genes responsible for the high growth rate of an H5N1 vaccine seed virus in chicken embryonated eggs, the growth of reassortant H5N1 viruses possessing PR8(UW) or PR8(Cambridge) internal genes in chicken embryonated eggs was assessed (FIG. 7). The HA and NA genes of all of the reassortant viruses were derived from A/Vietnam/1194/2002. All other genes were derived from either PR8(UW) or PR8 (Cambridge), which also provided the non-HA and -NA genes of the NIBRG-14 vaccine strain. Higher titers were obtained when the majority of internal genes were from PR8 (UW).

The effect of the M and NS genes on the growth of viruses in chicken embryonated eggs is shown in FIG. 8. For PR8 (UW)/1194-CamM and PR8(UW)/1194-CamNS, the M and NS gene segments, respectively, were derived from PR8 (Cambridge), while the other internal segments came from PR8(UW). The HA and NA segments were derived from A/Vietnam/1194/2004. Highest titers were with the M gene segment of PR8(UW) and the NS gene of PR8 (Cambridge).

The results in FIGS. 7-8 show that the polymerase subunit (PA, PB1, and PB2) and NP genes of PR8(UW) enhanced the growth of an H5N1 vaccine seed virus in chicken embryonated eggs. Also, the NS gene of PR8(Cambridge) enhanced the growth of an H5N1 vaccine seed virus in chicken embryonated eggs.

To identify the gene and amino acid(s) responsible for the high growth rate of the H5N1 vaccine seed virus in MDCK cells, the growth of PR8(UW)/1194 and NIBRG-14 virus in MDCK cells was assessed. The data in FIG. 9 show that the growth of PR8(UW)/1194 was significantly better than that of NIBRG-14 in MDCK cells. Moreover, the PB2 segment of PR8(UW) enhanced the growth of the vaccine seed virus in MDCK cells (FIG. 10). The tyrosine residue at position 360 in

PB2 of PR8(UW) is likely responsible for the high growth rate of the vaccine seed virus in MDCK cells (FIG. 11).

To identify a combination of genes responsible for the high growth of an H5N1 vaccine seed virus in MDCK cells, the growth rates in MDCK cells of reassortants with different HA, NA, and NS genes was determined. NS from PR8(Cambridge) and NA with a long stalk (e.g., from A/Hong Kong/213/2003 or VN1203Fill) enhanced virus growth in MDCK cells (FIG. 12).

To determine which amino acids in NS are responsible for the high growth rate of the H5N1 vaccine seed virus in MDCK cells, the growth in MDCK cells of the H5N1 vaccine seed virus containing a heterologous NS segment was measured. An amino acid substitution from K [PR8(UW)NS] to E [PR8(Cambridge)] at position 55 of NS1 enhanced the growth of the H5N1 vaccine seed viruses in MDCK cells (FIG. 13).

FIG. 14 summarizes the genotype of an H5N1 seed virus with high growth capacity in chicken embryonated eggs or MDCK cells.

REFERENCES

- Avery's *Drug Treatment: Principles and Practice of Clinical Pharmacology and Therapeutics*, 3rd edition, ADIS Press, Ltd., Williams and Wilkins, Baltimore, Md. (1987).
- Aymard-Henry et al., *Virology: A Practical Approach*, Oxford IRL Press, Oxford, 119-150(1985).
- Bachmeyer, *Intervirology*, 5:260 (1975).
- Berkow et al., eds., *The Merck Manual*, 16th edition, Merck & Co., Rahway, N.J. (1992).
- Bridgen et al., *Proc. Natl. Acad. Sci. U.S.A.*, 93:15400 (1996).
- Castrucci & Kawaoka, *J. Virol.*, 67:759 (1993).
- Castrucci et al., *J. Virol.*, 69:2725 (1995).
- Chen et al., *Emerg. Infect. Dis.*, 10:630 (2004).
- Chen et al., *Vaccine*, 18:3214 (2000).
- Claas et al., *Lancet*, 351:472 (1998).
- Conzelmann et al., *J. Gen. Virol.*, 77:381 (1996).
- Conzelmann et al., *Trends Microbiol.*, 4:386 (1996).
- Conzelmann, *Annu. Rev. Genet.*, 32:123 (1998).
- Cozelmann et al., *J. Virol.*, 68:713 (1994).
- Edwards, *J. Infect. Dis.*, 169: 68 (1994).
- Enami et al., *Proc. Natl. Acad. Sci. U.S.A.*, 87:3802 (1990).
- Enami et al., *Virology*, 185:291 (1991).
- Fodor et al., *J. Virol.*, 73:9679 (1999).
- Grand and Skehel, *Nature, New Biology*, 238:145 (1972).
- Hatta et al., *Science*, 293:1840 (2001).
- Horimoto et al., *J. Virol.*, 68:3120 (1994).
- Horimoto et al., *Vaccine*, 24:3669 (2006).
- Keitel et al., in *Textbook of Influenza*, eds. Nickolson, K. G., Webster, R. G., and Hay, A. (Blackwell, Oxford), pp. 373-390 (1998).
- Kendal et al., *Infect. Immunity*, 29:966 (1980).
- Kilbourne et al., *J. Virol.*, 2:281 (1968).
- Kilbourne, *Bull. M2 World Health Org.*, 41: 653 (1969).
- Kilbourne, *Bull. World Health Org.*, 41:643 (1969).
- Kobasa et al., *Nature*, 431:703 (2004).
- Kovesdi et al., *J. Curr. Opin. Biotechnol.*, 8:583 (1997).
- Laver & Webster, *Virology*, 69:511 (1976).
- Lawson et al., *Proc. Natl. Acad. Sci. U.S.A.*, 92:4477 (1995).
- Li et al., *Nature*, 430:209 (2004).
- Marriott et al., *Adv. Virus Res.*, 53:321 (1999).
- Mizrahi, (ed.), *Viral Vaccines*, Wiley-Liss, New York, 39-67 (1990).
- Munoz et al., *Antiviral Res.*, 46:91 (2000).

Murphy et al., *New Engl. J. Med.*, 286:1329 (1972).
 Murphy, *Infect. Dis. Clin. Pract.*, 2: 174 (1993).
 Muster et al., *Proc. Natl. Acad. Sci. USA*, 88: 5177 (1991).
 Nagai et al., *Microbiol. Immunol.* 43:613 (1999).
 Nagai, *Rev. Med. Virol.*, 9:83 (1999).
 Neumann et al., *Adv. Virus Res.*, 53:265 (1999).
 Neumann et al., *J. Gen. Virol.*, 83:2635 (2002).
 Neumann et al., *J. Virol.*, 71:9690 (1997).
 Neumann et al., *Proc. Natl. Acad. Sci. USA*, 96:9345
 (1999).
 Neumann et al., *Virology*, 287:243 (2001).
 Ogra et al., *J. Infect. Dis.*, 134: 499 (1977).
 Osol (ed.), *Remington's Pharmaceutical Sciences*, Mack
 Publishing Co., Easton, Pa. 1324-1341 (1980).
 Parks et al., *J. Virol.*, 73:3560 (1999).
 Peiris et al., *Lancet*, 363:617 (2004).
 Pekosz et al., *Proc. Natl. Acad. Sci. U. S. A.*, 96:8804
 (1999).
 Radecke et al., *EMBO J.*, 14:5773 (1995).
 Roberts et al., *Virology*, 247:1 (1998).
 Robertson et al., *Biologicals*, 20:213 (1992).
 Robertson et al., *Giornale di Igiene e Medicina Preventiva*,
 29:4 (1988).

Rose, *Proc. Natl. Acad. Sci. U.S.A.*, 93:14998 (1996).
 Schnell et al., *EMBO J.*, 13:4195 (1994).
 Stephenson et al., *Lancet Inf Dis.*, 4:499 (2004).
 Subbarao et al., *J. Virol.*, 67:7223 (1993).
 Subbarao et al., *Science*, 279:393 (1998).
 Subbarao et al., *Virology*, 305:192 (2003).
 Sugawara et al., *Biologicals*, 30:303 (2002).
 Webby & Webster et al., *Science*, 302:1519 (2003).
 Webby et al., *Lancet*, 363:1099 (2004).
 Wood & Robertson, *Nat. Rev. Microbiol.*, 2:842 (2004).
 World Health Organization TSR No. 673 (1982).
 World Health Organization. Confirmed human cases of
 avian influenza A (H5N1). See the URL at [www.who.int/csr/](http://www.who.int/csr/disease/avian_influenza/country/en/index.html)
[disease/avian_influenza/country/en/index.html](http://www.who.int/csr/disease/avian_influenza/country/en/index.html).
 Xu et al., *Virology*, 261:15 (1999).
 All publications, patents and patent applications are incor-
 porated herein by reference. While in the foregoing specifi-
 cation this invention has been described in relation to certain
 preferred embodiments thereof, and many details have been
 set forth for purposes of illustration, it will be apparent to
 those skilled in the art that the invention is susceptible to
 additional embodiments and that certain of the details
 described herein may be varied considerably without depart-
 ing from the basic principles of the invention.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 38

<210> SEQ ID NO 1

<211> LENGTH: 2233

<212> TYPE: DNA

<213> ORGANISM: Influenza A

<400> SEQUENCE: 1

```

agcgaaagca ggtactgata caaaatggaa gattttgtgc gacaatgctt caatccgatg      60
attgtcagagc ttgcggaaaa aacaatgaaa gagtatgggg aggacctgaa aatcgaaaaa      120
aacaaatttg cagcaatatg cactcacttg gaagtatgct tcatgtattc agattttcac      180
ttcatcaatg agcaaggcga gtcaataatc gtagaacttg gtgatccaaa tgcacttttg      240
aagcacagat ttgaaataat cgagggaaga gatcgacaaa tggcctggag agtagtaaac      300
agtatttgca acactacagg ggctgagaaa ccaaagtctt taccagattt gtatgattac      360
aaggagaata gattcatcga aattggagta acaaggagag aagttcacat atactatctg      420
gaaaaggcca ataaaattaa atctgagaaa acacacatcc acattttctc gttcactggg      480
gaagaaatgg ccacaaaggc agactacact ctcatgaag aaagcagggc taggatcaaa      540
accagactat tcaccataag acaagaaatg gccagcagag gcctctggga ttcctttcgt      600
cagtcgaga gaggagaaga gacaattgaa gaaaggtttg aaatcacagg aacaatgcgc      660
aagcttgccg accaaagtct ccgcgcgaac ttctccagcc ttgaaaattt tagagcctat      720
gtggatggat tcgaaccgaa cggtacattt gagggcaagc tgtctcaaat gtccaaagaa      780
gtaaatgcta gaattgaacc ttttttgaaa acaacaccac gaccacttag acttccgaat      840
gggcctccct gttctcagcg gtccaaattc ctgctgatgg atgcctaaa attaagcatt      900
gaggacccaa gtcataagg agagggaata ccgctatatg atgcaatcaa atgcatgaga      960
acattctttg gatggaagga acccaatgtt gttaaaccac acgaaaaggg aataaatcca      1020
aattatcttc tgtcatggaa gcaagtactg gcagaactgc aggacattga gaatgaggag      1080
aaaattccaa agactaaaaa tatgaagaaa acaagtcagc taaagtgggc acttggtgag      1140

```

-continued

aacatggcac	cagaaaaggt	agactttgac	gactgtaaag	atgtagggtga	tttgaagcaa	1200
tatgatagtg	atgaaccaga	attgaggtcg	cttgcaagtt	ggattcagaa	tgagtttaac	1260
aaggcatgcg	aactgacaga	ttcaagctgg	atagagctcg	atgagattgg	agaagatgtg	1320
gctccaattg	aacacattgc	aagcatgaga	aggaattatt	tcacatcaga	ggtgtctcac	1380
tgagagacca	cagaatacat	aatgaaggga	gtgtacatca	atactgcctt	gcttaatgca	1440
tcttgtgcag	caatggatga	tttccaatta	attccaatga	taagcaagtg	tagaactaag	1500
gaggggaaggc	gaaagaccaa	cttgtatggt	ttcatcataa	aaggaagatc	ccacttaagg	1560
aatgacaccg	acgtggtaaa	cttgttgagc	atggagtttt	ctctcactga	ccaagactt	1620
gaaccacata	aatgggagaa	gtactgtggt	cttgagatag	gagatatgct	tataagaagt	1680
gccataggcc	aggtttcaag	gcccattgtc	ttgtatgtga	gaacaaatgg	aacctcaaaa	1740
attaaaatga	aatggggaat	ggagatgagg	cgttgccctc	tccagtcact	tcaacaaatt	1800
gagagtatga	ttgaagctga	gtcctctgtc	aaagagaaag	acatgaccaa	agagttcttt	1860
gagaacaaat	cagaacatg	gcccattgga	gagtccecca	aaggagtggg	ggaaagtctc	1920
attgggaagg	tctgcaggac	tttattagca	aagtcggtat	tcaacagctt	gtatgcatct	1980
ccacaactag	aaggattttc	agctgaatca	agaaaactgc	ttcttatcgt	tcaggctctt	2040
agggacaacc	tggaaacctg	gacctttgat	cttggggggc	tatatgaagc	aattgaggag	2100
tgccatgatta	atgatccctg	ggttttgctt	aatgcttctt	ggttcaactc	cttccttaca	2160
catgcattga	gttagttgtg	gcagtgtctac	tatttgctat	ccatactgtc	caaaaaagta	2220
ccttgtttct	act					2233

<210> SEQ ID NO 2

<211> LENGTH: 2341

<212> TYPE: DNA

<213> ORGANISM: Influenza A

<400> SEQUENCE: 2

agcgaaagca	ggcaaacat	ttgaatggat	gtcaatccga	ccttactttt	cttaaaagt	60
ccagcacaaa	atgctataag	cacaactttc	ccttatactg	gagaccctcc	ttacagccat	120
gggacaggaa	caggatacac	catggatact	gtcaacagga	cacatcagta	ctcagaaaag	180
ggaagatgga	caacaacac	cgaaactgga	gcaccgcaac	tcaaccgat	tgatgggcca	240
ctgccagaag	acaatgaacc	aagtggttat	gcccacacag	atttgtgtatt	ggaggcgatg	300
gctttccttg	aggaatccca	tcttggtatt	tttgaaaact	cgtgtattga	aacgatggag	360
gttggttcagc	aaacacgagt	agacaagctg	acacaaggcc	gacagacctg	tgactggact	420
ctaaatagaa	accaacctgc	tgcaacagca	ttggccaaca	caatagaagt	gttcagatca	480
aatggcctca	cggccaatga	gtctggaagg	ctcatagact	tccttaagga	tgtaatggag	540
tcaatgaaca	aagaagaaat	ggggatcaca	actcattttc	agagaaagag	acgggtgaga	600
gacaatatga	ctaagaaaat	gataacacag	agaacaatgg	gtaaaaagaa	gcagagattg	660
aacaaaagga	gttatcta	tagagcattg	accctgaaca	caatgaccaa	agatgctgag	720
agaggggaagc	taaaacggag	agcaattgca	acccagggga	tgcaataaag	ggggtttgta	780
tactttgttg	agacactggc	aaggagtata	tgtgagaaac	ttgaacaatc	agggttgcca	840
gttgagggca	atgagaagaa	agcaaagtgtg	gcaaatgttg	taaggaagat	gatgaccaat	900
tctcaggaca	ccgaactttc	tttcaccatc	actggagata	acaccaaagt	gaacgaaaat	960
cagaatcctc	ggatgttttt	ggccatgatc	acatatatga	ccagaaatca	gcccgaatgg	1020

-continued

```

ttcagaaatg ttctaagtat tgctccaata atgtttctcaa acaaaatggc gagactggga 1080
aaaggggtata tgtttgagag caagagtatg aaacttagaa ctcaaatacc tgcagaaatg 1140
ctagcaagca tcgatttgaa atatttcaat gattcaacaa gaaagaagat tgaaaaaatc 1200
cgaccgctct taatagaggg gactgcatca ttgagccctg gaatgatgat gggcatgttc 1260
aatatgttaa gcaactgtatt aggcgtctcc atcctgaatc ttggacaaaa gagatacacc 1320
aagactactt actggtggga tggctctcaa tcctctgacg attttgctct gattgtgaat 1380
gcaccaatc atgaagggat tcaagccgga gtcgacaggt tttatcgaa cgtgaagcta 1440
cttggaatca atatgagcaa gaaaaagtct tacataaaca gaacaggtag atttgaattc 1500
acaagttttt tctatcgta tgggtttgtt gccaatttca gcatggagct tcccagtttt 1560
ggggtgtctg ggatcaacga gtcagccgac atgagtattg gagttactgt catcaaaaac 1620
aatatgataa acaatgatct tgggtccagca acagctcaaa tggcccttca gttgttcac 1680
aaagattaca ggtacacgta ccgatgccat ataggtgaca cacaatatca aaccgaaga 1740
tcatttgaaa taaagaaact gtgggagcaa acccgttcca aagctggact gctggtctcc 1800
gacggaggcc caaatttata caacattaga aatctccaca ttctgaagt ctgcctaaaa 1860
tgggaattga tggatgagga ttaccagggg cgtttatgca acccactgaa cccatttgtc 1920
agccataaag aaattgaatc aatgaacaat gcagtgatga tgccagcaca tgggtccagcc 1980
aaaaacatgg agtatgatgc tgttgcaaca acacactcct ggatcccaa aagaaatcga 2040
tccatcttga atacaagtca aagaggagta cttgaggatg acaaatgta ccaaaggatg 2100
tgcaatttat ttgaaaaatt cttccccagc agttcataca gaagaccagt cgggatatcc 2160
agtatggtgg aggctatggt ttccagagcc cgaattgatg cacggattga tttcgaatct 2220
ggaaggataa agaaagaaga gttcactgag atcatgaaga tctgttccac cattgaagag 2280
ctcagacggc aaaaatagtg aatttagctt gtccttcacg aaaaaatgac ttgtttctac 2340
t 2341

```

```

<210> SEQ ID NO 3
<211> LENGTH: 2341
<212> TYPE: DNA
<213> ORGANISM: Influenza A

```

```

<400> SEQUENCE: 3

```

```

agcgaaagca ggtcaattat attcaatatg gaaagaataa aagaactacg aaatctaag 60
tcgcagtctc gcaccgcga gatactcaca aaaaccaccg tggaccatat ggccataatc 120
aagaagtaca catcaggaag acaggagaag aaccagcac ttaggatgaa atggatgatg 180
gcaatgaaat atccaattac agcagacaag aggataacgg aaatgattcc tgagagaaat 240
gagcaaggac aaactttatg gagtaaaatg aatgatgccg gatcagaccg agtgatggta 300
tcacctctgg ctgtgacatg gtggaatagg aatggaccaa taacaaatac agttcattat 360
ccaaaaatct acaaaactta ttttgaaaga gtcgaaaggc taaagcatgg aacctttggc 420
cctgtccatt ttagaaacca agtcaaaata cgtcgagag ttgacataaa tcctggtcac 480
gcagatctca gtgcaagga ggcacaggat gtaatcatgg aagttgtttt ccctaacgaa 540
gtgggagcca ggatactaac atcggaatcg caactaacga taaccaaaga gaagaaagaa 600
gaactccagg attgcaaaat ttctcctttg atggttgcat acatgttga gagagaactg 660
gtccgcaaaa cgagattcct cccagtggtt ggtggaacaa gcagtgtgta cattgaagtg 720

```

-continued

ttgcatttga ctcaaggaa atgctgggaa cagatgtata ctccaggagg ggaagtggag	780
aatgatgatg ttgatcaaag cttgattatt gctgctagga acatagttag aagagctgca	840
gtatcagcag atccactagc atctttattg gagatgtgcc acagcacaca gattggtgga	900
attaggatgg tagacatcct taggcagaac ccaacagaag agcaagccgt ggatatatgc	960
aaggctgcaa tgggactgag aattagctca tcttcagtt ttggtggatt cacatttaag	1020
agaacaagcg gatcatcagt caagagagag gaagaggtgc ttacgggcaa tcttcaaaca	1080
ttgaagataa gagtgcataa gggatatgaa gagttcacia tgggtgggag aagagcaaca	1140
gccatactca gaaaagcaac caggagattg attcagctga tagtgagtgg gagagacgaa	1200
cagtcgattg ccgaagcaat aattgtggcc atggtatttt cacaagagga ttgtatgata	1260
aaagcagtca gaggtgatct gaatttcgtc aataggcgca atcaacgatt gaatcctatg	1320
catcaacttt taagacatct tcagaaggat gcgaaagtgc tttttcaaaa ttggggagtt	1380
gaacctatcg acaatgtgat gggaatgatt gggatattgc ccgacatgac tccaagcatc	1440
gagatgtcaa tgagaggagt gagaatcagc aaaatgggtg tagatgagta ctccagcacg	1500
gagagggtag tggtgagcat tgaccgtttt ttgagaatcc gggaccaacg aggaaatgta	1560
ctactgtctc ccgaggaggt cagtgaacaa cagggaacag agaaactgac aataacttac	1620
tcactgtcaa tgatgtggga gattaatggt cctgaatcag tgttggtaa tacctatcaa	1680
tggatcatca gaaactggga aactgttaaa attcagtggg ccagaaccc tacaatgcta	1740
tacaataaaa tggaatttga accatttcag tctttagtac ctaaggccat tagaggccaa	1800
tacagtgggt ttgtaagaac tctgttccaa caaatgaggg atgtgcttgg gacatttgat	1860
accgcacaga taataaaact tcttcccttc gcagccgctc caccaaagca aagtagaatg	1920
cagttctctc catttactgt gaatgtgagg ggatcaggaa tgagaatact tgtaaggggc	1980
aattctcctg tattcaacta taacaaggcc acgaagagac tcacagttct cggaaaggat	2040
gtggtcactt taactgaaga ccagatgaa ggcacagctg gagtggagtc cgctgttctg	2100
aggggattcc tcattctggg caaagaagac aagagatatg ggccagcact aagcatcaat	2160
gaactgagca accttgcgaa aggagagaag gctaattgtc taattgggca aggagacgtg	2220
gtgttggtaa tgaacgggaa acgggactct agcatactta ctgacagcca gacagcgacc	2280
aaaagaattc ggatggccat caattagtgt cgaatagttt aaaaacgacc ttgtttctac	2340
t	2341

<210> SEQ ID NO 4

<211> LENGTH: 1565

<212> TYPE: DNA

<213> ORGANISM: Influenza A

<400> SEQUENCE: 4

agcaaaagca gggtagataa tcaactactg agtgacatca aaatcatggc gtctcaaggc	60
accaaagcat cttacgaaca gatggagact gatggagaac gccagaatgc cactgaaatc	120
agagcatccg tcggaaaaat gattggtgga attggacgat tctacatcca aatgtgcacc	180
gaactcaaac tcagtgatta tgagggacgg ttgatccaaa acagcttaac aatagagaga	240
atggtgctct ctgcttttga cgaaaggaga aataaatacc ttgaagaaca tcccagtgcg	300
gggaaagatc ctaagaaaac tggaggacct atatacagga gagtaaacgg aaagtggatg	360
agagaactca tcccttatga caaagaagaa ataaggcgaa tctggcgcca agctaataat	420
ggtgacgatg caacggctgg tctgactcac atgatgatct ggcattccaa ttgtaatgat	480

-continued

gcaacttata agaggacaag agctcttggt cgcaccggaa tggatcccag gatgtgctct	540
ctgatgcaag gttcaactct ccctaggagg tctggagccg caggtgctgc agtcaaagga	600
gttggaacaa tggatgatga attggtcaga atgatcaaac gtgggatcaa tgatcggaac	660
ttctggagggt gtgagaatgg acgaaaaaca agaattgctt atgaaagaat gtgcaacatt	720
ctcaaaggga aatttcaaac tgctgcacaa aaagcaatga tggatcaagt gagagagagc	780
cggaaaccag ggaatgctga gttcgaagat ctcaacttttc tagcacggtc tgcactcata	840
ttgagagggt cgggtgctca caagtctgc ctgcctgcct gtgtgtatgg acctgccgta	900
gccagtgggt acgactttga aaggagaggga tactctctag tcggaataga ccctttcaga	960
ctgcttcaaa acagccaagt gtacagccta atcagaccaa atgagaatcc agcacacaag	1020
agtcaactgg tgtggatggc atgccattct gccgcatttg aagatctaag agtattaagc	1080
ttcatcaaa ggacgaaggt gctccaaga gggaagcttt ccactagagg agttcaaatt	1140
gcttccaatg aaaatatgga gactatggaa tcaagtacac ttgaactgag aagcaggtag	1200
tggggccataa ggaccagaag tggaggaaac accaatcaac agagggcgc tgcggggccaa	1260
atcagcatac aacctacgtt ctcagtacag agaaatctcc cttttgacag aacaaccatt	1320
atggcagcat tcaatgggaa tacagagggg agaacatctg acatgaggac cgaaatcata	1380
aggatgatgg aaagtgaag accagaagat gtgtctttcc agggggcgggg agtcttcgag	1440
ctctcgagc aaaaggcagc gagcccgatc gtgccttcct ttgacatgag taatgaagga	1500
tcttatttct tcggagacaa tgcagaggag tacgacaatt aaagaaaaat acccttgttt	1560
ctact	1565

<210> SEQ ID NO 5

<211> LENGTH: 1027

<212> TYPE: DNA

<213> ORGANISM: Influenza A

<400> SEQUENCE: 5

agcaaaagca ggtagatatt gaaagatgag tcttctaacc gaggtcgaaa cgtacgtact	60
ctctatcacc ccgtcaggcc ccctcaaagc cgagatcgca cagagacttg aagatgtctt	120
tgcagggaag aacaccgatc ttgaggttct catggaatgg ctaaagacaa gaccaatcct	180
gtcacctctg actaagggga ttttaggatt tgtgttcacg ctcaaccgtc ccagtgcgag	240
aggactgcag cgtagacgct ttgtccaaaa tgcccttaat gggaacgggg atccaaataa	300
catggacaaa gcagttaaac tgtataggaa gctcaagagg gagataacat tccatggggc	360
caaagaaatc tcactcagtt attctgctgg tgcacttgcc agttgtatgg gcctcatata	420
caacaggatg ggggctgtga ccactgaagt ggcatttggc ctggtatgtg caacctgtga	480
acagattgct gactcccagc atcgggtctca taggcaaatg gtgacaacaa ccaatccact	540
aatcagacat gagaacagaa tggttttagc cagcactaca gctaaggcta tggagcaaat	600
ggctggatcg agtgagcaag cagcagaggc catggagggt gctagtgcag ctagacaaat	660
gggtgcaagc atgagaacca ttgggactca tcctagctcc agtgctggtc tgaaaaatga	720
tcttcttgaa aatttgcagg cctatcagaa acgaatgggg gtgcagatgc aacggttcaa	780
gtgatcctct cactattgcc gcaaatatca ttgggactct gcacttgaca ttgtggattc	840
ttgatcgtct ttttttcaaa tgcatttacc gtgcgtttta atacggactg aaaggagggc	900
cttctacgga aggagtgcc aagtctatga gggaagaata tcgaaaggaa cagcagagtg	960

-continued

ctgtggatgc tgacgatggt cattttgtca gcatagagct ggagtaaaaa actaccttgt	1020
ttctact	1027

<210> SEQ ID NO 6
 <211> LENGTH: 890
 <212> TYPE: DNA
 <213> ORGANISM: Influenza A

<400> SEQUENCE: 6

agcaaaagca gggtgacaaa aacataatgg atccaaacac tgtgtcaagc tttcaggtag	60
attgttttct ttggcatgtc cgaaacgag ttgcagacca agaactaggc gatgccccat	120
tccttgatcg gcttcgcga gatcagaaat ccctaagagg aaggggcagt actctcggtc	180
tggacatcaa gacagccaca cgtgctggaa agcagatagt ggagcggatt ctgaaagaag	240
aatccgatga ggcacttaaa atgacatgg cctctgtacc tgcgtcgcgt tacctaactg	300
acatgactct tgaggaaatg tcaagggact ggtccatgct catacccaag cagaaagtgg	360
caggccctct ttgtatcaga atggaccagg cgatcatgga taagaacatc atactgaaag	420
cgaacttcag tgtgattttt gaccggctgg agactctaatt attgctaagg gctttcaccg	480
aagagggagc aattgttggc gaaatttcac cattgccttc tcttccagga catactgctg	540
aggatgtcaa aaatgcagtt ggagtcctca tcggaggact tgaatggaat gataacacag	600
ttcgagtctc tgaactcta cagagattcg cttggagaag cagtaatgag aatgggagac	660
ctccactcac tccaaaacag aaacgagaaa tggcgggaac aattaggtca gaagtttgaa	720
gaaataagat gggtgattga agaagtgaga caaaaactga agataacaga gaatagtttt	780
gagcaataa catttatgca agcctcat ctattgcttg aagtggagca agagataaga	840
actttctcgt ttcagcttat ttagtactaa aaaacacct tgtttctact	890

<210> SEQ ID NO 7
 <211> LENGTH: 1775
 <212> TYPE: DNA
 <213> ORGANISM: Influenza A

<400> SEQUENCE: 7

agcaaaagca ggggaaaata aaaacaacca aaatgaaggc aaacctactg gtctgttat	60
gtgcacttgc agctgcagat gcagacacaa tatgtatagg ctaccatgag aacaattcaa	120
ccgacactgt tgacacagta ctcgagaaga atgtgacagt gacacactct gttaacctgc	180
tcgaagacag ccacaacgga aaactatgta gattaaaagg aatagcccca ctacaattgg	240
ggaaatgtaa catcgccgga tggctcttgg gaaaccaga atgcgaccca ctgcttcag	300
tgagatcatg gtcctacatt gtagaaacac caaactctga gaatggaata tgttatccag	360
gagatttcat cgactatgag gagctgagg agcaattgag ctcagtgtca tcattcgaaa	420
gattcgaaat atttcccaa gaaagctcat ggcccaacca caacacaaac ggagtaacgg	480
cagcatgctc ccatgagggg aaaagcagtt ttacagaaa ttgctatgg ctgacggaga	540
aggagggctc atacccaaag ctgaaaaatt cttatgtgaa caaaaaggg aaagaagtcc	600
ttgtactgtg gggattcat caccgccta acagtaagga acaacagaat ctctatcaga	660
atgaaatgc ttatgtctct gtagtgactt caaattataa caggagattt accccgaaa	720
tagcagaaag acccaaagta agagatcaag ctgggaggat gaactattac tggaccttgc	780
taaaacccgg agacacaata atatttgagg caaatggaaa tctaatagca ccaatgtatg	840
ctttcgact gagtagaggc ttgggtccg gcatcatcac ctcaaacgca tcaatgcatg	900

-continued

```

agtgtaacac gaagtgtcaa acaccctgg gagctataaa cagcagtctc ccttaccaga    960
atatacacc agtcacaata ggagagtgcc caaaatacgt caggagtgcc aaattgagga    1020
tggttacagg actaaggaac attccgtcca ttcaatccag aggtctatct ggagccattg    1080
ccggttttat tgaaggggga tggactggaa tgatagatgg atgggtatgg tatcatcatc    1140
agaatgaaca gggatcaggc tatgcagcgg atcaaaaaag cacacaaaat gccattaacg    1200
ggattacaaa caaggtgaac actgttatcg agaaaatgaa cattcaattc acagctgtgg    1260
gtaaagaatt caacaaatta gaaaaaagga tggaaaattt aaataaaaaa gttgatgatg    1320
gatttctgga catttggaac tataatgcag aattgttagt tctactggaa aatgaaagga    1380
ctctggattt ccatgactca aatgtgaaga atctgtatga gaaagtaaaa agccaattaa    1440
agaataatgc caaagaaatc ggaaatggat gttttgagtt ctaccacaag tgtgacaatg    1500
aatgcatgga aagtgtgaag aatgggactt atgattatcc caaatattca gaagagtcaa    1560
agttgaacag ggaagagga gatggagtga aattggaatc aatggggatc tatcagattc    1620
tggcgatcta ctcaactgtc gccagttcac tgggtctttt ggtctcctg ggggcaatca    1680
gtttctggat gtgttctaata ggatctttgc agtgcagaat atgcatctga gattagaatt    1740
tcagagatat gaggaanaac acccttgttt ctact                                1775

```

```

<210> SEQ ID NO 8
<211> LENGTH: 1413
<212> TYPE: DNA
<213> ORGANISM: Influenza A

```

```

<400> SEQUENCE: 8

```

```

agcaaaagca ggggtttaaa atgaatccaa atcagaaaat aataaccatt ggatcaatct    60
gtctggtagt cggactaatt agcctaatat tgcaaatagg gaataataat tcaatatgga    120
ttagccattc aattcaaaact ggaagtcaaa accatactgg aatatgcaac caaaacatca    180
ttacctataa aaatagcacc tgggtaaagg acacaacttc agtgatatta accggcaatt    240
catctctttg tcccatccgt ggggtgggcta tatacagcaa agacaatagc ataagaattg    300
gttccaaagg agacgttttt gtcataagag agccctttat ttcattgtct cacttggaat    360
gcaggacctt ttttctgacc caagggtcct tactgaatga caagcattca agtgggactg    420
ttaaggacag aagcccttat agggccttaa tgagctgccc tgctcggtgaa gctccgtccc    480
cgtacaattc aagatttgaa tcgggttgctt ggtcagcaag tgcattgat gatggcatgg    540
gctggctaac aatcggaatt tcagggtccag ataattggagc agtggctgta ttaaaataca    600
acggcataat aactgaaacc ataaaaagtt ggaggaagaa aatattgagg acacaagagt    660
ctgaatgtgc ctgtgtaaat ggttcattgt ttactataat gactgatggc ccgagtgatg    720
ggctggcctc gtacaaaatt ttcaagatcg aaaaggggaa ggttactaaa tcaatagagt    780
tgaatgcacc taattctcac tatgaggaat gtccctgtta ccctgatacc ggcaagtgaa    840
tgtgtgtgtg cagagacaat tggcatgggt cgaaccggcc atgggtgtct ttcgatcaaa    900
acctggatta tcaaatagga tacatctgca gtgggggttt cggtgacaac ccgctgccc    960
aagatggaac aggcagctgt ggtccagtgt atgttgatgg agcaaacgga gtaaagggat    1020
tttcatatag gtatggtaat ggtgtttgga taggaaggac caaaagtcac agttccagac    1080
atgggtttga gatgatttgg gatcctaata gatggacaga gactgatagt aagttctctg    1140
tgaggcaaga tgtgtgggca atgactgatt ggtcagggta tagcggaagt ttcgttcaac    1200

```

-continued

```

atcctgagct gacagggcta gactgtatga ggccgtgctt ctgggttgaa ttaatcaggg 1260
gacgacctaa agaaaaaaca atctggacta gtgcgagcag catttctttt tgtggcgtga 1320
atagtatac tgtagattgg tcttggccag acggtgctga gttgccattc agcattgaca 1380
agtagtctgt tcaaaaaact cctgttttct act 1413

```

```

<210> SEQ ID NO 9
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A synthetic oligonucleotide

```

```

<400> SEQUENCE: 9

```

```

cacacacggt ctccgggagc gaaagcaggc a 31

```

```

<210> SEQ ID NO 10
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A synthetic oligonucleotide

```

```

<400> SEQUENCE: 10

```

```

ccaggacact gaaatttctt tcac 24

```

```

<210> SEQ ID NO 11
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A synthetic oligonucleotide

```

```

<400> SEQUENCE: 11

```

```

cacacaggtc tcctattagt agaacaagg cattt 35

```

```

<210> SEQ ID NO 12
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A synthetic oligonucleotide

```

```

<400> SEQUENCE: 12

```

```

cacacaggtc tccgggagcg aaagcaggtc 30

```

```

<210> SEQ ID NO 13
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A synthetic oligonucleotide

```

```

<400> SEQUENCE: 13

```

```

cacacacgtc tccatcatatc aatcctcttg 30

```

```

<210> SEQ ID NO 14
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A synthetic oligonucleotide

```

```

<400> SEQUENCE: 14

```

```

ctcctctgat ggtggcatatc 20

```


-continued

<210> SEQ ID NO 15
<211> LENGTH: 36
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A synthetic oligonucleotide

<400> SEQUENCE: 15

cacacaggtc tcctattagt agaaacaagg tcgttt 36

<210> SEQ ID NO 16
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A synthetic oligonucleotide

<400> SEQUENCE: 16

cacacacgtc tccgggagcg aaagcaggta c 31

<210> SEQ ID NO 17
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A synthetic oligonucleotide

<400> SEQUENCE: 17

cacacacgtc tcctattagt agaaacaagg tactt 35

<210> SEQ ID NO 18
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A synthetic oligonucleotide

<400> SEQUENCE: 18

cacacacgtc tccgggagca aaagcagggg 30

<210> SEQ ID NO 19
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A synthetic oligonucleotide

<400> SEQUENCE: 19

cacacacgtc tcctattagt agaaacaagg gtgtttt 37

<210> SEQ ID NO 20
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: A synthetic oligonucleotide

<400> SEQUENCE: 20

cacacacgtc tccgggagca aaagcagggt a 31

<210> SEQ ID NO 21
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: A synthetic oligonucleotide

<400> SEQUENCE: 21

cacacacgtc tcctattagt agaaacaagg gtattttt 38

<210> SEQ ID NO 22

<211> LENGTH: 31

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: A synthetic oligonucleotide

<400> SEQUENCE: 22

cacacaggtc tccgggagca aaagcaggag t 31

<210> SEQ ID NO 23

<211> LENGTH: 38

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: A synthetic oligonucleotide

<400> SEQUENCE: 23

cacacaggtc tggattagt agaaacaagg agtttttt 38

<210> SEQ ID NO 24

<211> LENGTH: 31

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: A synthetic oligonucleotide

<400> SEQUENCE: 24

cacacacgtc tccgggagca aaagcaggta g 31

<210> SEQ ID NO 25

<211> LENGTH: 38

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: A synthetic oligonucleotide

<400> SEQUENCE: 25

cacacacgtc tcctattagt agaaacaagg tagttttt 38

<210> SEQ ID NO 26

<211> LENGTH: 31

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: A synthetic oligonucleotide

<400> SEQUENCE: 26

cacacacgtc tccgggagca aaagcagggt g 31

<210> SEQ ID NO 27

<211> LENGTH: 37

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: A synthetic oligonucleotide

<400> SEQUENCE: 27

cacacacgtc tcctattagt agaaacaagg gtgtttt 37

-continued

<210> SEQ ID NO 28
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: A synthetic oligonucleotide

<400> SEQUENCE: 28

gggtttgtat ttgtgtgtca cc 22

<210> SEQ ID NO 29
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Influenza A

<400> SEQUENCE: 29

Arg Glu Arg Arg Arg Lys Lys Arg
 1 5

<210> SEQ ID NO 30
 <211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: A synthetic peptide

<400> SEQUENCE: 30

Thr Glu Thr Arg
 1

<210> SEQ ID NO 31
 <211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: A synthetic peptide

<400> SEQUENCE: 31

Arg Glu Thr Arg
 1

<210> SEQ ID NO 32
 <211> LENGTH: 20
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: A synthetic peptide

<400> SEQUENCE: 32

Cys Asn Gln Ser Ile Ile Thr Tyr Glu Asn Asn Thr Trp Val Asn Gln
 1 5 10 15

Thr Tyr Val Asn
 20

<210> SEQ ID NO 33
 <211> LENGTH: 2341
 <212> TYPE: DNA
 <213> ORGANISM: Influenza A

<400> SEQUENCE: 33

agcgaaagca ggtcaattat attcaatatg gaaagaataa aagaactaag aaatctaatag 60

tcgcagtctc gcacccgcca gataactaca aaaaccaccg tggaccatat ggccataatc 120

aagaagtaca catcaggaag acaggagaag aaccacgac ttaggatgaa atggatgatg 180

gcaatgaaat atccaattac agcagacaag aggataacgg aaatgattcc tgagagaaat 240

-continued

gagcaaggac aaactttatg gagtaaaatg aatgatgccg gatcagaccg agtgatggta	300
tcacctctgg ctgtgacatg gtggaatagg aatggaccaa tgacaaatac agttcattat	360
ccaaaaatct acaaaactta ttttgaaga gtcgaaaggc taaagcatgg aacctttggc	420
cctgtccatt ttagaaacca agtcaaaata cgtcggagag ttgacataaa tcctggtc	480
gcagatctca gtgccaagga ggcacaggat gtaatcatgg aagttgtttt ccctaacgaa	540
gtgggagcca ggatactaac atcggaaatcg caactaacga taaccaaga gaagaaagaa	600
gaactccagg attgcaaaat ttctcctttg atggttgcac acatgttggg gagagaactg	660
gtccgcaaaa cgagattcct ccagtggtg ggtggaacaa gcagtgtgta cattgaagtg	720
ttgcatttga ctcaaggaa atgctgggaa cagatgtata ctccaggagg ggaagtgaag	780
aatgatgatg ttgatcaaa cttgattatt gctgctagga acatagtggg aagagctgca	840
gtatcagcag acccactagc atctttattg gagatgtgcc acagcacaca gattggtgga	900
attaggatgg tagacatcct taagcagaac ccaacagaag agcaagccgt ggatatatgc	960
aaggctgcaa tgggactgag aattagctca tccttcagtt ttggtggatt cacatttaag	1020
agaacaagcg gatcatcagt caagagagag gaagaggtgc ttacgggcaa tcttcaaca	1080
ttgaagataa gagtgcata gggatctgaa gagttcaca tgggtgggag aagagcaaca	1140
gccatactca gaaaagcaac caggagattg attcagctga tagtgagtgg gagagacgaa	1200
cagtcgattg ccgaagcaat aattgtggcc atggtatatt cacaagagga ttgtatgata	1260
aaagcagtta gaggtgatct gaatttcgtc aatagggcga atcagcgact gaatcctatg	1320
catcaacttt taagacattt tcagaaggat gcgaaagtgc tttttcaaaa ttggggagtt	1380
gaacctatcg acaatgtgat gggaatgatt gggatattgc ccgacatgac tccaagcatc	1440
gagatgtcaa tgagaggagt gagaatcagc aaaatgggtg tagatgagta ctccagcacg	1500
gagagggtag tggtagcat tgaccggttc ttgagagtca gggaccaacg aggaaatgta	1560
ctactgtctc ccgaggaggt cagtgaaca cagggaacag agaaactgac aataacttac	1620
tcacgtcaa tgatgtggga gattaatggt cctgaatcag tggtagtcaa tacctatcaa	1680
tggatcatca gaaactggga aactgttaaa attcagtggg ccagaaacc tacaatgcta	1740
tacaataaaa tggaatttga accatttcag tcttttagta ctaaggccat tagaggccaa	1800
tacagtgggt ttgtaagaac tctgttccaa caaatgagg atgtgcttgg gacatttgat	1860
accgcacaga taataaaact tcttccttc gcagccgctc caccaaagca agtagaatg	1920
cagttctcct catttactgt gaatgtgagg ggtacaggaa tgagaatact tgtaaggggc	1980
aattctcctg tattcaacta caacaaggcc acgaagagac tcacagtctc cggaaggat	2040
gctggcactt taaccgaaga ccagatgaa ggcacagctg agtgaggatc cgctgttctg	2100
aggggattcc tcattctggg caaagaagac aggagatatg ggcagcatt aagcatcaat	2160
gaactgagca accttgcaaa aggagagaag gctaattgtc taattgggca aggagacgtg	2220
gtgttggtaa tgaacgaaa acgggactct agcatactta ctgacagcca gacagcgacc	2280
aaaagaatcc ggatggccat caattagtgt cgaatagttt aaaaacgacc ttgtttctac	2340
t	2341

<210> SEQ ID NO 34

<211> LENGTH: 2341

<212> TYPE: DNA

<213> ORGANISM: Influenza A

-continued

<400> SEQUENCE: 34

agcgaaagca ggcaaacccat ttgaatggat gtcaatccga ccttactttt cttaaaagtg	60
ccagcacaaa atgctataag cacaactttc ccttataccg gagaccctcc ttacagccat	120
gggacaggaa caggatacac catggatact gtcaacagga cacatcagta ctgagaaaag	180
ggaagatgga caacaaacac cgaaactgga gcaccgcaac tcaacccgat tgatgggcca	240
ctgccagaag acaatgaacc aagtgggtat gcccaaacag attgtgtatt ggaagcaatg	300
gctttccttg aggaatccca tcctgggtatt ttgaaaact cgtgtattga aacgatggag	360
gttggtcagc aaacacgagt agacaagctg acacaaggcc gacagaccta tgactggact	420
ttaaatagaa accagcctgc tgcaacagca ttggccaaca caatagaagt gttcagatca	480
aatggcctca cggccaatga gtcaggaagg ctcatagact tccttaagga tgtaatggag	540
tcaatgaaaa aagaagaaat ggggatcaca actcattttc agagaaagag acgggtgaga	600
gacaatatga ctaagaaaat gataacacag agaacaatag gtaaaaggaa acagagattg	660
aacaaaaggg gttatctaata tagagcattg accctgaaca caatgaccaa agatgctgag	720
agagggaagc taaaacggag agcaattgca accccaggga tgcaataag ggggtttgta	780
tactttgttg agacactggc aaggagtata tgtgagaaac ttgaacaatc agggttgcca	840
gttggaggca atgagaagaa agcaaagttg gcaaatgttg taagggaagat gatgaccaat	900
tctcaggaca ccgaactttc ttccaccatc actggagata acaccaaag gaacgaaaaat	960
cagaatcctc ggatgttttt ggccatgatc acatatatga ccagaaatca gccggaatgg	1020
ttcagaaatg ttctaagtat tgctccaata atgttctcaa acaaaatggc gagactggga	1080
aaagggtata tgtttgagag caagagtatg aaacttagaa ctcaaatacc tgcagaaatg	1140
ctagcaagca ttgatttgaa atatttcaat gattcaacaa gaaagaagat tgaaaaaatc	1200
cgaccgctct taatagaggg gactgcatca ttgagccctg gaatgatgat gggcatgttc	1260
aatatgttaa gcactgtatt aggcgtctcc atcctgaatc ttggacaaa gagatacacc	1320
aagactactt actggtggga tggctctcaa tcctctgacg attttgctct gattgtgaat	1380
gcacccaatc atgaagggat tcaagccgga gtcgacaggt tttatcgaa ctgtaagcta	1440
cttggaatca atatgagcaa gaaaaagtct tacataaaca gaacaggtag atttgaattc	1500
acaagttttt tctatcgta tgggtttggt gccaatttca gcatggagct tcccagtttt	1560
ggggtgtctg ggatcaacga gtcagcggac atgagtattg gagttactgt catcaaaac	1620
aatatgataa acaatgatct tggtcacgca acagctcaaa tggcccttca gttgttcac	1680
aaagattaca ggtacacgta ccgatgccat agagggtgaca cacaataca aacccaaga	1740
tcatttgaaa taaagaaact gtgggagcaa acccgttcca aagctggact gctggctctc	1800
gacggaggcc caaatattata caacattaga aatctccaca ttcctgaagt ctgcctaaaa	1860
tgggaattga tggatgagga ttaccagggg cgtttatgca acccactgaa cccatttgtc	1920
agccataaag aaattgaatc aatgaacaat gcagtgatga tgccagcaca tgggtccagcc	1980
aaaaacatgg agtatgatgc tgttgcaaca acacactcct ggatcccaa aagaaatcga	2040
tccatcttga atacaagtca aagaggagta cttgaagatg aacaaatgta ccaagggtgc	2100
tgcaatttat ttgaaaaatt cttccccagc agttcataca gaagaccagt cgggatatcc	2160
agtatggtgg aggtatggt ttccagagcc cgaattgatg cacggattga tttcgaatct	2220
ggaaggataa agaaagaaga gttcactgag atcatgaaga tctgttccac cattgaagag	2280
ctcagacggc aaaaatagtg aatttagctt gtccttcacg aaaaaatgcc ttgtttctac	2340

-continued

t	2341
 <210> SEQ ID NO 35	
<211> LENGTH: 2233	
<212> TYPE: DNA	
<213> ORGANISM: Influenza A	
 <400> SEQUENCE: 35	
agcgaaagca ggtactgatt caaaatggaa gattttgtgc gacaatgctt caatccgatg	60
attgtcgagc ttgcggaaaa aacaatgaaa gagtatgggg aggacctgaa aatcgaaaca	120
aacaaatttg cagcaatatg cactcacttg gaagtatgct tcatgtattc agatttccac	180
ttcatcaatg agcaaggcga gtcaataatc gtagaacttg gtgatcctaa tgcacttttg	240
aagcacagat ttgaataat cgagggaaga gatcgacaaa tggcctggac agtagtaaac	300
agtatttgca acactacagg ggctgagaaa ccaaagtctt taccagattt gtatgattac	360
aagggaaaata gattcatcga aattggagta acaaggagag aagttcacat atactatctg	420
gaaaaggcca ataaaattaa atctgagaaa acacacatcc acattttctc gttcactggg	480
gaagaaatgg ccacaagggc cgactacact ctgatgaag aaagcagggc taggatcaaa	540
accaggctat tcaccataag acaagaaatg gccagcagag gcctctggga ttctttctgt	600
cagtccgaga gaggagaaga gacaattgaa gaaaggtttg aaatcacagg aacaatgcgc	660
aagcttgccg accaaagtct cccgccgaac ttctccagcc ttgaaaattt tagagcctat	720
gtggatggat tcgaaccgaa cggctacatt gagggcaagc tgtctcaaat gtccaaagaa	780
gtaaatgcta gaattgaacc ttttttgaaa acaacaccac gaccacttag acttccgaat	840
gggcctccct gttctcagcg gtccaaattc ctgctgatgg atgccttaaa attaagcatt	900
gaggacccaa gtcatgaagg agagggaata ccgctatatg atgcaatcaa atgcatgaga	960
acattctttg gatggaagga acccaatggt gttaaaccac acgaaaaggg aataaatcca	1020
aattatcttc tgtcatggaa gcaagtactg gcagaactgc aggacattga gaatgaggag	1080
aaaattccaa agactaaaaa tatgaaaaaa acaagtcagc taaagtgggc acttggtgag	1140
aacatggcac cagaaaaggt agactttgac gactgtaaag atgtagggtga tttgaagcaa	1200
tatgatagtg atgaaccaga attgaggtcg cttgcaagtt ggattcagaa tgagttcaac	1260
aaggcatgcg aactgacaga ttcaagctgg atagagcttg atgagattgg agaagatgtg	1320
gtccaattg aacacattgc aagcatgaga aggaattatt tcacatcaga ggtgtctcac	1380
tgcagagcca cagaatacat aatgaagggg gtgtacatca atactgcctt acttaatgca	1440
tcttgtgcag caatggatga tttccaatta attccaatga taagcaagtg tagaaactaag	1500
gagggaaggc gaaagaccaa cttgtatggt ttcatacataa aaggaagatc ccacttaagg	1560
aatgacaccg acgtggtaaa ctttgtgagc atggagtttt ctctcactga cccaagactt	1620
gaaccacaca aatgggagaa gtactgtggt cttgagatag gagatattgt tctaagaagt	1680
gccataggcc aggtttcaag gcccatgttc ttgtatgtga ggacaaatgg aacctcaaaa	1740
attaaaatga aatggggaat ggagatgagg cgttgtctcc tccagtcact tcaacaaatt	1800
gagagtatga ttgaagctga gtcctctgtc aaagagaaag acatgaccaa agagtctctt	1860
gagaacaaat cagaaacatg gccattgga gagtctccca aaggagtgga ggaaagtctc	1920
attgggaagg tctgcaggac tttattagca aagtcggtat ttaacagctt gtatgcattc	1980
ccacaactag aaggattttc agctgaatca agaaaactgc ttcttatcgt tcagggtctt	2040

-continued

```

agggacaatc tggaaacctgg gacctttgat cttggggggc tatatgaagc aattgaggag 2100
tgccataatta atgatccctg ggttttgctt aatgcttctt gggtcaactc cttccttaca 2160
catgcattga gttagtgtg gcagtgtac tatgtgtat ccatactgtc caaaaaagta 2220
ccttgtttct act 2233

```

```

<210> SEQ ID NO 36
<211> LENGTH: 1565
<212> TYPE: DNA
<213> ORGANISM: Influenza A

```

```

<400> SEQUENCE: 36

```

```

agcaaaagca gggtagataa tcaactcactg agtgacatca aaatcatggc gtcccaaggc 60
accaaacggt cttacgaaca gatggagact gatggagaac gccagaatgc cactgaaatc 120
agagcatccg tcgaaaaaat gattggtgga attggacgat tctacatcca aatgtgcaca 180
gaacttaaac tcagtgatta tgagggacgg ttgatccaaa acagcttaac aatagagaga 240
atggtgctct ctgcttttga cgaaaggaga aataaatacc tggaagaaca tcccagtgcg 300
gggaaagatc ctaagaaaac tggaggacct atatacagaa gagtaaaccg aaagtggatg 360
agagaactca tcctttatga caaagaagaa ataaggcgaa tctggcgcca agctaataat 420
ggtgacgatg caacggctgg tctgactcac atgatgatct ggcatccaa tttgaatgat 480
gcaacttata agaggacaag ggctcttgtt cgcaccggaa tggatcccag gatgtgctct 540
ctgatgcaag gttcaactct cctaggagg tctggagcgg caggtgctgc agtcaaagga 600
gttgaacaa tggatgatga attggtcagg atgatcaaac gtgggatcaa tgatcggaac 660
ttctggaggg gtgagaatgg acgaaaaaca agaattgctt atgaaagaat gtgcaacatt 720
ctcaaaggga aatttcaaac tgctgcacaa aaagcaatga tggatcaagt gagagagagc 780
cggaaccagg ggaatgtga gttcgaagat ctcaactttc tagcacggtc tgcactcata 840
ttgagagggt cgggtgtcga caagtctgc ctgcctgctt gtgtgtatgg acctgccgta 900
gccagtgggt acgactttga aagagaggga tactctctag tcggaataga ccctttcaga 960
ctgcttcaaa acagccaagt gtacagccta atcagacca atgagaatcc agcacacaag 1020
agtcaactgg tgtggatggc atgccattct gccgcatttg aagatctaag agtattgagc 1080
ttcatcaaa ggacgaaggt ggtcccaaga gggaagcttt ccactagagg agttcaaatt 1140
gcttccaatg aaaatatgga gactatggaa tcaagtacac ttgaactgag aagcaggtac 1200
tgggccataa ggaccagaag tggaggaaac accaatcaac agagggcatc tgcgggccaa 1260
atcagcatac aacctacgtt ctcagtacag agaaatctcc cttttgacag aacaaccggt 1320
atggcagcat tcaactggaa tacagagggg agaacatctg acatgaggac cgaaatcata 1380
aggatgatgg aaagtgaag accagaagat gtgtctttcc aggggcgggg agtcttcgag 1440
ctctcggacg aaaaggcagc gagcccgatc gtgccttctt ttgacatgag taatgaagga 1500
tcttatttct tcggagacaa tgcagaggag tacgacaatt aaagaaaaat acccttgttt 1560
ctact 1565

```

```

<210> SEQ ID NO 37
<211> LENGTH: 1027
<212> TYPE: DNA
<213> ORGANISM: Influenza A

```

```

<400> SEQUENCE: 37

```

```

agcaaaagca ggtagatatt gaaagatgag tcttctaacc gaggtcgaaa cgtacgttct 60

```

-continued

```

ctctatcatc cgcgcagccc ccctcaaagc cgagatcgca cagagacttg aagatgtctt 120
tgcagggaag aacaccgatc ttgaggttct catggaatgg ctaaagacaa gaccaatcct 180
gtcacctctg actaagggga ttttaggatt tgtgttcacg ctcaccgtgc ccagtgcgag 240
aggactgcag cgtagacgct ttgtccaaaa tgccttaaat gggaacgggg atccaaataa 300
catggacaaa gcagttaaac tgtataggaa gctcaagagg gagataacat tccatggggc 360
caaagaaato tcaactcagtt attctgctgg tgcacttgcc agttgtatgg gcctcatata 420
caacaggatg ggggctgtga ccactgaagt ggcatttggc ctggtatgtg caacctgtga 480
acagattgct gactcccagc atcgggtctca taggcaaagtg gtgacaacaa ccaaccact 540
aatcagacat gagaacagaa tggttttagc cagcactaca gctaaggcta tggagcaaat 600
ggctggatcg agtgagcaag cagcagaggc catggagggt gctagtcagg ctaggcaaat 660
ggtgcaagcg atgagaacca ttgggactca tcctagctcc agtgctggtc tgaaaaatga 720
tcttcttgaa aatttgcagg cctatcagaa acgaatgggg gtgcagatgc aacggttcaa 780
gtgatcctct cgctattgcc gcaaatatca ttgggatctt gcaactgata ttgtggattc 840
ttgatcgtct ttttttcaaa tgcatttacc gtcgctttaa atacggactg aaaggagggc 900
cttctacgga aggagtgcga aagtctatga ggaagaata tcgaaaggaa cagcagagtg 960
ctgtggatgc tgacgatggt cattttgtca gcatagagct ggagtaaaaa actaccttgt 1020
ttctact 1027

```

```

<210> SEQ ID NO 38
<211> LENGTH: 890
<212> TYPE: DNA
<213> ORGANISM: Influenza A
<400> SEQUENCE: 38

```

```

agcaaaagca gggtgacaaa gacataatgg atccaaacac tgtgtcaagc tttcaggtag 60
attgctttct ttggcatgtc cgcaaacgag ttgcagacca agaactaggt gatgcccact 120
tccttgatcg gcttcgcgga gatcagaaat ccctaagagg aaggggcagc actcttggtc 180
tggacatcga gacagccaca cgtgctggaa agcagatagt ggagcggatt ctgaaagaag 240
aatccgatga ggcacttaaa atgaccatgg cctctgtacc tgcgtcgcgt tacctaaccg 300
acatgactct tgaggaaatg tcaagggaat ggtccatgct catacccaag cagaaagtgg 360
caggccctct ttgtatcaga atggaccagg cgatcatgga taaaaacatc atactgaaag 420
cgaacttcag tgtgattttt gaccggctgg agactctaat attgctaagg gctttcaccg 480
aagagggagc aattgttggc gaaatttcac cattgccttc tcttccagga catactgctg 540
aggatgtcaa aaatgcagtt ggagtcctca tcggaggact tgaatggaat gataacacag 600
ttcgagtctc tgaaactcta cagagattcg cttggagaag cagtaatgag aatgggagac 660
ctccactcac tccaaaacag aaacgagaaa tggcgggaac aattaggtea gaagtttgaa 720
gaaataagat ggttgattga agaagtgaga cacaaactga aggtaacaga gaatagtttt 780
gagcaataaa catttatgca agccttacat ctattgcttg aagtggagca agagataaga 840
actttctcat ttcagcttat ttaataataa aaaacaccct tgtttctact 890

```

What is claimed is:

1. A plurality of influenza virus vectors for a reassortant, comprising
 - a) a vector for vRNA production comprising a promoter operably linked to an influenza virus PA cDNA linked to

65

a transcription termination sequence, a vector for vRNA production comprising a promoter operably linked to an influenza virus PB1 cDNA linked to a transcription termination sequence, a vector for vRNA production comprising a promoter operably linked to an influenza virus

PB2 cDNA linked to a transcription termination sequence, a vector for vRNA production comprising a promoter operably linked to an influenza virus HA cDNA linked to a transcription termination sequence, a vector for vRNA production comprising a promoter operably linked to an influenza virus NP cDNA linked to a transcription termination sequence, a vector for vRNA production comprising a promoter operably linked to an influenza virus NA cDNA linked to a transcription termination sequence, a vector for vRNA production comprising a promoter operably linked to an influenza virus M cDNA linked to a transcription termination sequence, and a vector for vRNA production comprising a promoter operably linked to an influenza virus NS cDNA linked to a transcription termination sequence, wherein the vectors for the reassortant include cDNAs for PB1, PB2, PA, NP, M, and NA with sequences that encode a polypeptide having at least 97% contiguous amino acid sequence identity to a corresponding polypeptide encoded by SEQ ID NOs:1-5 and a polypeptide having at least 90% contiguous amino acid sequence identity to a corresponding polypeptide encoded by SEQ ID NO:8, a cDNA for NS that has a Glu residue at position 55 corresponding to position 55 in SEQ ID NO:38 and encodes a polypeptide having at least 97% contiguous amino acid sequence identity to a corresponding polypeptide encoded by SEQ ID NO:6, and a cDNA for HA that is from a different influenza virus strain than a strain having gene segments with sequences corresponding to the cDNAs for PB1, PB2, PA, NP, M, NS, and NA having SEQ ID Nos. 1-6 and 8, and wherein the cDNA for PB2 has a Tyr at residue 360; and

b) a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus PA, a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus PB1, a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus PB2, and a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus NP, and optionally a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus HA, a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus NA, a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus M1, a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus M2, or a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus NS2, which plurality of vectors, when introduced to an avian or mammalian cell, result in production of the reassortant which is capable of enhanced replication in embryonated eggs and MDCK cells relative to a corresponding influenza virus that does not have a NA with sequences that encode polypeptide having at least 90% contiguous amino acid sequence identity to a corresponding polypeptide encoded by SEQ ID NO:8 or does not have a NS with a Glu residue at position 55 corresponding to position 55 in SEQ ID NO:38 or a PB2 with a Tyr residue at position 360.

2. The plurality of vectors of claim 1, wherein the cDNA for NS encodes a polypeptide encoded by SEQ ID NO:38.

3. The plurality of vectors of claim 1, wherein the promoter is a RNA polymerase I promoter, a RNA polymerase II promoter, a RNA polymerase III promoter, a T3 promoter or a T7 promoter.

4. The plurality of vectors of claim 1, wherein the HA is a type A HA.

5. The plurality of vectors of claim 1 wherein the HA is a type B HA.

6. The plurality of vectors of claim 1, wherein the NA is N1.

7. The plurality of vectors of claim 1, wherein the HA is H5.

8. The plurality of vectors of claim 1, wherein a plurality of the vectors of a) comprise a RNA polymerase I promoter or a RNA polymerase II promoter.

9. The plurality of vectors of claim 8, wherein the RNA polymerase I promoter is a human RNA polymerase I promoter.

10. The plurality of vectors of claim 1, wherein all of the vectors of b) comprise a RNA polymerase II promoter.

11. The plurality of vectors of claim 1, wherein each vector of a) is on a separate plasmid.

12. The plurality of vectors of claim 1, wherein each vector of b) is on a separate plasmid.

13. The plurality of vectors of claim 1, wherein the each of the vectors of b) further comprise a RNA transcription termination sequence.

14. The plurality of vectors of claim 1, wherein the HA is a chimeric HA.

15. The plurality of vectors of claim 1, wherein the cDNA for HA does not encode a polypeptide corresponding to the polypeptide encoded by SEQ ID NO:7.

16. The plurality of vectors of claim 1, wherein the NA comprises a stalk region that is greater than 20 amino acids in length.

17. The plurality of vectors of claim 1, wherein the HA is an avirulent H5 HA.

18. A method to prepare influenza virus, comprising: contacting a cell with a vector for vRNA production comprising a promoter operably linked to an influenza virus PA cDNA linked to a transcription termination sequence, a vector for vRNA production comprising a promoter operably linked to an influenza virus PB1 cDNA linked to a transcription termination sequence, a vector for vRNA production comprising a promoter operably linked to an influenza virus PB2 cDNA linked to a transcription termination sequence, a vector for vRNA production comprising a promoter operably linked to an influenza virus HA cDNA linked to a transcription termination sequence, a vector for vRNA production comprising a promoter operably linked to an influenza virus NP cDNA linked to a transcription termination sequence, a vector for vRNA production comprising a promoter operably linked to an influenza virus NA cDNA linked to a transcription termination sequence, a vector for vRNA production comprising a promoter operably linked to an influenza virus M cDNA linked to a transcription termination sequence, and a vector for vRNA production comprising a promoter operably linked to an influenza virus NS cDNA linked to a transcription termination sequence, wherein the vectors for the 6:1 reassortant include cDNAs for PB1, PB2, PA, NP, M, and NA with sequences that encode a polypeptide having at least 97% contiguous amino acid sequence identity to a corresponding polypeptide encoded by SEQ ID NOs:1-5 and a polypeptide having at least 90% amino acid sequence identity to a corresponding polypeptide having SEQ ID NO:8, a cDNA for NS that has a Glu residue at position 55 corresponding to position 55 in SEQ ID NO:38 and encodes a polypeptide having at least 97% contiguous amino acid sequence identity to a cor-

75

responding polypeptide encoded by SEQ ID NO:6, and a cDNA for HA that is from a different influenza virus strain than a strain having gene segments with sequences corresponding to the cDNAs for PB1, PB2, PA, NP, M, NS, and NA having SEQ ID Nos. 1-6 and 8, and wherein the cDNA for PB2 has a Tyr at residue 360 ; and a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus PA, a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus PB1, a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus PB2, and a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus NP, and optionally a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus HA, a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus NA, a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus M1, a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus M2, or a vector for mRNA production comprising a promoter operably linked to a DNA segment encoding influenza virus NS2; in an amount effective to yield infectious influenza virus, wherein the infectious influenza virus has enhanced replication in embryonated eggs and MDCK cells relative to a corresponding influenza virus that does not have a NA with sequences that encode a polypeptide having at least 90% contiguous amino acid sequence identity to a corresponding polypeptide encoded by SEQ ID NO:8 or does not have a NS with a Glu residue at position 55 corresponding to position 55 in SEQ ID NO:38 or a PB2 with a Tyr residue at position 360.

19. The method of claim 18 further comprising isolating the virus.

20. Virus obtained by the method of claim 19.

21. An isolated recombinant influenza virus comprising a viral segment for PB1, PB2, PA, NP, M, and NA that has sequences for a polypeptide having at least 95% amino acid sequence identity to a corresponding polypeptide encoded by SEQ ID NOs:1-5 and at least 90% amino acid sequence identity to a corresponding polypeptide encoded by SEQ ID NO:8, a viral segment for NS with a Glu residue at position 55 corresponding to position 55 in SEQ ID NO:38 and at least

76

95% amino acid sequence identity to a corresponding polypeptide encoded by SEQ ID NO:6, and a viral segment for a heterologous HA, wherein the PB2 has a Tyr residue at position 360.

22. The isolated recombinant virus of claim 21, wherein the influenza virus that replicates to high titers is PR8HG.

23. The isolated recombinant influenza virus of claim 21, wherein the viral segment for HA is for H5.

24. An inactivated influenza virus vaccine comprising the isolated recombinant virus of claim 21.

25. The plurality of vectors of claim 1, wherein the cDNAs for PB1, PB2, PA, NP, and M encode a polypeptide having at least 99% contiguous amino acid sequence identity to a corresponding polypeptide encoded by SEQ ID NOs:1-5.

26. The isolated virus of claim 21, wherein the viral segments for PB1, PB2, PA, NP, and M have sequences for a polypeptide having at least 99% amino acid sequence identity to a corresponding polypeptide encoded by SEQ ID NOs:1-5.

27. An isolated recombinant influenza virus comprising a viral segment for PB1, PB2, PA, NP, M, and NS that has sequences for a polypeptide having at least 99% amino acid sequence identity to a corresponding polypeptide encoded by SEQ ID NOs:1-6, a viral segment for a heterologous HA, and a viral segment for NA, wherein the viral segment for NS encodes a NS1 protein has a Glu residue at position 55 corresponding to position 55 in SEQ ID NO:38, and wherein the viral segment for PB2 encodes a PB2 with a Tyr residue at position 360.

28. The recombinant virus of claim 27, wherein the PB1, PB2, PA, NP, and/or M have one or two substitutions relative to the corresponding polypeptide encoded by SEQ ID NOs:1-5.

29. The recombinant virus of claim 27, wherein the NA is N1.

30. The recombinant virus of claim 27, wherein the PB1, PB2, PA, NP, and M are encoded by SEQ ID NOs:1-5.

31. The plurality of vectors of claim 1, wherein the cDNAs for PB1, PB2, PA, NP, and/or M encode a polypeptide having one or two substitutions relative to the corresponding polypeptide encoded by SEQ ID NOs:1-5.

32. The plurality of vectors of claim 1, wherein the cDNAs for PB1, PB2, PA, NP, and M encode a polypeptide encoded by SEQ ID NOs:1-5.

* * * * *